PPIUCD Pre-insertion Screening

In preparation for insertion of the IUCD, confirm the following information about the woman and her clinical situation:

Ask the woman whether she still desires the IUCD for Immediate PPFP	❒ No	❒ Yes
Review her antenatal record and be certain that:		
• her antenatal screening shows that an IUCD is an appropriate method for her	❒ No	❒ Yes
• she has had FP counseling while not in active labor and there is evidence of consent in her chart **OR**	❒ No	❒ Yes
• she is being counseled in the post partum period	❒ No	❒ Yes
Review the course of her labor and delivery and ensure that none of the following conditions are present:		
If planning an *immediate post placental insertion*, check whether any of the following conditions are present:		
• Chorioamnionitis (during labor)	❒ Yes	❒ No
• More than 18 hours from rupture of membranes to delivery of baby	❒ Yes	❒ No
• Unresolved postpartum hemorrhage	❒ Yes	❒ No
If planning a *immediate postpartum insertion*, check whether any of the following conditions are present:		
• Puerperal sepsis	❒ Yes	❒ No
• Postpartum endometritis/metritis	❒ Yes	❒ No
• Continued excessive postpartum bleeding	❒ Yes	❒ No
• Extensive genital trauma where the repair would be disrupted by immediate postpartum placement of an IUCD	❒ Yes	❒ No
Confirm that sterile instruments are available*	❒ No	❒ Yes
Confirm that IUCDs are available and accessible on the labor ward*	❒ No	❒ Yes
	If **ANY** box is checked in this column, defer insertion of the IUCD and provide the woman with information about another method.	If **ALL** the boxes in this column are ticked, then proceed with IUCD insertion.

** If correct instruments or sterile IUCDs are not available, proceed with IUCD insertion if they become available within an appropriate time period.*

USAID FROM THE AMERICAN PEOPLE

NATIONAL RURAL HEALTH MISSION

MCHIP Maternal and Child Health Integrated Program