

PD-AAW-714  
53048

AGENCY FOR INTERNATIONAL DEVELOPMENT  
WASHINGTON, D.C. 20523

DATE: 9/10/87

MEMORANDUM

TO: AID/PPC/CDIE/DI, room 209 SA-18  
FROM: AID/SCI, Victoria Ose *VO*  
SUBJECT: Transmittal of AID/SCI Progress Report(s)

Attached for permanent retention/proper disposition is the following:

AID/SCI Progress Report No. 6. 233  
pd. Sept 1, 1986 - Jan 31, 1987  
rec'd 9/10

Attachment

*1 cy*

Semi- Annual progress Report

September 1, 1986 to January 31, 1987

Project No. 6.233 "Genetic Engineering of rhizobia for nodulation efficiency"  
Grant No. DPE-5542-G-SS-6039-00

Background.

Bradyrhizobium japonicum strain 61A76 is naturally composed of two distinct subpopulations of cells; one forms translucent and friable colonies on yeast extract mannitol gluconate medium (YEMG) while other forms slimy compact colonies on this medium. The former type breeds true on subculturing, while the latter produces both friable and slimy populations on subculturing. Regardless of the culture age, approximately 70% of the population in broth cultures of the slimy type consists of the friable true breeding variant.

Scope of the project and research done between September 1, 1986 and December 31, 87.

One of the objectives of this proposal was to understand the genetic basis of the mixed colony morphology in the slimy variant. In order to understand this phenomenon, we have constructed a gene bank of the strain 61A76 in the reconstructed cosmid vector pLAFRI. The original vector pLAFRI has a tetracyclin resistance gene as the marker. Our preliminary results indicated that strain 61A76 possessed high intrinsic resistance to tetracyclin. Therefore we replaced the tetracyclin marker gene in the vector with the Neomycin phosphoryltransferase gene (NPT) cloned from Tn5. The reconstructed vector possessed kanamycin resistance as the marker.

During this period we have also obtained variants of 61A76 that possess only the slimy compact phenotype. To our surprise two such variants tested failed to nodulate soybean cv. Williams. They also failed to deform root hairs. Experiments are underway to mate the gene bank into these variants and select the recombinants for mixed colony and Nod phenotype. These experiments will identify the clones that possess the genes present in the parent that are responsible for its mixed colony phenotype.

cy sent to PP/CDIE  
L. Fudenberg has copy

Rec'd in SO: SEP 10 1987