

PN-ABD-863

RESEARCH GUIDE
GUIA DE INVESTIGACION
GUIDE DE RECHERCHE

2392

CIP

CIP Research Guide 3

**TISSUE CULTURE
FOR
ELIMINATION OF PATHOGENS**

1989

Rolando Lizárraga, Nelson Espinoza,
Upali Jayasinghe, John H. Dodds



INTERNATIONAL POTATO CENTER (CIP)
CENTRO INTERNACIONAL DE LA PAPA (CIP)
CENTRE INTERNATIONAL DE LA POMME DE TERRE (CIP)

CIP Research Guide 3

**TISSUE CULTURE
FOR
ELIMINATION OF PATHOGENS**

1989

Rolando Lizárraga, Nelson Espinoza, Upali Jayasinghe, John H. Dodds

CIP
P.O. Box 5969
Lima, Perú

Location:
Av. La Universidad s/n
La Molina - Lima

Tel. 366920
Telex 25672 PE
Cable CIPAPA, Lima

CIP Research Guides (CRGs)

CIP Research Guides (CRGs) describe technologies that have been developed and used by CIP and National Potato Programs. The CRGs have been produced to promote exchange of information among scientists and are regularly updated to ensure that they describe the most recent advances.

Lizárraga, R.; Espinoza, N.; Jayasinghe, U.; Dodds, J. H. 1989. Tissue culture for elimination of pathogens. CIP Research Guide 3. International Potato Center, Lima, Peru. 21 pp.

**TISSUE CULTURE
FOR
ELIMINATION OF PATHOGENS**

- 1 Nature of pathogens
- 2 Thermotherapy
- 3 Chemotherapy
- 4 Surface sterilization
- 5 Antibiotics
- 6 Meristem isolation and culture
- 7 Maintenance of cultures
- 8 Media
- 9 Bibliography

Plant pathogens, such as nematodes, fungi, bacteria, and viruses, can be transmitted from diseased to healthy plants. However, not all cells become infected; the meristematic tissues are sometimes disease-free. It is possible to recover non-infected plants by in vitro meristem culture techniques and to grow them into healthy plants.

This document describes the methods that can be applied to remove pathogens from infected material and to produce "pathogen tested" plants for international distribution and propagation in a potato seed program.

1 NATURE OF PATHOGENS

Plant pathogens include nematodes, fungi, bacteria, rickettsia, mycoplasma, viruses and viroids. They are contagious and can be transmitted from diseased to healthy plants. In potatoes until now no reports indicate rickettsia as a pathogen.

The relative size of these pathogens varies greatly. Among the plant pathogens, nematodes are the largest and can be seen easily with a stereo microscope. Viruses and viroids are the smallest and one needs an electron microscope to visualize them.

The occurrence of a disease does not solely depend on the presence of a host and the pathogen. Environmental conditions especially humidity and temperature play an important role in the development of a disease. Thus a disease can be defined as a product of the interaction of the host, the pathogen and the environment. This concept of a disease is known as the "disease triangle".

The distribution of the different pathogens within a diseased plant also varies greatly. Pseudomonas solanacearum, potato leafroll virus, and mycoplasma are restricted to the vascular tissue of a plant. Erwinia carotovora, and potato virus X (PVX), invades both vascular and non-vascular plant tissues. Not all cells in a diseased plant are infected with pathogens. The meristematic tissues of root and shoots of an infected plant are sometimes free of viruses. In certain cases, such as PVX and tobacco rattle virus in potatoes, only the apical dome and the first young primordial leaf are free of viruses. The exact reason for this is not known, however, it is believed that one or all the following factors are responsible.

-
- High metabolic activity: Viruses replicate by taking over the host metabolic pathways. Due to the high metabolic activity in these cells viruses are unable to take over control of the host biosynthetic machinery.

 - Lack of vascular system: Viruses spread rapidly through the vascular system. Phloem restricted viruses (PLRV) cannot invade the meristematic tissues due to the absence of cell differentiation. In this meristematic region, viruses which infect non-vascular tissues spread from cell to cell through the plasmodesmata. This is a slow process which makes it relatively difficult for viruses to completely infect the rapidly dividing cells.

 - High auxin concentration: Plant meristematic tissues have a higher auxin concentration than tissues from the other plant regions. These auxins have been reported to inhibit the replication of viruses.

Since meristematic tissues are sometimes free of pathogens, it is possible to recover non-infected plants by in vitro meristem culture techniques and to grow them into healthy plants.

2 THERMOTHERAPY

Thermotherapy at elevated temperature (37 °C) does not eliminate the potato spindle tuber viroid (PSTV). PSTV consists of a single-stranded, ring-shaped RNA that is twisted in the form of a supercoil. In this form it is resistant to nucleases. Elevated temperature, far from decreasing the concentration of the viroid, favors its multiplication (Sanger and Ramm 1975). Therefore, a first test for PSTV should be carried out at the end of the thermotherapy period.

A method that permits the eradication of PSTV (Lizárraga et al. 1980) is based on the observation that in plants grown at low temperatures, the viroid concentration is low. In an experiment, plants were grown at 8°C for 4 months. Then apical domes were excised. 30 % of the plantlets regenerated were free of PSTV, even in the second tuber generation. A clear (negative) relationship between meristematic explant size and eradication success was observed (Lizárraga et al. 1982).

This method, however, is not suitable as a routine technique, since it is time consuming and costly. It may be useful in specific cases in which a valuable clone is heavily infected and no pathogen - tested material available.

Experiments carried out with different virus host systems have shown that treatment of plants with elevated temperatures (thermotherapy) leads to a reduction in virus concentration (titre) in the plant (Kassanis 1957; Quak 1977). Different reasons have been given to explain this phenomenon; probably not one alone, but a combination of several factors may be the cause for reduction in virus titre. These can include competition for sites of synthesis of nucleic acids and proteins between the fast dividing host cells and the virus particles, which may lead to a change in the

balance between synthesis and degradation of virus particles. Also the nucleic acid of the virus, the carrier of its genetic information, is usually protected from attacks by degrading enzymes because of a coat, made of many protein subunits. At elevated temperatures the linkage between these subunits becomes weaker, temporary holes may open and permit the attack of nucleases, leading to inactivation of virus and decrease of virus concentration.

Thermotherapy has been applied to dormant potato tubers. A reduction of virus concentration, mainly of potato leafroll virus (PLRV) has been observed. However, elimination was not achieved except for PLRV.

Thermotherapy applied to the whole plant as well as to sprouted tubers followed by meristem culture has been successfully used as a standard procedure for elimination of many viruses in potato (Stace-Smith and Mellor 1970; Pennazio and Redolfi 1973).

In the standard procedure used at CIP best results have been obtained when the plant is decapitated before introduction into thermotherapy and axillary buds grown while undergoing heat treatment. A daily temperature regime of 36 °C for 16 hours and 30 °C for 8 hours and continuous light of high intensity (10,000 lux) improved elimination rates. Plants are kept under these conditions for 4 weeks. From axillary buds as well as apical buds, meristems are isolated and cultivated, as shown in Section 6.

In vitro thermotherapy is an innovative method that can be used instead of the standard procedure. Nodal cuttings from in vitro plants are placed in plastic boxes with propagation medium C using agar as gelling agent. When the plantlets reach 3 to 4 cm height and have a good root system, they are ready for thermotherapy treatment (same as standard procedure). After 1 month treatment, apical meristems are isolated and cultured.

3 CHEMOTHERAPY

As an alternative to thermotherapy, chemotherapy has been tried in potato. A nucleoside analogue, Virazole, known for its broad spectrum against animal DNA and RNA viruses has shown variable results when applied to the potato plant as spray or in hydroponic culture followed by meristem culture. Preliminary results show some promise when culturing excised meristem tips in the presence of 100 ppm Virazole in the medium.

However, some reports indicate that antiviral chemicals may cause mutations in the plant. Therefore, at the present time, thermotherapy is definitely the preferred method of pre-treatment prior to meristem culture.

4 SURFACE STERILIZATION

If the surface of plant material is contaminated by pathogens or saprophytes, some of the contaminants quickly outgrow and can kill the plant introduced to in vitro. Most surface contaminants, such as bacteria and fungi, can be eliminated by surface sterilizing the plant material with a suitable sterilizing agent.

Surface sterilizing agents are normally applied for 10 - 15 minutes. Under aseptic conditions, the sterilizing solution is removed and the plant material washed 3 or 4 times for 5 minutes each time by agitation in sterile distilled water. Washing is important to remove excess sterilizing agent which inhibits plant growth.

Ethanol (Alcohol). Alcohol is a common surface sterilizing agent for killing bacteria and fungi, and is often given as a brief wash prior to applying other surface sterilizing treatments. It has a low surface tension and can easily penetrate between leaf hairs and wet the surface of the plant. 70 % Ethanol is a more effective surface sterilizing agent than 95-100 %.

Sodium or calcium hypochlorite. Plant material may also be surface sterilized with an aqueous solution of either sodium hypochlorite (NaOCl) or calcium hypochlorite (Ca[OCl]). The calcium salt is preferred as it is less phytotoxic. Many laboratories use a household bleach such as Clorox. These commercial products usually contain 5.25 % NaOCl as the active agent. When diluted with water (1 part bleach : 9 parts water), the final sterilizing solution should contain not less than 0.5 % NaOCl .

Because of complete dissociation, hypochlorite has relatively little activity at pH over 8.0, and it is much more effective by buffering the solution at about pH 6.0.

Freshly excised tissue, completely immersed in the hypochlorite solution, is surface sterilized after an exposure of 10-15 minutes. Following the hypochlorite treatment the plant material must be thoroughly rinsed with several changes of sterile distilled water to remove the disinfectant completely.

Mercuric chloride. Mercuric chloride (HgCl_2) has been used as a disinfectant, although it is extremely toxic. A solution of mercuric chloride is volatile at room temperature and can cause mercury poisoning. Therefore, we do not advise its use as a sterilizing agent !

Bactericides and fungicides. With material that is heavily contaminated some researchers recommend washing in a commercial bactericide/fungicide mix prior to surface sterilization. It should be remembered however that this treatment has no effect on systemic infections. Section 5 gives a list of bactericides and fungicides together with information about their use.

5 ANTIBIOTICS

Although employed routinely in animal cell cultures to prevent bacterial contamination, antibiotics have not been widely used in plant tissue cultures. Physiologists in the late 1950's were aware that these natural products may alter the growth and development of plant tissues cultures in vitro (Gautheret, 1959; Butenko, 1964). In fact, plant tissues are sensitive to antibiotics and show variable responses according to their genotype.

Most tissue culture specialists do not rely on antibiotics to remove superficial contaminants from explants. These compounds are expensive and no one is effective against all possible types of contaminating organisms. Antibiotics are only employed where microorganisms are difficult to eliminate by other means.

The following antibiotics or combinations of antibiotics have been successfully applied:

- Gentamycin
- Rifampicin
- Nystatin + Carbenecillin
- Gentamycin + Amphotericin B
- Vancomycin-HCl + Mycostatin
- Streptomycin + Carbenecillin

Toxicity has been reported to be caused in some cultures by:

- Penicillin
 - Streptomycin
 - Bactericin
 - Sparsomycin
-

6 MERISTEM ISOLATION AND CULTURE

The active growing point of the plant shoot is the meristem. It is a small region composed of rapidly dividing (meristematic) cells.

The dome of a shoot apical meristem contains the truly meristematic cells and is surrounded by leaf primordia and primary leaves. Since the more differentiated vascular tissues are found distant from the meristem (towards to older tissue of the stem), the vascular elements of the leaf primordia are still incipient, and have not yet made contact with the main strand of the vascular system in the stem. Therefore, virus particles, which may be present in the vascular system, can reach the meristematic region of the apex only through cell to cell movement; a slow process. This is one of the main reasons why in a virus infected plant, virus concentration decreases acropetally toward the meristem of both the apical and the axillary buds.

Isolation of the apical portion, called the meristem tip, under aseptic conditions and its culture on an adequate aseptic nutrient medium, leads to the development of plantlets. This developmental sequence, in principle, follows a pattern similar to that in the normal plant growth. The cells of the meristem divide and the differentiation of new tissues continues. The nutrition of the excised portion by the plant is supplied by the artificial medium. This technique, called meristem culture, was first applied for virus eradication some 30 years ago by Morel and Martin (1952) on Dahlia, and leads to pathogen-free plants.

The aseptic dissection of the meristem is a delicate progress and requires many hours of practice.

Stems are cut from the plant that has just undergone thermotherapy into segments each containing one node with its axillary bud. The leaves are carefully removed. The stem segments are disinfected for 30 seconds in 70 % alcohol, followed by 2.5 % calcium hypochlorite for 15 minutes. Then the stems are washed four times 5 minutes each with sterile distilled water to remove excess hypochlorite.

Under a binocular dissecting microscope, the leaflets surrounding the growing point are removed until only the apical dome and a few leaf primordia remain. The dome and two leaf primordia are excised and transferred to meristem culture medium A (section 8). The excised apical dome is transferred weekly to fresh medium B. After 6-8 weeks, the small plantlets are subcultured for further growth and micropropagation (Espinoza et al. 1989).

After regeneration from the cultured meristems, plants are tested (indexed) to detect any persisting virus infections. See the following references in Section 9:

- | | | |
|----------------------------------|---|-----------------|
| Fribourg, C., Nakashima, J. 1987 | - | Latex test |
| Salazar, L.F. 1981 | - | PSTV detection |
| Salazar, L.F. 1982 | - | Virus detection |
| Salazar, L.F. 1983 | - | ELISA |

7 MAINTENANCE OF CULTURES

In vitro material can be conserved in culture indefinitely when sufficient care is taken to avoid contamination, and transfer to fresh media is made at appropriate intervals. Pathogen-tested in vitro plantlets can be kept as base collections for propagation in a seed program.

For short-term maintenance, plants are grown in tubes on propagation medium C (section 8). The tubes are sealed with autoclavable plastic caps. These are better than cotton wool plugs that are easily penetrated by air-borne fungal spores and insects.

The growth rate of the plants depends on incubation temperature, medium composition, and genotype.

Long-term storage media E and H (section 8) exert an osmotic stress on the plantlets and can be used for storage at 25 °C; the stress reduces the growth rate and causes short internodes. Many nodes are thus available when propagation of the stored material is initiated. Material stored under these conditions normally needs to be transferred once a year only.

If the temperature can be lowered to 6 °C. This significantly reduces the growth rate of the plantlets and transfers are only required once every 2 or 3 years.

It is possible, therefore, to maintain a germplasm collection under the storage conditions outlined above. When a request is received for export of a particular pathogen tested genotype, it can be taken out from storage, micropropagated, and distributed in vitro.

8 MEDIA

All media used for this work are based on the salts of Murashige and Skoog (1962). The salt stock solutions are normally prepared in four separate parts:

- Salts
- $MgSO_4$
- Iron
- Vitamins

Salts stock solution: dissolve each salt in 200 cm³ distilled water

-	NH_4NO_3	35	g
-	KNO_3	40	g
-	$CaCl_2 \cdot 2H_2O$	9	g
-	KH_2PO_4	3.5	g
-	H_3BO_3	0.1	g
-	$MnSO_4 \cdot 4H_2O$	0.5	g
	($MnSO_4 \cdot H_2O$)	0.4	g)
-	$ZnSO_4 \cdot 7H_2O$	0.2	g
	($ZnSO_4 \cdot H_2O$)	0.1	g)
-	KI	0.02	g
-	$Na_2MoO_4 \cdot 2H_2O$	0.005	g

Dissolve 5 mg (0.005 g) of the following salts together in 10 cm³ of water:
add 1 cm³ of this solution to 200 cm³ water for the stock solution

- $CuSO_4 \cdot 5H_2O$ and $CoCl_2 \cdot 6H_2O$

Mix the ten salt solutions together to make 2 l (dm³) of the salt stock solution.

MgSO₄ stock solution

- MgSO₄.7H₂O 3.7 g in 100 cm³ distilled water

Iron stock solution

- Na₂EDTA 0.75 g
- FeSO₄.7H₂O 0.55 g

Dissolve FeSO₄.7H₂O in 20 cm³ distilled water: Na₂EDTA in 20 cm³ warm distilled water. Mix the solutions, cool, and make up to 100 cm³ with distilled water.

Vitamin stock solution

- Thiamine HCl (0.4 ppm) 20 mg
- Glycine (2.0 ppm) 100 mg
- Nicotinic acid (0.5 ppm) 25 mg
- Pyridoxine HCl (0.5 ppm) 25 mg

Dissolve each vitamin and complete to 500 cm³ with distilled water. Dispense 10 cm³ of stock solution in small vials and keep frozen. Use one vial for 1 liter of medium.

Medium preparation. Prepare 1 liter (dm^3) of the Murashige-Skoog basic medium by mixing the stock solutions with Inositol in the following proportions:

Salts	100 cm^3
MgSO ₄	10 cm^3
Iron	5 cm^3
Vitamins	10 cm^3
Inositol	100 mg

Add the relevant hormones and sucrose (see below), and autoclave the medium either with or without a gelling agent for 15 minutes.

Additions to the basic MS medium for propagation of specific media type::

Medium	MS media	+	the following additions
A	0.1	ppm	Gibberellic acid
	0.04	ppm	Kinetin
	2.5	%	Sucrose
	0.6	%	Agar
B	0.1	ppm	Gibberellic acid
	20.0	ppm	Putrescine HCl
	2.5	%	Sucrose
	0.6	%	Agar
C	1.0	ppm	Folic acid
	50.0	cm ³	Coconut milk
	4.0	ppm	L-Arginine HCl
	2.0	ppm	Calcium pantothenate
	10.0	ppm	Putrescine HCl
	3.0	%	Sucrose
	0.2	%	Gelrite or Phytigel
E	4.0	%	Sorbitol
	2.0	%	Sucrose
	0.8	%	Agar
H	4.0	%	Mannitol
	3.0	%	Sucrose
	0.8	%	Agar

9 **BIBLIOGRAPHY**

- Batenko, R.G. 1964. Plant tissue culture and plant morphogenesis. Translated from Russian. Jerusalem. Israel Program for Scientific Translations 1968.
- Espinoza, N.; Lizárraga, R.; Silva-Rodríguez, D.; Buitrón, F.; Bryan, J.; Dodds, J.H. 1989. Tissue culture micropropagation, conservation, and export of potato germplasm. CIP Research Guide 1. International Potato Center, Lima, Peru. 22 pp.
- Fribourg, C.; Nakashima, J. 1987. Latex test for detecting potato viruses. Slide Set, 32 slides; Guide Book, 12 pp. International Potato Center, Lima, Peru.
- Gautheret, R.J. 1959. La culture des tissus végétaux: Techniques et réalisations. Mason, Paris.
- Kassanis, B. 1957. The use of tissue culture to produce virus free clones from infected potato varieties. *Annals Applied Biology* 45:422-427.
- Lizárraga, R.E.; Salazar, L.F.; Roca, W.H.; Schilde-Rentscheler, L. 1980. Elimination of potato spindle tuber viroid by low temperature and meristem culture. *Phytopathology* 70:754-755.
- Lizárraga, R.E.; Salazar, L.F. 1982. Effect of meristem size on eradication of potato spindle tuber viroid. pp. 118-119. In: Hooker, W.J. Research for the potato in the year 2000. International Potato Center, Lima, Peru. 199 pp.
-

-
- Morel, G.M.; Martin, C. 1952. Guérison de dahlias atteints d'une maladie à virus. Comptes rendus hebdomadaire des séances de l'Académie des Sciences, Paris. 235:1324-1325.
- Murashige, T.; Skoog, F.C. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum* 15:473-497.
- Pennazio, S.; Redolfi, P. 1973. Factors affecting the culture in vitro of potato meristem tips. *Potato Research* 16:20-29.
- Quak, F. 1977. Meristem culture and virus free plants. pp. 598-615. In: Reinert, J.; Bajaj, Y.P.S. (Eds.) *Applied and fundamental aspects of plant cell, tissue and organ culture*. Springer, Berlin.
- Salazar, L.F. 1981. Detection of PSTV by gel electrophoresis. Slide Set. International Potato Center, Lima, Peru. 38 slides: Guide Book, 10 pp.
- Salazar, L.F. 1982. Virus detection in potato seed production. Technical Information Bulletin 18. International Potato Center, Lima, Peru. 14 pp.
- Salazar, L.F. 1983. Detection of potato viruses with ELISA. Slide Set. International Potato Center, Lima, Peru. 32 slides: Guide Book, 12 pp.
- Sanger, H.L.; Ramm, K. 1975. Radioactive labelling of viroid RNA. pp. 230-253. In: Markham et al. (Eds.) *Modification of the information content of plant cells*. Proceedings of 2nd John Innes Symposium, Norwich U.K.
- Stace-Smith, R.; Mellor, F.C. 1970. Eradication of potato spindle tuber virus by thermotherapy and axillary bud culture. *Phytopathology* 60: 1957-1958.
-

XVII-TR-E-09-0-500

Processed and printed by the Training and Communications Department, CIP
August 1989

Copies printed: 500
