Nontarget Effects of Soil-Applied Chlorpyrifos on Defoliating Pests and Arthropod Predators in Peanut[1]

J. E. Funderburk*, L.B. Braxton, and R. E. Lynch[2]

ABSTRACT

Studies determined the effects of granular chlorpyrifos on densities of nontarget arthropods in peanuts. A portion of several commercial fields was treated during the early pod stage of peanut growth in 1987 and 1988; then, densities of defoliating pests and arthropod predators were estimated weekly. Populations of *Helicoverpa* spp. were significantly increased in 1987 by chlorpyrifos application, but were not significantly affected in 1988. Densities of the velvetbean caterpillar, *Anticarsia gemmatalis* Hubner, were low on most sample dates. On dates when densities were great, populations were directly suppressed by chlorpyrifos application. Densities of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith), were always low, and were not affected by chlorpyrifos application. Arthropod predators sampled included the red imported fire ant, *Solenopsis invicta* Buren; spiders (Araneae: Araneidae); ground beetles (Coleoptera: Carabidae); earwigs (Dermaptera: Labiduridae); damsel bugs (Hemiptera: Nabidae); and bigeyed bugs (Hemiptera: Lygaeidae). Densities of spiders were significantly decreased by chlorpyrifos application. The other predators were not significantly affected. These results demonstrate that chlorpyrifos application results in nontarget effects on defoliating pest populations. Treated fields should be scouted regularly because such fields may have an increased potential for outbreaks of some defoliating pests.

Key Words: Insecta *Arachis hypogaea* L., *Helicoverpa* spp., *Spodoptera frugiperda* (J. E. Smith), *Anticarsia gemmatalis* Hubner, chlorpyrifos

Soil insect pests, primarily lesser cornstalk borer (*Elasmopalpus lignosellus* Zeller) and southern corn root worm (*Diabrotica undecimpunctata howardi* Barber), are key insect pests of peanut in the southeastern U.S. (7). Infestations that occur anytime between beginning pod and maturity [i.e. the R3 to R8 peanut growth stages (2)] can result in serious economic losses (6). Chlorpyrifos is efficacious against the soil-insect pest complex and is the most commonly used soil insecticide in peanut. Under normal growing conditions, chlorpyrifos applied during the R3 stage of peanut growth will provide residual efficacy until the R8 stage of peanut growth (5).

Soil-insect pests are effectively managed when insecticides such as chlorpyrifos are applied before populations reach outbreak levels. Because soil insecticides require rainfall or irrigation to be effective, control frequently does not occur rapidly enough to prevent unacceptable economic losses when applications are made after populations reach outbreak densities (1). Consequently, peanut producers sometimes apply chlorpyrifos as a preventative management practice.

[1]The research was supported in part by a Cooperative Agreement No. 58-43YK-7-0034 between ARS-USDA and the University of Florida. Marwan Kharbouth of Auburn University identified most of the natural enemies collected in this study. Fla. Agric. Exp. Sta. Journal Series No. 10,005.

[2]Associate Professor, N. Fla. Res. & Educ. Ctr., Quincy, Fla.; Field Development, Dow Chemical Co., Tallahassee, Fla.; and Research Leader, Insect Biol. & Popn. Mgmt. Lab. ARS-USDA, Tifton, Ga.

*Corresponding author.

Peanut fields in the southeastern U.S. are inhabited by a guild of leaf feeding species. Important species include the bollworm, *Helicoverpa zea* (Boddie); the tobacco budworm, *H. virescens* (Fabricius); the velvetbean caterpillar, *Anticarsia gemmatalis* Hubner; and the fall armyworm, *Spodoptera frugiperda* (J. E. Smith). Peanut fields also are inhabited by a guild of arthropod predators, including the red imported fire ant (*Solenopsis invicta* Buren), bigeyed bugs, damsel bugs, ground beetles, earwigs, and spiders. Many of these predatory species are important biological control agents of insect pests, preventing or ameliorating the severity of outbreaks.

The objective of this study was to determine the effects of soil-applied chlorpyrifos in peanut on densities of *Helicoverpa* spp., velvetbean caterpillar, and fall armyworm. The effects on densities of common predators in peanut fields also were evaluated.

Materials and Methods

Experiments were conducted in conventionally tilled and planted peanut fields at a 91-cm row spacing located in Jackson Co. in Florida. Experimental design was a randomized complete block with four fields in 1987 and two fields in 1988 serving as blocks. Row spacing in all fields was 91 cm. Peanut varieties each year were Florunner and NC 7. Standard agronomic practices were used for weed and disease control. Treatments consisted of an untreated control and an application of granular chlorpyrifos (Lorsban 15G, Dow Chemical Co., Midland, Mich. 48641) at the rate of 2.24 kg (AI)/ha in a 30 to 35 cm band over the row. Chlopyrifos was applied during the early pod fill stage of peanut growth (2) with a tractor-mounted granular applicator (Gandy Co., Owatonna, MN 55060) on 27 July in 1987 and on 6 or 7 July in 1988. A 2-ha treated and untreated area within each field was sampled, beginning soon after chlorpyrifos application and ending at about harvest. No other insecticide applications were made in the fields.

Densities of larval velvetbean caterpillars, fall armyworms, and *Helicoverpa* spp. and adult and immature spiders, damsel bugs, and bigeyed bugs were estimated by sweep sampling across a row (4) with a 38-cm-diam sweep net. Ten random samples of 10 sweeps each were taken weekly in treated and untreated areas of each field. Individual sweep samples were placed in plastic bags, put in an ice chest and kept cool during transport to the laboratory, and frozen until processed.

Densities of adult red imported fire ants, adult earwigs, and adult and larval ground beetles were estimated by pitfall sampling (3). Five, 15-cm-diam pitfall traps were randomly placed in treated and untreated areas of each field immediately following chlorpyrifos application. Peanut plants from 30 cm of row were removed at each sample location prior to trap placement, and a pitfall trap established directly within the row. Aluminum, 30-cm-diam shields were hung directly over pitfall traps to deflect water, and reservoirs of 100% ethylene glycol were maintained in inside bottoms to kill and preserve collected arthropods. Individual trap captures were collected bimonthly and returned to the laboratory for processing.

Effects each year of chlorpyrifos on densities of individual pests and natural enemies were determined by using analyses of variance. Data were analysed as a split-plot in time randomized complete block, with insecticide treatments as whole plots and dates as subplots (8). The main effect of chlorpyrifos treatment was a comparison of seasonal density between treated and untreated areas. The insecticide treatment*date interaction was used to determine if the response to chlorpyrifos was the same on all sample dates.

Results and Discussion

Densities of fall armyworm were low on all sample dates, with estimates greatest on the last three sample dates each



Fig. 1. Mean numbers (± SEM) of larval fall armyworms, *Helicoverpa* spp., and velvetbean caterpillars per 10 sweep samples in chlorpyrifos-treated and untreated areas of peanut fields sampled on nine dates in 1987 and 1988 in Jackson Co., Florida.

year (Fig. 1). Seasonal density was not affected by chlorpyrifos treatment in 1987 (F = 1.8; df = 1,3; $P > 0.05$) or 1988 (F = 1.0; df = 1,1; $P > 0.05$). The date*chlorpyrifos treatment interaction was not significant in 1987 (F = 1.6; df = 8,48; $P > 0.05$) or 1988 (F = 0.3; df = 8,16; $P > 0.05$).

Helicoverpa spp. were the most abundant defoliating pests in both 1987 and 1988 (Fig. 1). In 1987, more *Helicoverpa* were found in chlorpyrifos treated areas than in untreated areas on all individual sample dates (F = 11.5; df = 1,3; $P < 0.05$). Because densities were very low in both treatments on the last two sample dates, the date*chlorpyrifos treatment interaction also was significant (F = 2.4; df = 8,48; $P < 0.05$). Seasonal density of *Helicoverpa* spp. was statistically similar in treated and untreated areas of the peanut fields in 1988 (F = 13.2; df = 1,1; $P > 0.05$). The date*chlorpyrifos interaction also was not significant (F = 2.5; df = 8,16; $P > 0.05$).

Velvetbean caterpillar populations were very low in treated and untreated areas of each field in 1987, except during the last three sample dates (Fig. 1). Seasonal density was less in treated than untreated areas (F = 11.7; df = 1,3; $P < 0.05$), and the date*chlorpyrifos treatment interaction also was

2



Fig. 2. Mean numbers (± SEM) of damsel bugs, bigeyed bugs, and spiders per 10 sweep samples in chlorpyrifos-treated and untreated areas of peanut fields sampled on nine dates in 1987 and 1988 in Jackson Co., Florida.

significant (F = 15.1; df = 8,48; P < 0.001). Velvetbean caterpillar populations were very low on all sample dates in 1988. Seasonal density was similar in treated and untreated areas (F = 0.1; df = 1,1; P > 0.05), and the date°chlorpyrifos treatment interaction was not significant (F = 0.1; df = 8,16; P > 0.05).

Predators captured by sweepnets were damsel bugs, bigeyed bugs, and spiders (Fig.2). Damsel bug species in each peanut field were *Reduviolus roseipennis* Reuter, *Nabis capsiformis* Germar and *R. alternatus* (Parshley). The abundance of damsel bugs was not affected by treatment with chlorpyrifos. The main effect of chlorpyrifos treatment on damsel bugs was not significant in 1987 (F = 2.2; df = 1,3; P > 0.05) or 1988 (F = 1.0; df = 1,1; P > 0.05). The date°chlorpyrifos interaction also was not significant in 1987 (F = 0.9; df = 8,48; P > 0.05) or 1988 (F = 0.3; df = 8,16; P > 0.05). Bigeyed bug species in each peanut field were *Geocoris punctipes* (Say) and *G. uliginosis* (Say). The main effect of chlorpyrifos treatment was not significant in 1987 (F = 0.0; df = 1,3; P > 0.05) or 1988 (F = 0.1; df = 1,1; P > 0.05), and the date°chlorpyrifos treatment interaction was not significant in 1987 (F = 0.8; df = 8,48; P > 0.05) or 1988 (F

3



Fig. 3. Mean numbers (± SEM) of ground beetles, red imported fire ants, and earwigs per pitfall trap in chlorpyrifos-treated and untreated areas of peanut fields sampled on four dates in 1987 in Jackson Co., Florida.

= 0.5; df = 8,16; $P > 0.05$).

Abundance of spiders declined in plots treated with chlorpyrifos in 1987, but not 1988 (Fig. 2). In 1987, density was lower in treated than in untreated areas for several weeks following application, but was similar in treated and untreated areas for the remainder of the growing season. The main effect of chlorpyrifos treatment was not significant (F = 2.3; df = 1,3; $P > 0.05$), but the date*chlorpyrifos treatment interaction was significant (F = 2.6; df = 8,48; $P < 0.05$). The main effect of chlorpyrifos treatment was not significant in 1988 (F = 15.9; df = 1,1; $P > 0.05$), and the date*chlorpyrifos treatment interaction was not significant (F = 0.5; df = 8,16; $P > 0.05$).

The predators captured in pitfall traps were ground beetles, red imported fire ants, and earwigs. Frequent heavy rains invalidated results for all sample dates in 1988. Precision of density estimates in 1987, based on SEM's, was poor for red imported fire ants, ground beetles, and earwigs (Fig. 3). No significant differences were noted. Few red imported fire ants were collected on any sample date in plots treated with chlorpyrifos. However, the main effect of chlorpyrifos treatment only approached significance (F = 5.7; df = 1,3; P = 0.09). Nearly all of the earwigs in the pitfall samples were *Labidura riparia* (Pallas). The main effect of chlorpyrifos treatment was not significant for earwigs in 1987 (F = 3.4; df = 1,3; $P > 0.05$), despite much lower densities in treated than in untreated areas on all sample dates. Densities of ground beetles (*Calosoma* spp.) were equivalent in treated and untreated areas on each sample date, and the main effect of chlorpyrifos treatment was not significant (F = 0.2; df = 1,3; $P > 0.05$). The response of populations of red imported fire ants, earwigs, and ground beetles to chlorpyrifos application was the same on all sample dates (F = 1.1, 0.2, and 1.4, respectively; df = 3,18; $P > 0.05$).

Granular chlorpyrifos applied during the early pod stage of peanut growth for control of soil-insect pests resulted in direct and indirect effects on populations of defoliating pests. Velvetbean caterpillar populations declined in plots treated with chlorpyrifos. *Helicoverpa* spp. populations sometimes increased due to chlorpyrifos application. Increases in defoliator populations in treated areas probably resulted from indirect effects of chlorpyrifos on populations of natural enemies. Spider densities were reduced by chlorpyrifos treatment. Although results from pitfall sampling were inconclusive, the data indicate that red imported fire ants and possibly earwigs were directly reduced by chlorpyrifos application. However, other natural enemies were not affected, including damsel bugs, bigeyed bugs, and ground beetles. Our results demonstrate that chlorpyrifos for control of soil-insect pests will result in nontarget effects on defoliating pests. However, these effects can increase or decrease the possibility of an outbreak. Based on our findings, peanut fields should continue to be treated with soil insecticide when economically justified. However, treated fields should be scouted regularly, because such fields may have an increased potential for outbreaks of some defoliating pests.

Literature Cited

1. All, J.N., W. A. Gardner, E. F. Suber, and B. Rogers. 1982. Lesser cornstalk borer as a pest of corn and sorghum, pp. 36-46 *in* H. H. Tippins [ed.], A Review of Information on the Lesser Cornstalk Borer, *Elasmopalpus lignosellus* (Zeller). Georgia Agricultural Experiment Station Special Publication 17.
2. Boote, K.J. 1982. Growth stages of peanut (*Arachis hypogaea* L.) Peanut Sci. 9:35-40.
3. Jones, D., and M. H. Bass. 1979. Evaluation of pitfall traps for sampling lesser cornstalk borer larvae in peanuts. J. Econ. Entomol. 72:289-290.
4. Kogan, M., and H. N. Pitre, Jr. 1980. General sampling methods for above-ground populations of soybean arthropods, pp. 30-60. *in* M. Kogan and D. C. Herzog [eds.] Sampling Methods in Soybean Entomology. Springer, New York.
5. Mack, T. P., J. E. Funderburk, R. E. Lynch, M. G. Braxton, and C. B. Backman. 1989. Efficacy of chlorpyrifos in soil in Florunner peanut fields to the lesser cornstalk borer (Lepidoptera: Pyralidae). J. Econ. Entomol. 82:1224-1229.
6. Morgan, L. W., M. H. Bass, and H. Womack. 1982. Lesser cornstalk

- 4

borer as a pest of peanuts, pp. 56-64. *in* H. H. Tippins [ed.], A Review of Information on the Lesser Cornstalk Borer *Elasmopalpus lignosellus* (Zeller). Georgia Agricultural Experiment Station Special Publication 17.

Smith, J. W., Jr., and C. S. Barfield. 1982. Management of preharvest insects, pp. 250-325. *in* Peanut Science and Technology, H. E. Pattee and C. T. Young, [eds.]. Peanut Research and Education Society, Inc., Yoakum, Texas.

S. Steel, R. G.D., and J. H. Torrie. 1960. Principles and procedures of statistics with special reference to the biological sciences. McGraw-Hill, Inc., 481 pp.

Accepted December 17, 1990

A Morphological Appraisal of Wild Species in Section *Arachis* of Peanuts[1]

H. T. Stalker[2]

ABSTRACT

The cultivated peanut, *Arachis hypogaea* L., is a member of section *Arachis nom. nud.* along with its tetraploid progenitor, *A. monticola* Krap. et Rig., four validly described diploid species, eight diploid species whose names have never been validly published, and a large collection of taxa discovered since 1975. Systematic relationships and possible species circumscriptors are assessed in section *Arachis* by means of numerical taxonomy. Seventy-three accessions were grown in the field and three randomly selected specimens of each accession were evaluated. Numerical techniques in the form of cluster and principal components analyses were used on 56 characters, including 20 reproductive, 30 vegetative, and six created variables. Most variation was observed for leaflet size and shape, followed by branching habits and flower size. Although grouping of accessions did not always conform to expectations based on published species descriptions, general relationships among taxa are evident from the analyses. A total of 20 to 25 species will likely exist in section *Arachis* based on the observed variability.

Key Words: *Arachis*, species, numerical taxonomy.

The genus *Arachis* comprises a large and diverse group of diploid and tetraploid taxa native to South America. Species were first described by Linneaus in 1753 and monographs have since been published by Chevalier (1), Hoehne (7), and Hermann (6). Due to inadequate herbarium specimens and the lack of field collections from over the range of the genus, Gregory *et al.* (4) considered these works to be greatly deficient. Three different infrageneric classifications have also been published (5, 8, 9) wherein the genus was divided into sections. However, none of the sectional names have been validly published according to the International Code of Botanical Nomenclature (10). Although 70 or more species may exist in the genus (3, 18), only 22 species, one variety and one hybrid have been validly described. Names for at least 12 additional species are commonly found in the literature, but these names have never been validated with descriptors. Collections made in South America during recent years have nearly tripled the number of accessions maintained under cultivation (12), and the taxonomy of *Arachis* has become increasingly deficient and confusing.

Species of section *Arachis nom. nud. sensu* Gregory *et al.* (5) are of special interest because the cultivated peanut *A. hypogaea* L. ($2n = 4x = 40$) is a member of the group. Both annual and perennial species in this section are widely distributed east of the Andes and north of 35° south latitude in South America. The known distribution of section *Arachis* has been extended to more than 1000 km northwest in Bolivia, 1500 km to the northeast in Brazil, and 350 km north along the Atlantic coast since 1976 (11). Section *Arachis* is distinguished from other sections of the genus by having tap roots but no adventitious, roots or rhizomes, vertical pegs which do not grow in a horizontal direction after penetrating the soil, and flowers without prominent red veins (5, 10). Six species (*A. batizocoi* Krap. et Greg., *A. diogoi* Hoehne, *A. helodes* Martius ex Krap. et Rig., *A. hypogaea*, *A. monticola* Krap. et Rig. and *A. villosa* Benth. var. *villosa*) have been described and names for nine other species have been used (Table 1). Estimates of the number of species which eventually will be described in section *Arachis* have ranged from 15 to 20 by Stalker and Moss (18) and from 34 to 46 by Valls *et al.* (19).

The purpose of this paper is to present results from a morphological study of wild *Arachis* species referable to section *Arachis*. A large number of living accessions have been used in an attempt to better describe taxonomic relationships in section *Arachis* and to delimit groups which may eventually be recognized as species. It is hoped that the results from this analysis will lay the foundation for a more complete taxonomic synthesis of section *Arachis* in the future.

Materials and Methods

Seventy-three wild peanut accessions belonging to section *Arachis* were transplanted in a field at the Sandhills Agricultural Research Station, NC in May 1986 (Table 2). Forty-three of these accessions were obtained from D. J. Banks, U.S. Curator at that time for *Arachis* species at Stillwater, OK in 1979 and 1981; 14 accessions were obtained in 1982 from C. E. Simpson of Texas A&M University, Stephenville, TX; and 14 accessions plus a small leaf (sl) mutant of 10035 were already maintained at North Carolina State University prior to recent introductions. Two additional collections (6536 and 7762) were obtained in Brazil during a collection expedition in 1984 by H. T. Stalker, C. E. Simpson, J. F. M. Valls, I. J. Godoy, and W. L. Werneck.

Three plants were randomly selected from each accession grown in the field and scored for 20 reproductive and 30 vegetative traits 90 to 95 days after planting. Data for the following traits were collected: presence or absence of flowers on the main stem; main stem and lateral branches erect or prostrate; length of main stem; length of longest lateral branch; relative amount of lateral branching; annual or perennial; length and width of main stem and lateral leaves and leaflets; shape of main stem and lateral leaflets (lanceolate = 1 to obcordate = 6); shape of the leaflet apex on main stem and lateral leaves (acuminate-abrupt = 1 to emarginate = 7); stipule length; petiole length of main stem and lateral leaves; internode length between leaflets on main stem and lateral leaves; numbers of trichomes on stems, upper and lower leaflet surface, leaflet margins and pegs; length and type

[1]Paper No. 12576 of the journal series of the North Carolina Agricultural Research Service, Raleigh, NC 27695-7643. This publication was supported in part by the Peanut CRSP, USAID grant number DAN-4048-G-SS-2065-00. Recommendations do not represent an official position or policy of USAID.

[2]Professor of Crop Science, North Carolina State University, Raleigh, NC 27695.