

## Managing native and legume-fixed nitrogen in lowland rice-based cropping systems\*

T. GEORGE<sup>1,2</sup>, J.K. LADHA<sup>1</sup>, R.J. BURESH<sup>1,3</sup> and D.P. GARRITY<sup>1</sup>

<sup>1</sup>International Rice Research Institute (IRRI), P.O. Box 933, Manila, Philippines,

<sup>2</sup>Nitrogen Fixation by Tropical Agricultural Legumes (NifTAL) Project, University of Hawaii, 1000 Holomua Avenue, Paia, HI 96779, USA and

<sup>3</sup>International Fertilizer Development Center (IFDC), P.O. Box 2040, Muscle Shoals, AL 35662, USA

**Key words:** biological nitrogen fixation, denitrification, fallow, flooded soil, leaching, legume, nitrate, nitrogen balance, nitrogen loss, *Oryza sativa*, rice, weeds

### Table of contents

- Abstract
- Introduction
- Major lowland rice-based cropping systems
  - Year-round lowland rice culture
  - Lowland rice culture with short non-flooded season
  - Lowland rice culture with long non-flooded season
- Soil nitrogen dynamics in rice lowlands
  - Soil nitrate accumulation
  - Loss of nitrate from soil
  - Nitrogen dynamics of major lowland rice-based cropping systems
  - Impacts of nitrate loss from rice lowlands
- Legumes in lowland rice-based cropping systems
- Biological nitrogen fixation by legumes in rice lowlands
  - Rhizobial determinants
  - Plant determinants
  - Environmental determinants
- Nitrogen contribution from legumes in rice lowlands
  - Food and forage legumes
  - Stem-nodulating aquatic green manure legumes
  - Legumes in the post-rice as opposed to the pre-rice niche
- Maximizing total nitrogen capture by legumes in rice lowlands
  - Managing soil nitrate supply
    - Allowing for increased nitrate uptake
    - Minimizing soil supply of nitrate
  - Managing inefficient nitrate uptake by legumes

\* IRRI-NifTAL-IFDC joint contribution.

Use of legume nitrogen for lowland rice

Belowground residues

Aboveground residues

Nitrogen release from legume residues

Conclusions and research directions

References

**Abstract**

Lowlands comprise 87% of the 145 M ha of world rice area. Lowland rice-based cropping systems are characterized by soil flooding during most of the rice growing season. Rainfall distribution, availability of irrigation water and prevailing temperatures determine when rice or other crops are grown. Nitrogen is the most required nutrient in lowland rice-based cropping systems. Reducing fertilizer N use in these cropping systems, while maintaining or enhancing crop output, is desirable from both environmental and economic perspectives. This may be possible by producing N on the land through legume biological nitrogen fixation (BNF), minimizing soil N losses, and by improved recycling of N through plant residues. At the end of a flooded rice crop, organic- and  $\text{NH}_4\text{-N}$  dominate in the soil, with negligible amounts of  $\text{NO}_3$ . Subsequent drying of the soil favors aerobic N transformations. Organic N mineralizes to  $\text{NH}_4$ , which is rapidly nitrified into  $\text{NO}_3$ . As a result,  $\text{NO}_3$  accumulates in soil during the aerobic phase. Recent evidence indicates that large amounts of accumulated soil  $\text{NO}_3$  may be lost from rice lowlands upon the flooding of aerobic soil for rice production. Plant uptake during the aerobic phase can conserve soil  $\text{NO}_3$  from potential loss. Legumes grown during the aerobic phase additionally capture atmospheric N through BNF. The length of the nonflooded season, water availability, soil properties, and prevailing temperatures determine when and where legumes are, or can be, grown. The amount of N derived by legumes through BNF depends on the interaction of microbial, plant, and environmental determinants. Suitable legumes for lowland rice soils are those that can deplete soil  $\text{NO}_3$  while deriving large amounts of N through BNF. Reducing soil N supply to the legume by suitable soil and crop management can increase BNF. Much of the N in legume biomass might be removed from the land in an economic crop produce. As biomass is removed, the likelihood of obtaining a positive soil N balance diminishes. Nonetheless, use of legumes rather than non-legumes is likely to contribute higher quantities of N to a subsequent rice crop. A whole-system approach to N management will be necessary to capture and effectively use soil and atmospheric sources of N in the lowland rice ecosystem.

**Introduction**

Rice (*Oryza sativa*) is the major staple food crop of the tropical world and rice-based cropping patterns are prevalent in much of the tropics. Rice is grown on 145 M ha, 87% of which is in wetland (lowland) culture, with the remaining grown on dry lands. The annual cultivation period for rice is determined by the distribution of rainfall, availability of irrigation water, and temperatures. Upland crops are typically grown during the nonflooded period, particularly in the transition seasons either at the end of the rains or before the onset of heavy rainfall. Cool-season crops, such as wheat and winter legumes,

are grown during the nonflooded season in areas where cool temperatures limit rice production.

Lowland rice is grown in banded fields and is surface-flooded during most of the growing season. Rice soils are typically flooded and puddled, and rice seedlings are then transplanted. Alternatively, germinated seeds are broadcast onto flooded soil that is either puddled or non-puddled. Dry seeding, where seeds are dibbled or broadcast onto aerobic soil, is also practiced. The soil is flooded when there is sufficient water to be impounded.

Nitrogen is the input required in largest quantities for lowland rice production. Soil N and biological nitrogen fixation (BNF) by associated

organisms are major sources of N for lowland rice. More than 50% of the N used by flooded rice receiving fertilizer N is derived from the combination of mineralization of soil organic N (Bouldin, 1986; Broadbent, 1984) and BNF by free-living and rice plant-associated bacteria (Roger and Ladha, 1992). The remaining N requirement is normally met with fertilizer. However, the efficiency of fertilizer N uptake by rice is low due to large N losses from flooded soil (De Datta and Buresh, 1989; De Datta and Patrick, 1986; Savant et al., 1982). Further, declining or stagnating yields have recently been noticed with flooded rice culture receiving fertilizer N (Flinn and De Datta, 1984). An additional concern is that the capacity of soil to supply N may decline with continuous intensive rice cropping necessitating the application of successively larger quantities of fertilizer N to maintain yields (Bacon, 1990). On the other hand, large amounts of soil  $\text{NO}_3$  may be lost from rice lowlands upon the flooding of aerobic soil before the establishment of rice (Buresh et al., 1989).

Reducing N fertilizer use while maintaining the native soil N resource and enhancing crop N output is desirable from both environmental and economic perspectives. This may be possible by obtaining more N on the land through biological nitrogen fixation (BNF), reducing losses of N, and by recycling the N captured in vegetation during the off-season. Buresh and De Datta (1991) recently reviewed the dynamics and management of N in rice-legume cropping systems. Their review indicates that soil N losses may be reduced by fallow management, and legumes can partially contribute to the N requirement of a succeeding rice crop. Thus, the management of indigenous soil N and N derived in situ through legume BNF poses potentials for enhancing the N nutrition and N use efficiency of crops and the total N output from a lowland rice-based cropping system. The integrated application of possible N management tactics will require detailed knowledge of the N dynamics of the cropping system as influenced by variation in hydrological characteristics and diversity of cropping practices.

In this paper, we address the N dynamics for a range of major tropical rice-growing ecosystems. We attempt to define the major lowland rice-

based cropping systems, and to document their hydrological, climatic and management effects on soil N dynamics. We then discuss the use of legumes in the capture and use of indigenous soil nitrogen and fixed atmospheric nitrogen in major rice-based cropping systems.

### Major lowland rice-based cropping systems

Rice-based cropping patterns are prevalent in those tropical areas with 1500 mm or more annual rainfall with at least three consecutive months of 200 mm per month. The amount and duration of rainfall normally determine the number of crops possible. The rates of onset and decline of rainfall determine the type of rice culture, other crops in the pattern, and difficulty in tillage (Harwood and Price, 1976). Growing upland crops after paddy rice requires soil conversion from an anaerobic puddled state to an aggregated aerobic condition. This is often difficult in soils high in montmorillonite clays. These soils are often left fallow during the dry months even when irrigation water is available (Palaniappan, 1985).

The number of rice crops grown in lowlands varies with supplemental water availability and prevailing temperatures. The type of upland crops that can be grown are influenced by the durations and timing of non-flooded periods. For purposes of this discussion, we distinguish three broad classes of tropical lowland rice culture in terms of the annual period in which the soil is not flooded: (1) continuously flooded or saturated year-round (Fig. 1A), (2) short non-flooded (Fig. 1B), and (3) long non-flooded (Fig. 1C). The three classes may be further subdivided on the basis of other important determinants of N dynamics and upland crop adaptation, such as temperature and water availability.

#### *Year-round lowland rice culture*

Continuous rice production systems were rare prior to the introduction of photoperiod-insensitive rice cultivars. However, they have become increasingly common. At the most intensive limit, three rice crops are successively cultivated




Fig. 1. Major lowland rice-based cropping systems from a soil nitrogen dynamics perspective. (A) permanent flooding (or soil saturation), (B) short, non-flood period, (C) long, non-flood period; (i) warm tropics and (ii) cool tropics/subtropics.

per annum under full irrigation, and the fields remain constantly saturated or flooded. The total area is not large, but this system is locally prominent in such areas as north Java and Bali, Indonesia.

Much more important are the irrigated

double-crop ricelands, which are very common in countries at latitudes within 14° of the equator. Huke (1982) estimated double-cropped ricelands at 8.8 M ha or 13% of the total irrigated rice area worldwide. Much of the double-cropped land remains flooded or saturated virtu-

ally year-round. An example is the double-cropped pump irrigated areas in Central Luzon, Philippines. Some very poorly drained rainfed and deepwater ricelands also experience year-round waterlogging.

*Lowland rice culture with short non-flooded season*

In much of Asia's irrigated double-cropped rice-land there is a short dry period of one to two months between rice crops. This generally occurs in the late wet season before the dry-season crop is established, and/or after the dry-season crop is harvested.

Some rainfed areas that experience long wet seasons support two successive rice crops, leaving a short dry soil period of two to three months. The fields are usually left fallow due to lack of water or difficulty in tillage. In some areas legumes are produced on residual soil moisture, or established as relay crops on supplemental irrigation. Examples of this cropping system are rice-rice-grain legumes in southern India and Philippines, rice-rice-soybeans in Indonesia and rice-rice-*Astragalus* sp. in China.

*Lowland rice culture with long non-flooded season*

Single-crop rice production is the dominant rice cultural system. The soil remains non-flooded from four to eight months of the year. This is the dominant case in the tropics because most rice-land is rainfed, or can be irrigated only in the wet season (Garrity et al., 1986). It is also dominant in the subtropics because although the land is fully irrigated, rice is constrained by cool temperatures during the winter season. Temperature also affects the soil N dynamics. Therefore, we recognize two subcategories of single-crop lands, those in the warm tropics, and those with cool winters.

In single-crop ricelands, the rice crop may occupy the field from 90 to 210 days. There is great diversity in the rainfed rice ecosystems. They range from favorable rainfed conditions, to drought or submergence-prone shallow (0 to 25 cm), medium deep (25 to 50 cm), or deep water (>50 cm) flooding regimes. The highly

variable surface hydrology has a profound influence on the nature of the N dynamics. Particularly significant is the unpredictable occurrence of aerobic or dry soil surface conditions during the growing season, alternating with surface flooding.

During the off-season, rainfed ricelands are typically fallowed (De Datta, 1981). The straw and fallow weed vegetation are subject to grazing by farm livestock. In a minor fraction of the area with conducive residual soil water-holding capacity, and/or a high groundwater table, upland crops, including legumes, are grown in the post-rice season. This practice is most common where the soil textures are loamy and exhibit superior tillage properties.

In well-drained ricelands with a gradual onset of rainfall in the dry-to-wet transition period, upland crops are grown prior to rice. Very early duration is advantageous, to permit maturity before the soil waterlogs. Mungbean is a very common grain legume in the pre-rice niche.

At latitudes above 14°N rice may experience significant cool-temperature stress during the coolest periods of the year. The establishment of dry-season rice crops is deliberately delayed to avoid this damage. Above 20° N dry-season rice cultivation is generally not feasible due to cold stress. Winter cereals or legumes are often grown in rotation with rice. The rice-wheat cropping pattern is dominant, occupying some 18 M ha in South Asia and China. Rice-legume patterns are also common. These often include chickpea, lentil, or soybean.

**Soil nitrogen dynamics in rice lowlands**

Fertilizer N transformations in flooded rice fields have been researched extensively (De Datta and Buresh, 1989; De Datta and Patrick, 1986). However, little attention has been paid to N transformations during the dry season when the soil is usually aerated (Buresh and De Datta, 1991). Knowledge is particularly lacking on the dynamics of soil N during the transition periods between dry and wet seasons.

A conceptualization of the dynamics of soil mineral N in a lowland, rice-growing soil undergoing a flooded-nonflooded sequence is pre-




Fig. 2. Dynamics of soil mineral nitrogen in a lowland rice soil.

sented in Figure 2. Under saturated or flooded conditions, nitrification is restricted by a limited  $O_2$  supply and  $NH_4$  accumulates in the soil. At the end of a flooded rice crop, organic N and  $NH_4$ -N dominate and  $NO_3$  present is typically negligible. Subsequent drying of the soil favors aerobic N transformations. Organic N mineralizes to  $NH_4$ , which is rapidly nitrified into  $NO_3$ . As a result,  $NO_3$  accumulates in the soil during the aerobic phase, but is rapidly lost upon soil flooding.

Plant growth can influence the magnitudes and forms of soil mineral N (Fig. 3). Nitrate, the




Fig. 3. Conceptual role of plants in conserving soil mineral and atmospheric nitrogen in a lowland rice-based cropping system.

predominant form of soil mineral N during the nonflooded season, will be assimilated by plants growing during the dry season and dry-to-wet transition periods. Buresh et al. (1989) found that soil  $NO_3$  during the dry season in a lowland rice soil correlated inversely with the amount of N accumulated in weeds. Legumes grown during this season additionally accumulate atmospheric N through BNF. Part or all of the N in the plant biomass is removed from the land through the harvestable products of an economic crop. The remaining plant N can be returned to the soil as residues.

Decomposition of plowed-under dry-season plant residues releases  $NH_4$  into the flooded soil. Rice uptake decreases  $NH_4$ , the predominant N form in the flooded soil. Declining  $NH_4$  level in flooded soil during the later part of rice season is partly replenished through BNF in flooded soil (Eskew et al., 1981; Yoshida and Yoneyama, 1980) and from decomposition of rice roots and stubbles in addition to mineralization of soil N.

#### Soil nitrate accumulation

Nitrate-N accumulates when lowland rice soils become dry and aerated (Patrick and Wyatt, 1964; Ventura and Watanabe, 1978). Nitrate-N levels prior to flooding for wet season rice in several Philippine lowland rice soils ranged from 5 to  $39 \text{ mg kg}^{-1}$  soil (Ponnamperuma, 1985). Buresh et al. (1989) measured 39 to  $91 \text{ kg soil } NO_3\text{-N ha}^{-1}$  in the top 60 cm layer at the end of the dry season at three Philippine sites.

The production of  $NO_3$  depends on the amount of organic N available for mineralization. Organic N varies considerably in lowland rice soils, depending on the level of native organic matter, and the history of cropping and organic amendments. The amount of N mineralized is increased by higher soil organic N levels (Broadbent, 1984; Hadas et al., 1986; Sahrawat, 1983). Bacon (1990) reported that successive soil incorporation of rice stubble gradually increased the N mineralization potential.

Residual soil water from the preceding wet-season rice crop, intermittent rains, and irrigation cause fluctuating soil water conditions. Soil water fluctuation generally results in a large production of  $NO_3$  (Birch, 1960; Herlihy, 1979;

Sahrawat, 1980; Seneviratne and Wild, 1985). In a greenhouse study, continuous soil flooding during the dry season prevented  $\text{NO}_3$  accumulation (Ventura and Watanabe, 1978). Drying, as well as alternate drying and wetting treatments resulted in the accumulation of soil  $\text{NO}_3$ .

Dry-season fallows are subjected to tillage for upland crop establishment or for weed control. Tillage increases soil aeration and tends to enhance the nitrification of mineralized  $\text{NH}_4$  into  $\text{NO}_3$  (Dowdell et al., 1983). Herridge (1986) reported more  $\text{NO}_3$  in a cultivated fallow compared with a non-tilled fallow in a rainfed environment in Australia.

#### *Loss of nitrate from soil*

Flooding of aerated soil has generally been accepted as a major factor of  $\text{NO}_3$  loss from rice soils (Bacon et al., 1986; Buresh et al., 1989; Janssen and Metzger, 1928; Patrick and Wyatt, 1964; Strickland, 1969; Wallihan, 1938; Yamane, 1957). With the use of  $^{15}\text{N}$  microplots, Bacon et al. (1986) reported 90% reduction in  $\text{NO}_3$  concentration after flooding of a rice soil. Loss of up to  $91 \text{ kg ha}^{-1}$  of native  $\text{NO}_3\text{-N}$  after flooding of fallow rice fields in the Philippines has been reported by Buresh et al. (1989).

Soil flooding can promote leaching losses of  $\text{NO}_3$ . Downward water flow carries  $\text{NO}_3$  to lower soil layers (White, 1988). Two direct determinants of  $\text{NO}_3$  leaching are the net downward flux of water and the concentration of  $\text{NO}_3$  in the leaching water. Measuring these variables presents considerable problems because of difficulty in sampling and spatial and temporal variability in water flux and  $\text{NO}_3$  concentration (Keeney, 1986; White, 1988). Large  $\text{NO}_3$  losses through leaching occur generally from aggregated soils and from soils under bare fallow (Legg and Meisinger, 1982; White, 1988). These conditions exist in lowland soils during the aerobic phase before puddling for rice culture.

Soil saturation depletes soil of its  $\text{O}_2$ , thus creating favorable conditions for denitrifying organisms to reduce  $\text{NO}_3$ . Denitrification results in the production of  $\text{N}_2$  and  $\text{N}_2\text{O}$  (Knowles, 1982). The rates of  $\text{N}_2$  and  $\text{N}_2\text{O}$  evolution from soils will be higher under conditions of fluctuating redox potentials (wetting and drying cycles) than

when the redox potential is continuously high (aerobic condition) or continuously low (anaerobic condition) (Letey et al., 1981; Reddy and Patrick, 1975). Mosier et al. (1986) measured large fluxes of  $\text{N}_2$  and  $\text{N}_2\text{O}$  from both fallow and cropped aerobic soils following water applications. Conditions favorable for  $\text{N}_2$  and  $\text{N}_2\text{O}$  emissions may occur during the dry-to-wet transition period in lowland rice-based cropping systems.

Bacon et al. (1986) attributed rice soil  $\text{NO}_3$  loss primarily to denitrification. An incubation study with the use of  $^{15}\text{N}$ -labeled  $\text{NO}_3$  by Buresh et al. (1989) indicated that  $\text{NO}_3$  completely disappeared after 9 days of flooding, and 5% or less of the added  $\text{NO}_3$  remained in the soil as either  $\text{NH}_4$  or organic N. Since conversion of  $\text{NO}_3$  to  $\text{NH}_4$  was negligible, either denitrification or leaching appeared to be the major loss mechanism.

#### *Nitrogen dynamics of major lowland rice-based cropping systems*

Soil N is essentially conserved under year-round flooded rice culture. Maintaining flooding between and during rice crops increases the soil N supply (Ponnamperuma, 1985; Santiago-Ventura et al., 1986; Ventura and Watanabe, 1978). Nitrate does not accumulate in the soil since soil drying between rice crops is minimal. Therefore, soil  $\text{NO}_3$  losses associated with flooding events are not experienced. Rice field-associated BNF contributes to the maintenance of available soil N in these soils (Koyama and App, 1979).

In warm, tropical single-crop rice lowlands, large quantities of  $\text{NO}_3$  might accumulate in soil over the prolonged aerated soil phase, then be lost during the dry-to-wet transition period. The quantity of  $\text{NO}_3$  accumulated will depend on how the fallow is managed. Tillage and occasional rains might increase soil  $\text{NO}_3$ .

In intensively cultivated rice-wheat lowlands, fertilizer N application in excess of wheat crop requirements may lead to large quantities of residual mineral N after wheat harvest. Chaney (1990) observed significant increases in residual soil  $\text{NO}_3$  levels after winter wheat at several sites in England when the N applications rates ex-

1

ceeded the optimum. Soil  $\text{NO}_3$  after wheat in a sandy loam soil in India averaged  $82 \text{ kg N ha}^{-1}$  (Sharma et al., 1985), but the portion originated from fertilizer might have been low (Macdonald et al., 1989). Sharma et al. (1985) also found that  $\text{NO}_3$  accumulation during a mungbean crop following wheat was substantial. Prior to the start of the wet season, the soil contained  $223 \text{ kg NO}_3\text{-N ha}^{-1}$ .

#### *Impacts of nitrate loss from rice lowlands*

It is well understood that the loss of  $\text{NO}_3$  via leaching and denitrification has undesirable environmental consequences. Leaching leads to  $\text{NO}_3$  pollution of ground-water and  $\text{N}_2\text{O}$  emission from denitrification poses a threat to the stability of the ozone layer (Pretty and Conway, 1989). The significance, if any, of  $\text{NO}_3$  loss from rice lowlands on the environment, and the negative impacts of  $\text{NO}_3$  loss on the long-term N fertility of lowland rice soil, have thus far been largely ignored by researchers.

Available reports suggest possible high losses of  $\text{NO}_3$  during soil flooding for rice production, but do not account for the cumulative loss that can occur during the entire dry and dry-to-wet transition periods. Lowland rice soils undergo short-term wetting or flooding, especially from intermittent early rain near the end of the dry season. During short-term wetting of the soil,  $\text{NO}_3$  can be lost through denitrification (Mosier et al., 1986). Heavy rain would cause substantial downward water flow, especially in sandy soils, leading to  $\text{NO}_3$  loss through leaching (White, 1988). It is likely that the total  $\text{NO}_3$  loss is greater than the reported disappearances of  $\text{NO}_3$  after soil flooding for rice production.

#### **Legumes in lowland rice-based cropping systems**

Legumes are grown in rice lowlands primarily for food, and to a lesser extent for green manure and forage. The importance of legumes in rice lowlands is enhanced by their ability to meet part of their N requirements from the atmosphere

through BNF and their potential to enhance soil N fertility. The major food legumes grown include cowpea (*Vigna unguiculata*), mungbean (*Vigna radiata*), peanut (*Arachis hypogaea*), pigeon pea (*Cajanus cajan*), chick pea (*Cicer arietinum*), and soybean (*Glycine max*). Major green manure legumes include milk vetch (*Astragalus sinicus*) and *Sesbania* spp. in China, *Sesbania* spp., sunnhemp (*Crotalaria juncea*) and berseem clover (*Trifolium alexandrinum*) in India; and indigo (*Indigofera tinctoria*) in the Philippines (Garrity and Flinn, 1988; Singh et al., 1991; Yost and Evans, 1988). The most promising forage legumes include siratro (*Macroptilium atropurpureum*), lablab bean (*Lablab purpureus*), sunnhemp, and clitoria (*Clitoria ternatea*) (Carangal et al., 1988). Several of the green manure legumes such as berseem clover (*T. alexandrinum*) are also suitable as forages.

The length of the non-flooded season, water availability, soil properties and climate primarily determine when and where legumes can be grown. These together with socio-economic factors determine whether food, forage, or green manure legumes are grown on ricelands. Under continuous rice culture, the periods between rice crops are too short to include legumes. In lowlands where two rice crops are taken, the non-flooded fallow may be long enough to grow a short-duration legume. Green manure legumes could be planted during a 40-to-60 day transition period between rice crops (Garrity and Flinn, 1988). Relay cropping of pulse crops such as mungbean after the second rice crop is common in parts of India (Palaniappan, 1985).

In single rice crop lowlands of the warm tropics with a long nonflooded season, there are two niches for short-duration legumes. They are the post-rice dry season for grain or forage types and the pre-rice dry-wet transition for grain or green manure types (Garrity and Flinn, 1988). Long-duration forage, green manure and cover crop legumes are grown sometimes as well. They may occupy most of the non-flooded season. In single rice crop lowlands of the cool tropics/subtropics, short- and long-duration winter legumes are grown for food, forage or green manure.

**Biological nitrogen fixation by legumes in rice lowlands**

The determinants of BNF by legumes have been dealt with in detail in other recent papers (Bohlool et al., 1992; Peoples and Craswell, 1992; Peoples and Herridge, 1990). The coverage here will be limited to general principles and to specific determinants that are applicable to legume production in lowland rice-based cropping systems. In general, BNF can supply N to a legume as long as a factor other than N is not limiting plant growth and an effective *Rhizobium* symbiosis is ensured (Bohlool et al., 1992). Abundant mineral N supply, however, inhibits or replaces BNF as a N source for legumes (Munns, 1977; Streeter, 1988). The interplay of *Rhizobium*, plant, and environment determines the proportions of legume N derived from atmospheric and soil sources.

*Rhizobial determinants*

Indigenous soil rhizobia are often ineffective or partially effective in BNF, but are quite successful in competing for nodule sites with introduced elite strains. This frequently results in no observed increase in BNF from rhizobial inoculation (Dowling and Broughton, 1986; Ham, 1980; Meade et al., 1985; Weaver and Frederick, 1974). Nonetheless, a response to inoculation may not be obtained if the number of native effective soil rhizobia exceeds 20 to 50 cells g<sup>-1</sup> soil (Singleton and Tavares, 1986; Thies et al., 1991a).

Soil flooding during the rice season decreases rhizobial numbers, but strains differ in survival ability (Boonkerd and Weaver, 1982), and often sufficient numbers of rhizobia for effective nodulation thrive (Weaver et al., 1987). Ladha et al. (1989a) reported survival in high numbers of rhizobia of the aquatic legume *Sesbania rostrata* in flooded rice rhizosphere. Rain splash and flooding often promote stem nodulation by *S. rostrata*; however, stem inoculation is probably beneficial under dry conditions (Ladha et al., 1991). Upland legumes following prolonged soil flooding may require inoculation to ensure effective symbiosis establishment. With proper




Fig. 4. Model describing legume inoculation response. Model inputs are numbers of indigenous rhizobia and soil N mineralization potential. Adapted from Thies et al. (1991b).

characterization of native soil rhizobia and soil N supplying capacity, the inoculation requirements of legumes may be predicted for various environments using the models developed by Thies et al. (1991b). One model which incorporates the numbers of indigenous rhizobia and soil N mineralization potential is presented in Figure 4. The model input variables can be obtained through soil analysis prior to legume planting.

*Plant determinants*

The potential BNF capacity may be defined as the aggregate of the per-day deficits in mineral N uptake during the legume growth cycle (George and Singleton, 1992). Therefore, the higher the N yield potential of a legume for a given growth duration and soil N supply, the higher would be the proportion and amount of plant N derived through BNF (Fig. 5). For the sake of simplicity, it is assumed that the soil N uptake rate remains constant throughout the growth period which is unlikely in field environments.

The amount of N derived from BNF is also altered by the pattern of plant N assimilation (George and Singleton, 1992). The legume-*Rhizobium* symbiosis becomes functional in BNF only after the early rhizobial infection and nodule formation (Hardy et al., 1971; Vincent,

9




Fig. 5. Conceptual relationship between nitrogen yield potential and nitrogen derived from BNF under a constant soil nitrogen supply. It is assumed arbitrarily that the increasing nitrogen yield potential has no influence on the amount of nitrogen derived from soil.

1980). Therefore, during early legume growth, soil N is the major source of plant N. It is likely that a legume capable of meeting its early N requirement from the soil, but having a subsequent N requirement exceeding the soil N supply, will benefit from BNF. Consequently, soybean, which has a high N requirement during the podfill period, also has a high  $N_2$  fixation rate during the same period (Afza et al., 1987; George and Singleton, 1992; Imsande, 1988, 1989; Zapata et al., 1987).

The efficiency of mineral N uptake by legumes further interacts with the N yield potential and the pattern of N assimilation in determining the amount of N derived from BNF. Under similar soil mineral N levels, legumes that are more efficient in extracting soil mineral N will benefit less from BNF. Using  $^{15}N$ , George and Singleton

(1992) determined in the field that common bean scavenges soil N much more efficiently than soybean, resulting in a lower proportion of N derived from BNF (Table 1). In their study, soybean derived 70% of its N from BNF compared to 23% by common bean at a similar flowering date, but the mineral N uptake efficiency of soybean was less than half that of common bean as determined by  $^{15}N$ . Consequently, KCl-extracted mineral N in the soil under soybean was almost twice as high as that under common bean.

#### Environmental determinants

Major environmental determinants of optimal BNF performance are soil-related constraints and climatic stresses. The soil mineral N supply is of dominant importance. Increased soil mineral N supply invariably decreases the proportion of N derived from BNF (Deibert et al., 1979; Eaglesham et al., 1983; George et al., 1988; George and Singleton, 1992; Gibson and Harper, 1985; Herridge and Brockwell, 1988; Thies et al., 1991b). However, soil mineral N and BNF can be complementary in meeting the N requirement of the legume (Bergersen et al., 1985; Deibert et al., 1979; Eaglesham et al., 1983; Herridge and Brockwell, 1988; Herridge et al., 1984).

On the other hand, the supply of mineral N in early legume growth can enhance BNF, a phenomenon often referred to as the 'starter N' effect. Insufficient soil N can restrict initial plant growth which limits subsequent plant growth and N demand, resulting in reduced need for BNF-supplied N. Alleviation of early N stress increases the amount of N derived from BNF (Table 2). Soybean plants receiving small amounts of mineral N formed a higher number

Table 1. Plant N, N derived from BNF, N uptake efficiency<sup>a</sup> and KCl-extractable soil N at flowering of field grown soybean and common bean (George and Singleton, 1992)

Legume	Plant N (kg N ha <sup>-1</sup> )	N from BNF (%)	<sup>15</sup> N uptake (%)	KCl-extr. N (mg N kg <sup>-1</sup> soil)
Soybean	58	70	5	17
Common bean	80	23	12	10

<sup>a</sup>N uptake efficiency is determined by using  $^{15}N$  isotope as tracer and is the percent of applied  $^{15}N$  fertilizer (3 kg N ha<sup>-1</sup>) assimilated by the plant.

Table 2. Mineral N effects on nodulation, total N, N derived through BNF and trifoliolate leaf area by soybeans grown for 33 days in pots amended with 0.8% bagasse (Singleton P W. 1991; pers. commun.)

Nitrogen applied ( $\mu\text{g N g}^{-1}$ soil)	Nodule (no. pot <sup>-1</sup> )	Nodule weight (mg pot <sup>-1</sup> )	Total N	N from BNF <sup>a</sup>		Area of trifoliolate <sup>b</sup> (cm <sup>2</sup> )
				Amount	% of total	
0	148	440	189	168	99	30
50	173	651	316	280	95	41
100	144	667	343	251	78	53
200	87	459	541	219	42	51
400	66	147	593	0	0	42

<sup>a</sup>Determined by the N difference method.

<sup>b</sup>Area of trifoliolate at 16 days of growth.

and weight of nodules and derived larger amounts of N from BNF than plants receiving either no N or large amounts of mineral N. The 'starter N' effect was due to increased plant growth as reflected by the increased area of the trifoliolate at 16 days.

From the data presented in Table 2 and in other reports (Deibert et al., 1979; Eaglesham et al., 1983; George and Singleton, 1992; Herridge and Brockwell, 1988; Thies et al., 1991a,b), a generalized conceptual relationship between soil N supply and N derived from BNF by legumes can be illustrated (Fig. 6). Initially, a 'starter N' effect tends to increase BNF but N from BNF approaches zero at the highest level of soil N supply. Increasing N supply, however, tends to increase total legume N.




Fig. 6. A generalized conceptual relationship between soil N supply and nitrogen derived from soil or BNF. Data from Table 2 and data reported by Deibert et al. (1979); Eaglesham et al. (1983); George and Singleton (1992); Herridge and Brockwell (1988); Thies et al. (1991a,b) are considered in the proposed relationship.

Other soil constraints include acidity, salinity, nutrient toxicities, and nutrient deficiencies (Craswell et al., 1987; Munns, 1977; O'Hara et al., 1988; Singleton and Bohlool, 1983; Singleton et al., 1985). Application of P in soils of low P availability (Cassman et al., 1981; Ssali and Keya, 1986) and liming acidic soils (Torres et al., 1988) increases total N accumulation by legumes. Further, N<sub>2</sub>-fixing legumes may have differing requirements of mineral nutrients compared to mineral N-dependent plants. Insufficient supply of mineral elements such as Ca, Mo, B and Fe can particularly limit N<sub>2</sub> fixation (O'Hara et al., 1988; Robson, 1978). External P requirements of N<sub>2</sub>-fixing legumes are higher than for mineral N-dependent legumes because of limited proliferation of the root system (Cassman et al., 1980, 1981).

Likewise, waterlogging or water deficit negatively influences either plant growth or the symbiosis (Bennett and Albrecht, 1984; Chapman and Muchow, 1985; De Vries et al., 1989a,b, Kucey et al., 1988; Minchin and Summerfield, 1976; Muchow, 1985; Sinclair et al., 1987; Smith et al., 1988; Timsina, 1989). Water stress reduces total N accumulation (Chapman and Muchow, 1985) and, especially during late growth stages, limits pod addition and subsequent seed N demand in grain legumes (De Vries et al., 1989b), thereby reducing N derived from BNF.

Additionally, climatic stresses such as low temperature, low solar radiation and unfavorable photoperiod can deleteriously affect plant and symbiotic performance (Becker et al., 1990; George et al., 1988, 1990; Hodges and French, 1985; Lawn and Williams, 1987; Muchow, 1985). Becker et al. (1990) reported that the number of

days required to produce 100 kg N by *S. rostrata* was higher when the plants were grown during short days than during long days.

#### **Nitrogen contribution from legumes in rice lowlands**

Inclusion of legumes in the cropping pattern with rice can increase the total N output from the cropping system but does not necessarily result in a positive soil N contribution. The amount of N derived from BNF, the net effect of the legume on N losses and the amount of N removed with harvested plant material will determine whether there is a positive net soil N balance.

#### *Food and forage legumes*

The amount of N removed from the land and the amount available for recycling will vary with the N harvest index (NHI), which is the fraction of total N removed in the harvested produce. To achieve a positive soil N balance after legume harvest, it is essential that the amount of N derived from BNF exceed the amount of N removed in the legume produce, i.e. the proportion of total N derived from BNF should exceed NHI. Nitrogen removal through grain or forage may deplete soil N. Of the 23 cases of grain legumes compiled by Peoples and Craswell (1992), N balances were positive in 17 cases and negative or unchanged in six cases.

Examples of soil N gain or depletion by grain legumes under varying N availability and yield potentials are presented in Table 3. The data demonstrate that even with a large N input from BNF, the net soil N balance may be negative. Despite a high soybean crop N value (404 kg N ha<sup>-1</sup>) and high BNF input (290 kg N ha<sup>-1</sup>) the net soil N balance was negative because NHI exceeded the proportion of N derived from BNF (Table 3, Chapman and Myers, 1987). In the above example, a slight reduction in NHI or a slight increase in N derived from BNF would have resulted in a large soil N gain since the N yield was large. Since NHI is an attribute less amenable to field management, obtaining a posi-

tive soil N gain may primarily depend on strategies to increase BNF.

As Peoples and Craswell (1992) have also argued, even when there is little or no net soil N gain, the depletion of soil N is still much less after a grain legume than a nonlegume. Nevertheless, grain legumes increase the N output of the cropping system that is economically usable. Management approaches to limit the amount of N removed in an economic produce to the amount of N derived from BNF will at least prevent depletion of native soil N. One approach which meets the purposes of both grain N harvest and residue N production is the use of dual-purpose food legumes (Buresh and De Datta, 1991; Kulkarni and Pandey, 1988).

The effect of cutting and removing plant residues as fodder for farm animals or of allowing farm animals to graze on legume cover-cropped fallows has a similar effect as exporting N through harvested grains (Myers and Wood, 1987). The amount of N returned to the land depends on how the animal residues are managed. Nevertheless, grazing by animals is likely to remove less N from land than cutting and removing plant residues because part of the forage ingested by grazing animals is returned to the land via urine and dung.

#### *Stem-nodulating aquatic green manure legumes*

Stem-nodulating aquatic green manure legumes are fast-growing and can accumulate large amounts of N in 45 to 60 days of growth (Ladha et al., 1992). Data on the most frequently studied stem-nodulating aquatic green manure legume, *S. rostrata*, show that it derives larger amounts of N from BNF than do root-nodulating waterlogging-tolerant legumes, under either flooded or non-flooded conditions and it accumulates as much or more N under flooded conditions as do root-nodulating legumes such as *S. cannabina* and *S. sesban* grown under drained conditions (Morris et al., 1989; Ndoye and Dreyfus, 1988; Pareek et al., 1990). These features make stem-nodulating legumes uniquely suitable for areas prone to flooding in the pre-rice season.

The reported tolerance of BNF by stem-nodulating aquatic legumes to mineral N under

Table 3. Net soil nitrogen gains due to BNF following cropping of legumes at varying yield-potential and soil N availability levels<sup>a</sup>

Legume	N status at planting			Total crop N	% total crop N derived from BNF	% total crop N in produce removed	Net soil N gain due to BNF	Reference
	Total N (%)	Miner. N ( $\mu\text{g g}^{-1}$ )	Fert. N ( $\text{kg N ha}^{-1}$ )					
Cowpea	0.12	– <sup>b</sup>	30	84	19	67	– 40	Sisworo et al. (1990) <sup>c</sup>
	–	81	0	125	70	64	+ 7	Ofori et al. (1987)
	–	81	25	153	54	64	– 16	Ofori et al. (1987)
	–	–	25	100	70	49	+ 21	Eaglesham et al. (1982) <sup>d</sup>
Mungbean	–	–	100	98	36	53	– 17	Eaglesham et al. (1982)
	0.04	3.5	–	172	65	52	+ 23	Chapman and Myers (1987)
	0.16	–	0	60	67	52	+ 9	Suwanarit et al. (1986)
Peanut	0.16	–	20	62	56	53	+ 2	Suwanarit et al. (1986)
	0.16	–	0	241	60	48	+ 29	Suwanarit et al. (1986)
Soybean	0.16	–	20	250	62	47	+ 38	Suwanarit et al. (1986)
	0.04	3.5	–	404	72	73	– 6	Chapman and Myers (1987)
	0.12	–	30	90	33	62	– 26	Sisworo et al. (1990) <sup>e</sup>
	0.16	–	0	186	65	9	+104	Suwanarit et al. (1986)
	0.16	–	20	174	68	9	+103	Suwanarit et al. (1986)
	–	Low	0	246	68	62	+ 14	Thurlow and Hiltbold (1985) <sup>f</sup>
	0.30	–	33	222	46	73	– 60	Zapata et al. (1987)
0.13	38	0	347	9	47	–132	Bergersen et al. (1989) <sup>g</sup>	
0.13	19	0	267	56	68	– 32	Bergersen et al. (1989) <sup>h</sup>	

<sup>a</sup>Values derived from available data when originally not given. <sup>b</sup>Data not available. <sup>c</sup>Average for four crops. <sup>d</sup>Average for four cultivars. <sup>e</sup>Average for two crops. <sup>f</sup>Average for 'Lee' soybean at six sites. <sup>g</sup>Pre-fallowed soil; total N includes roots, nodules and fallen leaves. <sup>h</sup>Pre-cropped (oats) soil; total N includes roots, nodules and fallen leaves.

13

flooded conditions (Becker et al., 1986, 1990; Dreyfus and Dommergues, 1980; Eaglesham and Szalay, 1983; Ladha et al., 1992) may suggest that these legumes prefer N from BNF over soil N. Whether this characteristic leads to soil NO<sub>3</sub> accumulation in aerobic soil is yet to be determined. However, similar total N accumulation and similar or decreased N<sub>2</sub> fixation by *S. rostrata* in non-flooded compared to flooded soil (Morris et al., 1989; Ndoye and Dreyfus, 1988) may indicate a significant capacity for soil N uptake in aerobic soil.

#### *Legumes in the post-rice as opposed to the pre-rice niche*

Post-rice legumes with subsequent fallow periods before wet-season rice, and pre-rice legumes grown immediately before the wet-season rice, may have varied effects on native soil N and on N contribution to rice. It appears that soil NO<sub>3</sub> accumulation in fallows following legumes could be significant. Strong et al. (1986) presented data on soil mineral N (NO<sub>3</sub> and NH<sub>4</sub>) accumulation during 6 months of fallow following several legumes, oil seeds, and cereals. Following harvest, crop residues were left in the field, and the soil was tilled to a depth of 8 cm. Soil mineral N accumulation during the fallow period was generally higher following legumes than following oilseeds or cereals. They attributed differences in soil mineral N to differences in N content and C:N ratios of the crop residues.

High quantities of mineral N following the legumes are not advantageous to rice. Nitrogen

mineralization during the post-legume fallow, and subsequent N loss upon soil flooding will remove the mineral N from the system. There is evidence that, regardless of post-rice treatments, soil NO<sub>3</sub> may rapidly drop to uniformly low levels by the start of the wet-season rice crop. The higher soil NO<sub>3</sub> level following post-rice legumes compared to weeds reported by Buresh and De Datta (1991) in a Philippine soil persisted until the next sampling, 3 weeks later. However, NO<sub>3</sub> in all treatments declined to a low level before the start of the wet-season rice culture (Table 4, IRRI-NifTAL-IFDC collaborative research). Weed growth during the fallow period after the legume harvest probably assimilated some of the soil NO<sub>3</sub>, but NO<sub>3</sub> loss was likely to have been large in the treatments.

Results of Hamid et al. (1984) show no difference in rice yields when rice is grown six weeks after legume harvest or after approximately 18 weeks of continuous fallow. However, when sorghum was the post-rice crop the rice yield was severely depressed. In a study by John et al. (1989), yield of rice immediately after cowpea cropping with all aboveground residues removed or after a weedy fallow were similar. The addition of pre-rice cowpea residues, however, did increase rice yield.

#### **Maximizing total nitrogen capture by legumes in rice lowlands**

Maximizing N gains through legumes in rice lowlands entails approaches that maximize BNF

Table 4. Soil-nitrate accumulation during the fallow period following post-rice legumes and weedy fallow in a lowland rice-based cropping system

Post-rice treatment	Sampling time for nitrate		
	Legume harvest <sup>a</sup> (0–80 cm)	3 wk after harvest (0–60 cm)	6 wk after harvest <sup>b</sup> (0–60 cm)
	(kg NO <sub>3</sub> -N ha <sup>-1</sup> )		
Cowpea	53	47	8
Mungbean	53	52	9
Nodulating soybean	54	50	8
Nonnodulating soybean	47	40	8
Weedy fallow	32	28	7

<sup>a</sup>Data from Buresh and De Datta (1991).

<sup>b</sup>Start of wet-season rice culture.

while minimizing soil N losses. Soil and crop management should ensure establishment of an effective symbiosis, and prevent excessive accumulation and subsequent loss of soil  $\text{NO}_3$ . The quantity of mineral N present at planting and mineralized during the legume growing season will influence the amount of N derived from BNF.

#### *Managing soil nitrate supply*

Soil nitrate associated with the aerobic phase in ricelands can be a source of N for both legume and rice crop, if it can be managed effectively. A lack of response of legumes to external N application indicates that the initial N needs are generally met by native soil  $\text{NO}_3$ . Moreover, legume N needs may be limited by a reduced yield potential due to inadequate supplies of water and nutrients other than N. Nitrate-N assimilated by the legume is conserved in the system against possible loss due to subsequent soil flooding. The management objective then is to utilize soil  $\text{NO}_3$  that might otherwise be lost, and to maximize BNF.

#### *Allowing for increased nitrate uptake*

The total amount of N that can be accumulated by a legume crop in a given environment is fixed by physiological and environmental determinants. Legume management aims to meet this potential by maximizing the plant demand for N and by optimizing the soil mineral N supply to maximize aggregate plant N accumulation through soil uptake and BNF.

Soil water deficits often limit legume growth and plant N demand in lowland rice-based cropping systems. Establishing a post-rice legume crop prior to or immediately after the rice harvest is crucial in maximizing the use of the entire residual soil water reservoir. Water deficits are likely during the late growth stages of a post-rice legume because of a receding water table (Senthong and Pandey, 1989; Timsina, 1989).

In areas with limited soil-water availability, drought-tolerant legumes such as pigeon pea and indigo are suitable post-rice legumes that accumulate N throughout the dry season. Peanut is reported to withstand relatively larger soil water deficits than most grain legumes because of a

higher root density in the lower soil depths (De Vries et al., 1989a; Senthong and Pandey, 1989).

Early rains facilitate establishment of pre-rice legumes. However, intermittent waterlogging during this season is deleterious to crop growth and yield formation (Alam, 1989; Minchin et al., 1978; Timsina, 1989), thus limiting N derived from BNF. Alam (1989) observed that cowpea and mungbean are unsuitable in poorly drained soils where short-term flooding occurs during the pre-rice dry-to-wet transition period. Aquatic legumes can increase N input in these soils. Green manure legumes such as *S. rostrata* and *Aeschynomene afraspera* are adapted to flooded soil conditions (Ladha et al., 1992). Aquatic legumes can possibly conserve soil  $\text{NO}_3$  during growth in nonflooded soil. Stem nodule BNF can additionally provide N to the plant when the soil is flooded.

Increased soil water availability not only enhances legume growth, but also promotes soil  $\text{NO}_3$  production. Ideally, the previously accumulated  $\text{NO}_3$  should be depleted during early legume growth, and  $\text{NO}_3$  supply thereafter should be less than plant N demand. A reduced soil  $\text{NO}_3$  supply combined with sufficient water for legume growth can result in enhanced BNF as the legume enters the reproductive phase.

#### *Minimizing soil supply of nitrate*

Management may play a role in influencing the amount of  $\text{NO}_3$ -N formed in aerobic soil and its subsequent assimilation by plants or loss from the soil-plant system. Planting legumes with no-tillage is a strategy to decrease soil  $\text{NO}_3$ , since tillage can increase soil  $\text{NO}_3$  levels. Peoples and Craswell (1992) have presented data showing improved  $\text{N}_2$  fixation by soybean under no-tillage. Therefore, relay seeding of legumes before harvest of the preceding rice crop might be suitable for maximizing residual water use and decreasing  $\text{NO}_3$  accumulation.

Rice stubble management can influence the amount of  $\text{NO}_3$  produced in soil. Incorporating the stubble rather than leaving it on the soil surface or burning it following harvest, significantly reduces soil  $\text{NO}_3$  (Bacon, 1987, 1990). Bacon (1987) reported reduced soil  $\text{NO}_3$  levels for up to 12 weeks, and reduced N uptake by the following wheat crop, when rice stubble

(15 t ha<sup>-1</sup>) was incorporated soon after harvest.

Nitrogen depletion by crops preceding the legume or competition for N in a legume-nonlegume intercropping system, influences the amount of NO<sub>3</sub> available to the legume crop. Soil NO<sub>3</sub> is low immediately after a flooded rice crop. Planting a legume immediately after the rice crop might maximize BNF provided soil water is adequate. Total N derived from BNF by several legumes increased to 93–95% when grown immediately after a dry-season rice crop, compared to 65–72% when grown after a prolonged fallow (Chapman and Myers, 1987). Legumes following cereals usually fix higher amounts of N than legumes following a fallowed soil (Bergersen et al., 1985, 1989) because of decreased soil mineral N levels. Likewise, cereal-legume intercropping can enhance N derived from BNF by the legume (Fujita et al., 1992; Rerkasem et al., 1988; Rerkasem and Rerkasem, 1988) due to mineral N depletion by the cereal.

#### *Managing inefficient nitrate uptake by legumes*

Legumes that are inefficient in mineral N uptake might meet their N requirement through BNF because of a complementary relationship between soil N uptake and BNF. This might cause no substantial change in the amount of legume N accumulated but might lead to larger net NO<sub>3</sub> loss from lowland rice soils. While some reports indicate inefficient NO<sub>3</sub> uptake during the legume reproductive phase (Imsande, 1986; Imsande and Edwards, 1988) others indicate large N uptake from aerobic soil throughout the legume reproductive phase (Afza et al., 1987; George and Singleton, 1992). Herridge and Bergersen (1988) described the “sparing” of soil NO<sub>3</sub> by N<sub>2</sub>-fixing legumes, where more NO<sub>3</sub> remains in the soil after a legume compared to a nonlegume. Nonetheless, excess NO<sub>3</sub> may remain in a lowland soil after the growth of a legume. This NO<sub>3</sub> can be lost after soil flooding.

Larger amounts of residual soil mineral N have been reported after cropping of legumes than of other crops (Doughton and MacKenzie, 1984; McEwen et al., 1989; Stong et al., 1986). Reports on post-legume soil NO<sub>3</sub> levels in lowland rice soils are few. Buresh et al. (1989) reported 25 kg NO<sub>3</sub>-N ha<sup>-1</sup> in the top 60 cm soil

layer at harvest of mungbean grown after lowland rice. Table 4 shows identical soil NO<sub>3</sub> levels (52 kg NO<sub>3</sub>-N ha<sup>-1</sup>) in the top 80 cm layer at harvest of soybean, cowpea, mungbean and a non-nodulating soybean isolate in a lowland rice soil, but a reduced NO<sub>3</sub> level (32 kg NO<sub>3</sub>-N ha<sup>-1</sup>) under a weedy fallow. The corresponding NO<sub>3</sub> levels in the top 20 cm soil layer were 30 kg NO<sub>3</sub>-N ha<sup>-1</sup> for legumes and 12 kg NO<sub>3</sub>-N ha<sup>-1</sup> for a weedy fallow, indicating that the weeds were more effective than legumes either in depleting soil NO<sub>3</sub> or in maintaining low NO<sub>3</sub> levels by other mechanisms. Nitrogen fixation by the legumes, however, had not influenced the soil NO<sub>3</sub> levels, as the non-nodulating soybean and the N<sub>2</sub>-fixing legumes maintained similar soil NO<sub>3</sub> levels.

Weed N accumulation during the dry season in lowland rice soils reported by Buresh and De Datta (1991) ranged from 11 to 29 kg N ha<sup>-1</sup>. It is likely that most legumes that derived at least 50% of their N from BNF would have equal or larger amounts of N removed from soil compared to weeds; there might be other effects of weeds on soil NO<sub>3</sub> than uptake. Soil NO<sub>3</sub> levels are usually lower in grass and hay crop fields than in fields of row-planted crops such as maize (Magdoff, 1982).

It follows that strategies to develop or select legume-Rhizobium symbioses tolerant to soil NO<sub>3</sub> (Betts and Herridge, 1987; Carroll et al., 1985; Hansen et al., 1989; Herridge and Betts, 1988) or legumes with reduced ability to use NO<sub>3</sub> (Carroll and Gresshoff, 1986; Nelson et al., 1983) might not be advantageous in lowland rice-based cropping systems unless methods other than plant uptake are found to prevent the buildup and subsequent loss of NO<sub>3</sub>. Suitable legume genotypes for lowland rice soils might be those that can deplete soil NO<sub>3</sub> while still deriving large amounts of N from BNF. Part of the NO<sub>3</sub> could possibly also be assimilated by weeds either during a post-legume fallow period before rice or during the legume crop.

#### **Use of legume nitrogen for lowland rice**

Use of legume green manures and grain legume residues as a N source to rice has been well

reviewed (Buresh and De Datta, 1991; Singh et al., 1991). Soil incorporation of legume green manures (Ladha et al., 1989b; Morris et al., 1986a, 1989; Ventura et al., 1987) and of legume harvest residues (Alam, 1989; John et al., 1989; Norman et al., 1990) in lowland rice culture generally results in increased N uptake and yield of rice. However, the efficiency of residue N use by rice might be exaggerated, because most reports attribute the positive influence exclusively to a N effect.

#### Belowground residues

Data compiled by Buresh and De Datta (1991) showed legume root N accumulation ranging from 3 to 50 kg N ha<sup>-1</sup>. They assessed the contribution from belowground legume residues as a N source for rice to be negligible in most cases. This would be especially true with grain legumes grown to maturity because of the remobilization of N from roots and nodules to the grain. The belowground N contribution may be significant if legumes are incorporated as green manure during the flowering and early reproductive stages when root nodulation could be abundant. Nodule biomass at flowering of several grain legumes ranged approximately from 4 to 118 kg ha<sup>-1</sup> (Table 5, Abaidoo et al., 1990; George et al., 1987; Thies et al., 1991a). Assuming a maximum 7% N content, this nodule biomass contained approximately 0.3 to 8 kg N ha<sup>-1</sup>. Adding the amount of N present in

the roots, total belowground N could become significant in certain cases. However, the value of this N to lowland rice is questionable because N-rich nodule tissue would decompose and release N very rapidly. Morris et al. (1989) found no difference in rice N-gain between legume green manure grown in situ or that transported from a nearby field, suggesting no N contribution from belowground biomass. Nevertheless, quantitative data on belowground residue N are required to understand N cycling in rice-based cropping systems.

#### Aboveground residues

The amount of N that can be incorporated through some legume green manure crops (Ladha et al., 1992) is larger than the N needs of rice. Buresh and De Datta (1991) compiled data on amounts of N incorporated with aboveground green manure and grain legume residues in lowland rice-based cropping systems. Green manure N ranged from 21 to 267 kg N ha<sup>-1</sup>; N in aboveground residues of grain legumes from 17 to 101 kg N ha<sup>-1</sup>. Nitrogen fertilizer substitution for lowland rice by incorporated green manures ranged from 24 to 137 kg N ha<sup>-1</sup>, and by grain legume residues from 37 to 100 kg N ha<sup>-1</sup>. Similar data are presented by Singh et al. (1991). Furoc and Morris (1989) suggested that the addition of N in *S. rostrata* green manure in excess of 100 kg ha<sup>-1</sup> has limited value for increasing rice yield. Moreover, large application of N-rich legume residues has the potential to supply excessive N to rice, leading to crop lodging and yield reduction (Alam, 1989).

#### Nitrogen release from legume residues

Many studies demonstrate a rapid release of N, and an increase in soil NH<sub>4</sub>, followed by a plateau or decline in NH<sub>4</sub> levels after green manure incorporation (Khind et al., 1985; Nagarajah et al., 1989; Palm et al., 1988). The release of NH<sub>4</sub> from green manure is, therefore, not matched by the N uptake by rice. Thus, NH<sub>4</sub> released may be subject to immobilization or loss. Morris et al. (1986b) reported that recovery of N from green manure was 33%, similar to that from N fertilizer. In contrast, Ventura et al.

Table 5. Nodule biomass at flowering of legumes in tropical soils

Legume	Nodule biomass (kg ha <sup>-1</sup> )	Reference
Common bean	4– 68	Abaidoo et al., (1990) <sup>a</sup>
	8– 65	Thies et al., (1991a) <sup>b</sup>
Cowpea	38–105	Thies et al., (1991a)
Lima bean	13– 67	Thies et al., (1991a)
Soybean	16– 75	Abaidoo et al., (1990) <sup>a</sup>
	24– 87	Thies et al., (1991a) <sup>b</sup>
	26–118	George et al., (1987) <sup>c</sup>

<sup>a</sup>Across two sites of differing mean temperature and three N application regimes.

<sup>b</sup>Across several sites of varying soil N status and yield potential.

<sup>c</sup>Across three sites of varying soil N status and mean temperature.

(1987) reported up to 75% recovery of *S. rostrata* green manure N by rice. The differences in green manure N recoveries reported may be partly due to differences in the amount applied, in C:N ratio, and in timing and method of application. Recent data show that  $\text{NH}_4\text{-N}$  loss from green manure was lower than that with equivalent amounts of fertilizer N (Biswas, 1988). Rice recovery of green manure N is larger than from urea, and green manure increases the recovery of urea-N (Diekmann, 1991). Supplementing moderate amounts of basally applied green manure N with topdressed fertilizer N may maximize both green manure and fertilizer N recoveries (Morris et al., 1986b).

### Conclusions and research directions

The large amounts of  $\text{NO}_3\text{-N}$ , which accumulate in aerobic soil during the non-flooded season, can be lost for lowland rice upon flooding. Prevention of this loss might make a significant N contribution to crops in the dry-to-wet transition period or to a subsequent rice crop. In addition, the inclusion of legumes for grain, forage, and green manure production might increase the total N output from the cropping system and further add to soil N through BNF. The N derived from BNF by legumes can be increased by maximizing plant N demand and by minimizing soil  $\text{NO}_3$  supply. A post-rice nonlegume crop followed by a pre-rice green manure legume or intercropping legumes with nonlegumes during both the post-rice and pre-rice seasons might maximize the conservation and effective use of both soil  $\text{NO}_3$  and atmospheric N in lowland rice-based cropping systems. Legumes would be ideal crops for soils low in native N, where other upland crops may grow poorly without external N supply. Where soils are left fallow during the non-flooded season due to production constraints, increasing the legume component in the weed flora may be suitable for capturing both soil  $\text{NO}_3$  and atmospheric N.

Our knowledge of the processes and factors regulating  $\text{NO}_3$  accumulation and loss during the nonflooded season and the dry-to-wet season transition period and how legume BNF fits into this complex are still very limited. Research is

needed to understand the mechanisms of N gains and losses and to identify and refine appropriate soil and crop management practices to maximize soil N and legume BNF use in lowland rice-based cropping systems.

So far, research on N dynamics and balances in legume-lowland rice sequences has been primarily with pre-rice legumes grown just prior to wet season rice. Because dry-season legumes are also an important post-rice crop in lowland rice-based cropping system, there is a need to ascertain the effects of post-rice legumes with subsequent fallow periods before wet-season rice on native soil N dynamics and N contribution to rice.

Maintaining soil N levels in the rice-based farming system is not the dominant objective of the farmer in the choice of cropping pattern and management practices. Overall profitability shall continue to dominate as the basic farmer objective. However, there is a major potential for directing research toward identification of management practices aimed at maintaining long-term soil fertility while providing profitability in the short term. This will depend upon a more solid knowledge base of system-level N dynamics.

### References

- Abaidoo R C, George T, Bohlool B B and Singleton P W 1990 Influence of elevation and applied nitrogen on rhizosphere colonization and competition for nodule occupancy by different rhizobial strains on field-grown soybean and common bean. *Can. J. Microbiol.* 36, 92-96.
- Afza R G, Zapata H H and Danso S K A 1987 Effect of delayed soil and foliar N fertilization on yield and  $\text{N}_2$  fixation of soybean. *Plant and Soil* 97, 361-368.
- Alam M 1989 Stability and comparative nitrogen contribution of pre-rice legumes: Grain legume vs. green manure. M.Sc. thesis, University of the Philippines, Los Baños, Philippines.
- Bacon P E 1987 Effect of nitrogen fertilization and rice stubble management techniques on soil moisture content, soil nitrogen status, and nitrogen uptake by wheat. *Field Crops Res.* 17, 75-90.
- Bacon P E 1990 Effects of stubble and N fertilization management on N availability and uptake under successive rice (*Oryza sativa* L.) crops. *Plant and Soil* 121, 11-19.
- Bacon P E, McGarity J W, Hoult E H and Alter D 1986 Soil mineral nitrogen concentration within cycles of flood irrigation: Effect of rice stubble and fertilization management. *Soil Biol. Biochem.* 18, 173-178.

- Becker M, Alazard D and Ottow J C G 1986 Mineral nitrogen effect on nodulation and nitrogen fixation of the stem-nodulating legume *Aeschynomene afraspera*. Z. Pflanzenernaehr. Bodenk. 149, 485-491.
- Becker M, Ladha J K and Ottow J C G 1990 Growth and N<sub>2</sub> fixation of two stem-nodulating legumes and their effect as green manure on lowland rice. Soil Biol. Biochem. 22, 1109-1119.
- Bennett J M and Albrecht S L 1984 Drought and flooding effects on N fixation, water relations and diffusive resistance of soybean. Agron. J. 76, 735-740.
- Bergersen F J, Turner G L, Chase D L, Gault R R and Brockwell J 1985 The natural abundance of <sup>15</sup>N in an irrigated soybean crop and its use for the calculation of nitrogen fixation. Aust. J. Agric. Res. 36, 411-423.
- Bergersen F J, Brockwell J, Gault R R, Morthorpe L, Peoples M B and Turner G L 1989 Effects of available soil nitrogen and rates of inoculation on nitrogen fixation by irrigated soybeans and evaluation of  $\delta^{15}\text{N}$  methods for measurement. Aust. J. Agric. Res. 40, 763-780.
- Betts J H and Herridge D F 1987 Isolation of soybean capable of nodulation and nitrogen fixation under high levels of nitrate supply. Crop Sci. 27, 1156-1161.
- Birch H F 1960 Nitrification in soils after different periods of dryness. Plant and Soil 12, 81-96.
- Biswas T K 1988 Nitrogen dynamics and nitrogen-15 balance in lowland rice as affected by green manure and urea application. Ph.D. thesis, Indian Agric. Res. Inst., New Delhi, India.
- Bohlool B B, Ladha J K, Garrity D P and George T 1992 Biological nitrogen fixation for sustainable agriculture: A perspective. Plant and Soil 141, 1-11.
- Boonkerd N and Weaver R W 1982 Survival of cowpea rhizobia in soil by soil temperature and soil moisture. Appl. Environ. Microbiol. 43, 585-589.
- Bouldin D R 1986 The chemistry and biology of flooded soils in relation to the nitrogen economy in rice fields. In Nitrogen Economy of Flooded Rice Soils. Eds. S K De Datta and W H Patrick Jr. pp 1-14. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Broadbent F E 1984 Plant use of soil nitrogen. In Nitrogen in Crop Production. Ed. R D Hauck. pp 171-182. ASA-CSSA-SSSA, Madison, WI.
- Buresh R J and De Datta S K 1991 Nitrogen dynamics and management in rice-legume cropping systems. Adv. Agron. 45, 1-59.
- Buresh R J, Woodhead T, Shepherd K D, Flordelis E and Cabangon R C 1989 Nitrate accumulation and loss in a mungbean/lowland rice cropping system. Soil Sci. Soc. Am. J. 53, 477-482.
- Carangal V R, Tengco P L, Miah N I and Topar-Ngarm A 1988 Food and forage crop intercropping under rainfed lowland conditions. In Proceedings of the Crop-Animal Systems Research Workshop. pp 435-451. Serdang, Malaysia.
- Carroll B J and Gresshoff P M 1986 Isolation and initial characterization of constitutive nitrate reductase-deficient mutants NR328 and NR345 of soybean (*Glycine max*). Plant Physiol. 81, 572-576.
- Carroll B J, McNeil D L and Gresshoff P M 1985 A supernodulating and nitrate tolerant symbiotic (*nts*) soybean mutant. Plant Physiol. 78, 34-40.
- Cassman K G, Whitney A S and Stockinger K R 1980 Root growth and dry matter distribution of soybean as affected by phosphorus stress, nodulation, and N source. Crop Sci. 20, 239-244.
- Cassman K G, Whitney A S and Fox R L 1981 Phosphorus requirements of soybean and cowpea as affected by mode of N nutrition. Agron. J. 73, 17-22.
- Chaney K 1990 Effect of nitrogen fertilizer rate on soil nitrate nitrogen content after harvesting winter wheat. J. Agric. Sci. Cambridge 114, 171-176.
- Chapman A L and Muchow R C 1985 Nitrogen accumulated and partitioned at maturity by grain legumes grown under different water regimes in a semi-arid tropical environment. Field Crops Res. 11, 69-79.
- Chapman A L and Myers R J K 1987 Nitrogen contributed by grain legumes in rotation with rice on the Cununurra soils of the Ord Irrigation Area, Western Australia. Aust. J. Exp. Agric. 27, 155-163.
- Craswell E T, Loneragan J F and Keerati-Kasikorn P 1987 Mineral constraints to food legume crop production in Asia. In Food Legume Improvement for Asian Farming Systems: Proceedings of an International Workshop. Khon Kaen, Thailand, September 1986. ACIAR Proc. Ser. No. 18. Eds. E S Wallis and D E Byth. pp. 99-111.
- Deibert E J, Bijeriego M and Olson R A 1979 Utilization of <sup>15</sup>N fertilizer by nodulating and nonnodulating soybean isolines. Agron. J. 71, 717-722.
- De Datta S K 1981 Principles and Practices of Rice Production. Wiley, New York. 618 p.
- De Datta S K and Buresh R J 1989 Integrated nitrogen management in irrigated rice. Adv. Soil Sci. 10, 143-169.
- De Datta S K and Patrick W H 1986 Nitrogen Economy of Flooded Rice Soils. Martinus Nijhoff Publishers, Dordrecht, The Netherlands. 186 p.
- De Vries J D, Bennett J M, Albrecht S L and Boote K J 1989a Water relations, nitrogenase activity, and root development of three grain legumes in response to soil water deficits. Field Crops Res. 21, 215-226.
- De Vries J D, Bennett J M, Boote K J, Albrecht S L and Maliro C E 1989b Nitrogen accumulation and partitioning by three grain legumes in response to soil water deficits. Field Crops Res. 22, 33-44.
- Diekmann K H 1991 Nitrogen transformation processes involving <sup>15</sup>N labeled *Sesbania* and *Aeschynomene afraspera* green manure in lowland rice (*Oryza sativa* L.). Ph. D. Thesis. Justus-Liebig University, Giessen, Germany.
- Dowdell R J, Crees R and Cannell R Q 1983 A field study of contrasting methods of cultivation on soil nitrate content during autumn, winter and spring. J. Soil Sci. 34, 367-379.
- Dowling D N and Broughton W J 1986 Competition for nodulation of legume. Annu. Rev. Microbiol. 40, 131-157.
- Doughton J A and MacKenzie J 1984 Comparative effects of black and green gram (mung beans) and grain sorghum on soil mineral nitrogen and subsequent grain sorghum yields on the Eastern Darling Downs. Aust. J. Exp. Agric. Anim. Husb. 24, 244-249.
- Dreyfus B and Dommergues Y 1980 Non-inhibition de la fixation d'azote atmospherique par l'azote combine chez

- une légumineuse a nodules caulinaires. *Sesbania rostrata*. C.R. Acad. Sc. Paris 291. 767-770.
- Eaglesham A R J and Szalay A A 1983 Aerial stem nodules on *Aeschynomene* pp. Plant Sci. Lett. 29, 265-272.
- Eaglesham A R J, Ayanaba A, Rao V R and Eskew D L 1982 Mineral N effects on cowpea and soybean crops in a Nigerian soil. II. Amounts of N fixed and accrual to the soil. Plant and Soil 68, 183-192.
- Eaglesham A R J, Hassouna S and Seegers R 1983 Fertilizer-N effects on N<sub>2</sub> fixation by cowpea and soybean. Agron. J. 75, 61-66.
- Eskew D L, Eaglesham A R J and App A 1981 Heterotrophic N<sub>2</sub> fixation and distribution of newly fixed nitrogen in a rice-flooded soil system. Plant Physiol. 68, 48-52.
- Flinn J C and De Datta S K 1984 Trends in irrigated rice-yields under intensive cropping at Philippine Research Stations. Field Crops Res. 9, 1-15.
- Fujita K and Ofosu-Budu Ogata S 1992 Biological nitrogen fixation in mixed legume-cereal systems. Plant and Soil 141, 155-176.
- Furoc R E and Morris R A 1989 Apparent recovery and physiological efficiency of nitrogen in *Sesbania* incorporated before rice. Agron. J. 81, 797-802.
- Garrity D P and Flinn J C 1988 Farm-level management systems for green manure crops in Asian rice environments. In Sustainable Agriculture: Green Manure in Rice Farming. pp 111-130. International Rice Research Institute, Los Baños, Philippines.
- Garrity D P, Oldeman L R, Morris R A and Lenka D 1986 Rainfed lowland rice ecosystems: Characterization and distribution. In Progress in Rainfed Lowland Rice. pp 3-23. International Rice Research Institute, Los Baños, Philippines.
- George T and Singleton P W 1992 Relationship between nitrogen assimilation traits and dinitrogen fixation in soybean and common bean. Agron. J. (In press).
- George T, Bohlool B B and Singleton P W 1987 *Bradyrhizobium japonicum*-environment interactions: Nodulation and interstrain competition in soils along an elevational transect. Appl. Environ. Microbiol. 53, 1113-1117.
- George T, Singleton P W and Bohlool B B 1988 Yield soil nitrogen uptake and nitrogen fixation by soybeans from four maturity groups grown at three elevations. Agron. J. 80, 563-567.
- George T, Bartholomew D P and Singleton P W 1990 Effect of temperature and maturity group on phenology of field grown nodulating and nonnodulating soybean isolines. Biotronics 19, 49-59.
- Gibson A H and Harper J E 1985 Nitrate effect on nodulation of soybean by *Bradyrhizobium japonicum*. Crop Sci. 25, 497-501.
- Hadas A, Feigenbaum S, Feigin A and Portnoy R 1986 Nitrogen mineralization in profiles of differently managed soil types. Soil Sci. Soc. Am. J. 50, 314-319.
- Ham G E 1980 Inoculation of legumes with *Rhizobium* in competition with naturalized strains. In Nitrogen Fixation. Vol. II. Eds. W E Newton and W H Orme-Johnson. pp 131-138. University Park Press, Baltimore, MD.
- Hamid A, Paulsen G M and Zandstra H G 1984 Performance of rice grown after upland crops and fallow in the humid tropics. Trop. Agric. (Trinidad) 61, 305-310.
- Hansen A P, Peoples M B, Gresshoff P M, Atkins C A, Pate J S and Carroll B J 1989 Symbiotic performance of super-nodulating soybean (*Glycine max* (L.) Merrill) mutants during development on different nitrogen regimes. J. Exp. Bot. 40, 715-724.
- Hardy R W F, Burns R C, Herbert R R, Hosten R D and Jackson E K 1971 Biological nitrogen fixation: A key to world protein. Plant and Soil Spec. Vol. 561-590.
- Harwood R R and Price E C 1976 Multiple cropping in tropical Asia. In Multiple Cropping, pp 11-40. ASA special publication number 27, American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, WI.
- Herlihy M 1979 Nitrogen mineralization in soils of varying texture, moisture and organic matter. I. Potential and experimental values in fallow soils. Plant and Soil 53, 255-267.
- Herridge D F 1986 Nitrogen fixation dynamics by rainfed grain legume crops: Potential for improvement. In Transactions of the XIII Congress of International Society of Soil Science. pp 794-804. Hamburg, Germany.
- Herridge D F, Roughly R J and Brockwell J 1984 Effects of rhizobia and soil nitrate on the establishment and functioning of the soybean symbiosis in the field. Aust. J. Agric. Res. 35, 149-161.
- Herridge D F and Bergersen F J 1988 Symbiotic nitrogen fixation. In Advances in Nitrogen Cycling in Agricultural Ecosystems. Ed. J R Wilson. pp 46-65. CAB International, Wallingford, UK.
- Herridge D F and Betts J H 1988 Field evaluation of soybean genotypes selected for enhanced capacity to nodulate and fix nitrogen in the presence of nitrate. Plant and Soil 110, 129-135.
- Herridge D F and Brockwell J 1988 Contribution of fixed nitrogen and soil nitrate to the nitrogen economy of irrigated soybean. Soil Biol Biochem. 20, 711-717.
- Hodges T and French V 1985 Soyphen: Soybean growth stages modeled from temperature, daylength, and water availability. Agron. J. 77, 500-505.
- Huke R E 1982 Rice area by type of culture: South, Southeast, and East Asia. International Rice Research Institute, Los Baños, Philippines. 32 p.
- Imsande J 1986 Ineffective utilization of nitrate by soybean during pod fill. Physiol. Plant. 68, 689-694.
- Imsande J and Edwards D G 1988 Decreased rates of nitrate uptake during pod fill by cowpea, green gram, and soybean. Agron. J. 80, 789-793.
- Imsande J 1988 Interrelationship between plant development stage, plant growth rate, nitrate utilization and nitrogen fixation in hydroponically grown soybean. J. Exp. Bot. 39, 775-785.
- Imsande J 1989 Rapid dinitrogen fixation during soybean pod fill enhances net photosynthetic output and seed yield: A new perspective. Agron. J. 81, 549-556.
- Janssen G and Metzger W H 1928 Transformation of nitrogen in rice soil. J. Am. Soc. Agron. 20, 459-476.
- John P S, Pandey R K, Buresh R J and Prasad R 1989 Lowland rice response to urea following three cowpea cropping systems. Agron. J. 81, 853-857.
- Keeney D 1986 Sources of nitrate to groundwater. CRC Critical Reviews in Environ. Control 16, 257-304.

- Khind C S, Josan A S and Beri V 1985 Nitrogen release from sesbania green manure and effect of time of application of N fertilizer on lowland rice. *Int. Rice Res. Newsl.* 10, 26–27.
- Knowles 1982 Denitrification. *Microbiol. Rev.* 46, 43–70.
- Koyama T and App A 1979 Nitrogen balance in flooded rice soils. *In Nitrogen and Rice*. pp 95–104. International Rice Research Institute, Los Baños, Philippines.
- Kucey R M N, Chaiwanakupt P, Arayangkool T, Snitwongse P, Siripaibool C, Wadisirisuk P and Boonkerd N 1988 Nitrogen fixation ( $^{15}\text{N}$  dilution) with soybeans under Thai field conditions. II. Effect of herbicides and water application schedule. *Plant and Soil* 108, 87–92.
- Kulkarni K R and Pandey R K 1988 Annual legumes for food and as green manure in a rice-based cropping system. *In Sustainable Agriculture: Green Manuring in Rice Farming*. pp 289–299. International Rice Research Institute, Los Baños, Philippines.
- Ladha J K, Garcia M, Miyan S, Padre A T and Watanabe I 1989a Survival of *Azorhizobium caulinodans* in the soil and rhizosphere of wetland rice under *Sesbania rostrata*-rice rotation. *Appl. Environ. Microbiol.* 55, 454–460.
- Ladha J K, Miyan S and Garcia M 1989b *Sesbania rostrata* as a green manure for lowland rice: Growth,  $\text{N}_2$  fixation, *Azorhizobium* sp. Inoculation, and effects on succeeding crop yields and N balance. *Biol. Fert. Soils* 7, 191–197.
- Ladha J K, Pareek R P and Becker M 1992 Stem-nodulating legume-Rhizobium symbiosis and its agronomic use in lowland rice. *Adv. Soil Sci.* (*In press*).
- Lawn R J and Williams J H 1987 Limits imposed by climatological factors. *In Food Legume Improvement for Asian Farming Systems: Proceedings of an International Workshop*, Khon Kaen, Thailand, September 1986. ACIAR Proc. Ser. No. 18. Eds. E S Wallis and D E Byth. pp. 83–98.
- Legg J O and Meisinger J J 1982 Soil nitrogen budgets. *In Nitrogen in Agricultural Soils*. Ed. F J Stevenson. Agron. Monograph No. 22 pp 503–566. Am. Soc. Agron. Madison, WI.
- Letej J, Valoras N, Focht D D and Ryden J C 1981 Nitrous oxide production and reduction during denitrification as affected by redox potential. *Soil Sci. Soc. Am. J.* 45, 727–730.
- Macdonald A J, Powlson D S, Poulton P R and Jenkinson D S 1989 Unused fertiliser nitrogen in arable soils – Its contribution to nitrate leaching. *J. Sci. Food Agric.* 46, 407–419.
- Magdoff F R 1982 Soil nitrogen availability under grassland and cultivated corn. *Plant and Soil* 68, 395–398.
- McEwen J, Darby R J, Hewitt M V and Yeoman D P 1989 Effects of field beans, fallow, lupins, oats, oilseed, rape, peas, ryegrass, sunflowers and wheat on nitrogen residues in the soil and on the growth of a subsequent wheat crop. *J. Agric. Sci. Camb.* 115, 209–219.
- Meade J, Higgins P and O'Gara F 1985 Studies on the inoculation and competitiveness of a *Rhizobium leguminosarum* strain in soils containing indigenous rhizobia. *Appl. Environ. Microbiol.* 49, 899–903.
- Minchin F R and Summerfield R J 1976 Symbiotic nitrogen fixation and vegetative growth of cowpea (*Vigna unguiculata* (L.) Walp.) in waterlogged conditions. *Plant and Soil* 45, 113–127.
- Minchin F R, Summerfield R J, Eaglesham A R J and Stewart K A 1978 Effects of short-term waterlogging on growth and yield of cowpea. *J. Agric. Sci. Camb.* 90, 335–336.
- Morris R A, Furoc R E and Dizon M A 1986a Rice responses to a short duration green manure. I. Grain yield. *Agron. J.* 78, 409–412.
- Morris R A, Furoc R E and Dizon M A 1986b Rice responses to a short duration green manure. II. N recovery and utilization. *Agron. J.* 78, 413–416.
- Morris R A, Furoc R E, Rajbhandari N K, Marqueses E P and Dizon M A 1989 Rice response to waterlog-tolerant green manures. *Agron. J.* 81, 803–809.
- Mosier A R, Guenzi W D and Schweizer E E 1986 Soil losses of dinitrogen and nitrous oxide from irrigated crops in Northeastern Colorado. *Soil Sci. Soc. Am. J.* 50, 344–348.
- Muchow R C 1985 Phenology, seed yield, and water use of grain legumes grown under different soil water regimes in a semi-arid tropical environment. *Field Crops Res.* 11, 81–97.
- Munns D N 1977 Mineral nutrition of the legume symbiosis. *In A Treatise on Dinitrogen Fixation. IV. Agronomy and Ecology*. Eds. R W F Hardy and A H Gibson. pp 353–391. Wiley, New York.
- Myers R J K and Wood I M 1987 Food legumes in the nitrogen cycle of farming systems. *In Food Legume Improvements for Asian Farming Systems: Proceedings of an International Workshop*, Khon Kaen, Thailand, September 1986. ACIAR Proc. Ser. No. 18. Eds. E S Wallis and D E Byth. pp 46–52.
- Nagarajah S, Neue H U and Alberto M C R 1989 Effect of *Sesbania*, *Azolla* and rice straw incorporation on the kinetics of  $\text{NH}_4$ , K, Fe, Mn, Zn and P in some flooded rice soils. *Plant and Soil* 116, 37–48.
- Ndoye I and Dreyfus B 1988  $\text{N}_2$  fixation by *Sesbania rostrata* and *Sesbania sesban* estimated using  $^{15}\text{N}$  and total N difference methods. *Soil Biol. Biochem.* 20, 209–213.
- Nelson R S, Ryan S A and Harper J E 1983 Soybean mutants lacking constitutive nitrate reductase. I. Selection and initial plant characterization. *Plant Physiol.* 72, 503–509.
- Norman R J, Gilmour J T and Wells B R 1990 Mineralization of nitrogen from nitrogen-15 labeled crop residues and utilization by rice. *Soil Sci. Soc. Am. J.* 54, 1351–1356.
- Ofori F, Pate J S and Stern W R 1987 Evaluation of  $\text{N}_2$  fixation and nitrogen economy of a maize/cowpea inter-crop system using  $^{15}\text{N}$  dilution methods. *Plant and Soil* 102, 149–160.
- O'Hara G W, Boonkerd N and Dilworth M J 1988 Mineral constraints to nitrogen fixation. *Plant and Soil* 108, 93–110.
- Palaniappan S P 1985 *Cropping Systems in the Tropics*. Wiley Eastern Ltd., New Delhi, India. 215 p.
- Palm O, Weerakoon W L, De Silva M A P and Rosswall T 1988 Nitrogen mineralization of *Sesbania sesban* used as green manure for lowland rice in Sri Lanka. *Plant and Soil* 108, 201–209.
- Pareek R P, Ladha J K and Watanabe I 1990 Estimating  $\text{N}_2$  fixation by *Sesbania rostrata* and *S. cannabina* (syn. *S. aculeata*) in lowland rice soil by the  $^{15}\text{N}$  dilution method. *Biol. Fert. Soils* 10, 77–88.

- Patrick Jr W H and Wyatt R 1964 Soil nitrogen loss as a result of alternate submergence and drying. *Soil Sci. Soc. Am. Proc.* 28, 647-653.
- Peoples M B and Craswell E T 1992 Biological nitrogen fixation: Investments, expectations and actual contributions to agriculture. *Plant and Soil* 141, 13-39.
- Peoples M B and Herridge D F 1990 Nitrogen fixation by legumes in tropical and subtropical agriculture. *Adv. Agron.* 44, 156-223.
- Ponnampuruma F N 1985 Chemical kinetics of wetland rice soils relative to soil fertility. In *Wetland Soils Characterization, Classification, and Utilization*, pp 71-80. International Rice Research Institute, Los Baños, Philippines.
- Pretty J N and Conway G R 1989 Agriculture as a global polluter. Gatekeeper Series No. SA11, International Institute for Environment and Development, London, UK. 15 p.
- Reddy K R and Patrick W H Jr 1975 Effect of alternate aerobic and anaerobic conditions on redox potential, organic matter decomposition and nitrogen loss in a flooded soil. *Soil Biol. Biochem.* 7, 87-94.
- Rerkasem K and Rerkasem B 1988 Yields and nitrogen nutrition of intercropped maize and ricebean (*Vigna umbellata* [Thunb.] Ohwi and Ohashi). *Plant and Soil* 108, 151-162.
- Rerkasem B, Rerkasem K, Peoples M B, Herridge D F and Bergersen F J 1988 Measurement of  $N_2$  fixation in maize (*Zea mays* L.)-ricebean (*Vigna umbellata* [Thunb.] Ohwi and Ohashi) intercrops. *Plant and Soil* 108, 125-135.
- Robson A D 1978 Mineral nutrients limiting nitrogen fixation in legumes. In *Mineral Nutrition of Legumes in Tropical and Subtropical Soils*. Eds. C S Andrew and E J Kamprath. pp 277-293. CSIRO, Melbourne, Australia.
- Roger P A and Ladha J K 1992 Biological  $N_2$  fixation in wetland rice fields: Estimation and contribution to nitrogen balance. *Plant and Soil* 141, 41-55.
- Sahrawat K L 1980 Soil and fertilizer nitrogen transformations under alternate flooding and drying moisture regimes. *Plant and Soil* 55, 225-233.
- Sahrawat K L 1983 Mineralization of soil organic nitrogen under waterlogged conditions in relation to other properties of tropical rice soils. *Aust. J. Soil. Res.* 21, 133-138.
- Santiago-Ventura T, Bravo M, Daez C, Ventura V, Watanabe I and App A A 1986 Effects of N-fertilizers, straw, and dry fallow on the nitrogen balance of a flooded soil planted with rice. *Plant and Soil* 93, 405-411.
- Savant N K, De Datta S K and Craswell E T 1982 Distribution patterns of ammonium nitrogen and N uptake by rice after deep placement of urea supergranules in wetland soil. *Soil Sci. Soc. Am. J.* 46, 567-573.
- Seneviratne R and Wild A 1985 Effect of mild drying on the mineralization of soil nitrogen. *Plant and Soil* 84, 175-179.
- Senthong C and Pandey R K 1989 Response of five food legume crops to an irrigation gradient imposed during reproductive growth. *Agron. J.* 81, 680-686.
- Sharma K N, Bhandari A L, Kapur M L and Rana D S 1985 Influence of growing various crops in five different fixed rotations on the changes in nitrate and total nitrogen content of soils. *J. Agric. Sci. Camb.* 104, 609-613.
- Sinclair T R, Muchow R C, Bennett J M and Hammond L C 1987 Relative sensitivity of nitrogen and biomass accumulation to drought in field-grown soybean. *Agron. J.* 79, 986-991.
- Singh Y, Khind C S and Singh B 1991 Efficient management of leguminous green manures in wetland rice. *Adv. Agron.* 45, 135-189.
- Singleton P W and Bohlool B B 1983 The effect of salinity on the functional components of the soybean-*Rhizobium japonicum* symbiosis. *Crop Sci.* 23, 815-818.
- Singleton P W, AbdelMagid H M and Tavares J W 1985 Effect of phosphorus on the effectiveness of strains of *Rhizobium japonicum*. *Soil Sci. Soc. Am. J.* 49, 613-616.
- Singleton P W and Tavares J W 1986 Inoculation response of legumes in relation to the number and effectiveness of indigenous rhizobium populations. *Appl. Environ. Microbiol.* 51, 1013-1018.
- Sisworo W H, Mitrosuhardjo M M, Rasjid H and Myers R J K 1990 The relative roles of N fixation, fertilizer, crop residues and soil in supplying N in multiple cropping systems in a humid, tropical upland system. *Plant and Soil* 121, 73-82.
- Smith D L, Dijk M and Hume D J 1988 The effect of water deficit on nitrogen fixation (acetylene reduction) by white bean and soybean. *Can. J. Plant Sci.* 68, 957-968.
- Ssali H and Keya S O 1986 The effects of phosphorus and nitrogen fertilizer level on nodulation, growth and dinitrogen fixation of three bean cultivars. *Trop. Agric. (Trinidad)* 63, 105-109.
- Streeter J 1988 Inhibition of legume nodule formation and  $N_2$  fixation by nitrate. *CRC Critical Rev. Plant Sci.* 7, 1-23.
- Strickland R W 1969 Fluctuations in available mineral nitrogen in a flooded rice soil on the sub-coastal plains of the Adelaide River, N.T. *Aust. J. Exp. Agric. Anim. Husb.* 9, 532-540.
- Strong W M, Harbison J, Nielson R G H, Hall B D and Best E K 1986 Nitrogen availability in a Darling Downs soil following some cereal, oilseed and grain legume crops. 1. Soil nitrogen accumulation. *Aust. J. Exp. Agric.* 26, 347-351.
- Suwanarit A, Suwanarit C and Chotechaungmanirat S 1986 Quantities of fixed N and effects of grain legumes on following maize, and N and P status of soil as indicated by isotopes. *Plant and Soil* 93, 249-258.
- Thies J E, Singleton P W and Bohlool B B 1991a Influence of the size of indigenous rhizobial populations on establishment and symbiotic performance of introduced rhizobia on field-grown legumes. *Appl. Environ. Microbiol.* 57, 19-28.
- Thies J E, Singleton P W and Bohlool B B 1991b Modeling symbiotic performance of introduced rhizobia in the field by use of indices of indigenous population size and nitrogen status of the soil. *Appl. Environ. Microbiol.* 57, 29-37.
- Thurlow D L and Hiltbold A E 1985 Dinitrogen fixation by soybeans in Alabama. *Agron. J.* 77, 432-436.
- Timsina J 1989 Adaptation of diverse cowpea (*Vigna unguiculata* (L.) Walp.) cultivars to excess moisture and drought in riceland. Ph.D. Thesis, University of the Philippines, Los Baños, Philippines.
- Torres R O, Magbanua R D and Garrity D P 1988 Evaluation of cowpeas as legume component in acid upland rice-based cropping systems. *Philipp. J. Crop Sci.* 13, 91-98.
- Ventura W and Watanabe I 1978 Dry season soil conditions

- and soil nitrogen availability to wet season wetland rice. *Soil Sci. Plant Nutr.* 24, 535–545.
- Ventura W, Mascarina G B, Furoc R E and Watanabe I 1987 Azolla and Sesbania as biofertilizers for lowland rice. *Philipp. J. Crop Sci.* 12, 61–69.
- Vincent J M 1980 Factors controlling the legume-Rhizobium symbiosis. In *Nitrogen Fixation. II*. Eds. W E Newton and W H Orme-Johnson. pp 103–120. University Park Press, Baltimore, MD.
- Weaver R W and Frederick L R 1974 Effect of inoculum rate on competitive nodulation of *Glycine max* L. Merrill. II. Field studies. *Agron. J.* 66, 233–236.
- Weaver R W, Morris D R, Boonkerd N and Sij J 1987 Populations of *Bradyrhizobium japonicum* in fields cropped with soybean rice rotation. *Soil Sci. Soc. Am. J.* 51, 90–92.
- White R E 1988 Leaching. In *Advances in Nitrogen Cycling in Agricultural Ecosystems*. Ed. J R Wilson. pp 193–211. C.A.B. International, UK.
- Wallihan E F 1938 The effect of water logging on the rate of nitrification in soils subsequent to drainage. *Soil Sci. Soc. Am. Proc.* 2, 259–262.
- Yamane I 1957 Nitrate reduction and denitrification in flooded soils. *Soil Plant Food* 3, 100–103.
- Yoshida T and Yoneyama T 1980 Atmospheric dinitrogen fixation in the flooded rice rhizosphere determined by the N-15 isotope technique. *Soil Sci. Plant Nutr.* 26, 551–559.
- Yost R and Evans D 1988 Green manure and legume covers in the tropics. Hawaii Institute of Tropical Agriculture and Human Resources Research Series 055. College of Tropical Agriculture and Human Resources, University of Hawaii. 44 p.
- Zapata F, Danso S K A and Fried M 1987 Time course of nitrogen fixation in field grown soybean using nitrogen-15 methodology. *Agron. J.* 79, 172–176.

23