

1999 UKRAINE REPRODUCTIVE HEALTH SURVEY

Final Report

September 2001



KIEV INTERNATIONAL
INSTITUTE OF SOCIOLOGY



SAFER • HEALTHIER • PEOPLE™

DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention



U.S. Agency for
International Development

1999 UKRAINE

REPRODUCTIVE HEALTH SURVEY

FINAL REPORT

1999 UKRAINE
REPRODUCTIVE HEALTH SURVEY
FINAL REPORT

Kiev International Institute of Sociology

**Centers for Disease Control and Prevention,
Division of Reproductive Health, USA**

United States Agency for International Development

September, 2001

ACKNOWLEDGMENTS

This report describes the 1999 Ukraine Reproductive Health Survey and its major findings. This undertaking could not have taken place successfully without the participation, support, and cooperation of a large number of individuals and organizations, both in Ukraine and the United States. Funding for the survey was provided by the United States Agency for International Development (USAID) mission in Kiev, which was also the motivating force behind the activity. The survey was capably directed by Dr. Vladimir Paniotto, president of the Kiev International Institute of Sociology (KIIS). KIIS staff members in Kiev and elsewhere in Ukraine, particularly Elena Buslayeva and Victoria Zakhozha, did an outstanding job coordinating and implementing all survey activities. Dr. Howard Goldberg coordinated technical assistance from the United States Centers for Disease Control and Prevention's Division of Reproductive Health, which provided consultation in survey design, planning, questionnaire development, fieldwork implementation, data analysis, and report preparation. Important contributions were also made by many others working in the area of reproductive health in Ukraine, most notably at the Ukraine Ministry of Health and Johns Hopkins University/Population Communications Services. Finally, thanks go to the KIIS interviewing staff, who did an excellent job collecting the survey data, and to the many women across Ukraine who willingly gave their time and cooperation to provide the information that will be so valuable to improving the health of women and children in their country.

Contributors

This report was prepared by Howard Goldberg, Natalia Melnikova, Elena Buslayeva, and Victoria Zakhosha. The following organizations and individuals made important contributions to the 1999 Ukraine Women's Reproductive Health Survey and the preparation of this report:

Kiev International Institute of Sociology (KIIS)

Vladimir Paniotto, President, Survey Director
Elena Buslayeva, Assistant Director, KIIS
Victoria Zakhosha, Project Manager

Centers for Disease Control and Prevention (USA)

Howard Goldberg, Deputy Chief, Behavioral Epidemiology and Demographic
Research Branch (BEDRB), Division of Reproductive Health (DRH)
Natalya Melnikova, Visiting Fellow, BEDRB, DRH
Jay Friedman, Program Analyst, BEDRB, DRH
Leo Morris, Chief, BEDRB, DRH
David Gilbertz, Computer Programmer, DRH
J. Timothy Johnson, Chief, Program Services Evaluation Section, DRH

United States Agency for International Development/Kiev

Teresa Ingham, Women's Health Advisor (former), Office of Democratic and Social
Transition (ODST)
Marilynn Schmidt, Director, ODST
Catherine Fischer, Senior Health Advisor (former), ODST
Pamela Mandel, Deputy Director, ODST
Alina Yurova, Health Program Coordinator, ODST
Eliot Pearlman, Senior Health Advisor, ODST
Boris Uspensky, Women's Health Advisor, ODST

Johns Hopkins University/Population Communications Services (JHU/PCS)

Laurie Liskin, Chief, Division of Europe and New Independent States (DENIS)
Paul Nary, Program Officer, DENIS
Anne Sochan, Project Officer, DENIS

The Futures Group, Inc.

Monica Medrek, Country manager, POLICY Project
Karen Foreit, Evaluation Coordinator, POLICY Project

Others

Petr Velebil, Institute for the Care of Mother and Child, Prague, Czech Republic

Naomi Blumberg, Fellow, Office of Population, USAID/Washington

Tamara K. Irkina, Deputy Head of Administration, Head of Department of Maternal and
Child Health, Ukraine Ministry of Health

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGMENTS	i
CONTRIBUTORS	ii
LIST OF TABLES	viii
LIST OF FIGURES	xv
SUMMARY OF FINDINGS	1
CHAPTER I. INTRODUCTION	11
Background	
The Ukraine Women’s Reproductive Health Initiative	
The 1999 Ukraine Reproductive Health Survey	
CHAPTER II. SURVEY METHODOLOGY	15
Organizational Structure	
Questionnaire content	
Survey design	
Data collection	
Response rates	
CHAPTER III. CHARACTERISTICS OF THE RESPONDENTS	19
Demographic and social characteristics	
Economic characteristics	
Marriage	
CHAPTER IV. PREGNANCY AND CHILDBEARING	31
Fertility and pregnancy levels and patterns	
Pregnancy outcomes	
Planning status of pregnancies	
Pregnancy intentions	
CHAPTER V. INDUCED ABORTION	55
Proportions of women with any abortions	
Current incidence of abortion	
Types of abortion, complications, cost	

CHAPTER VI. INFERTILITY.....	71
Prevalence of infertility	
Treatment of infertility	
CHAPTER VII. CONTRACEPTION.....	79
Knowledge and ever use of contraceptive methods	
Current contraceptive prevalence	
Recent trends in contraceptive prevalence	
Source of contraceptive methods	
Reasons for not using contraception	
Unmet need for contraception	
Preference for other methods/Problems with current method	
Contraceptive failure and discontinuation	
Opinions about fertility control methods	
Use of non-supplied methods	
IUD use	
Oral contraceptive use	
Contraceptive sterilization	
CHAPTER VIII. CONTRACEPTIVE COUNSELING.....	127
Post-abortion/Post-partum counseling	
Content of counseling/Method selection	
Partner involvement in reproductive health decisions	
CHAPTER IX. INFORMATION, EDUCATION, AND COMMUNICATION ...	135
Television viewing habits	
Radio listening habits	
Newspaper readership	
Exposure to and attitudes about health messages in the media	
CHAPTER X. SEXUAL EXPERIENCE.....	149
Sexual experience of all women	
First sexual experience of young adults	
CHAPTER XI. MATERNAL AND CHILD HEALTH / WOMEN'S HEALTH...	165
Prenatal care	
Hospitalization during pregnancy	
Labor and delivery	
Cigarette smoking during pregnancy	
Breastfeeding	
Women's health behaviors	

CHAPTER XII. SEXUALLY TRANSMITTED INFECTIONS.....	187
Awareness of STI's and lifetime history of STI diagnosis	
Treatment for possible symptoms of STIs	
Knowledge about prevention/Perceived risk of acquiring STIs	
CHAPTER XII. DOMESTIC VIOLENCE.....	199
CHAPTER XIII. CONCLUSIONS.....	207
REFERENCES.....	211
APPENDIX. Survey Questionnaire.....	215

LIST OF TABLES

Chapter II

- 2.1 Percentage distribution of household and individual final interview status, by residence

Chapter III

- 3.1 Percentage distributions of socio-demographic characteristics of respondents, by residence
- 3.2 Percentage distributions of socio-demographic characteristics of respondents, by region of residence
- 3.3 Percentage distributions of current employment status and percent of women who live in homes with selected possessions, by residence
- 3.4 Percentage distributions of current marital status by age of respondent, by residence
- 3.5 Percent of respondents who have ever been in union and who are currently in union, by current age
- 3.6 Percent of women who were married or in a consensual union before selected ages by current age

Chapter IV

- 4.1 Percentage of respondents who have had one or more live births and mean number of live births, by age at interview, by residence
- 4.2 Percentage of respondents who have had one or more pregnancies and mean number of pregnancies, by age at interview, by residence
- 4.3 Age-specific and total fertility rates, by residence, region, and education
- 4.4 Age-specific and total pregnancy rates, by residence, region, and education
- 4.5 Percent of women ever in union with any live births and percent ever pregnant, by years since first union and residence
- 4.6 Percentage distribution of number of live births, by current age and residence
- 4.7 Percentage distribution of number of live births, by number of years since first union, by residence
- 4.8 Percentage of women who had a live birth before selected age, by current age
- 4.9 Percentage distribution of number of pregnancies, by current age, by residence
- 4.10 Percentage distributions of outcomes of pregnancies ending since January 1996, by selected characteristics
- 4.11 Percentage distribution of planning status of pregnancies ending since January 1996, by selected characteristics
- 4.12 Percentage distribution of outcomes of pregnancies ending since January 1996,
- 4.13 Number of additional children desired, by number of living children, by residence among fecund women currently in union
- 4.14 Percentage distribution of total number of children desired, by number desired at time of first marriage, by residence, fecund women currently in union

Chapter V

- 5.1 Percentage of respondents who have had any induced abortions or two or more induced abortions and mean number of induced abortions, by age at interview, by residence
- 5.2 Percentage of respondents who have had any induced abortions or two or more induced abortions and mean number of induced abortions, by selected characteristics
- 5.3 Percentage distribution of number of abortions, by current age, by residence
- 5.4 Age-specific abortion rates and other measures of induced abortion, by residence
- 5.5 Selected measures of induced abortion, by region and education
- 5.6 Total fertility, pregnancy, and induced abortion rates, by selected characteristics
- 5.7 Percentage distribution of types of abortions, by selected characteristics, all abortions since January 1994
- 5.8 Percentage of induced abortions since January 1994 resulting in complications requiring medical treatment, by type of abortion, by residence
- 5.9 Percentage distribution of primary reason for abortion since January 1994, by age at abortion, region, residence, and education
- 5.10 Percentage distribution of payment for most recent abortion since January 1997, by age at abortion, region, residence, and education
- 5.11 Percentage of women who received local anesthesia for their most recent abortion since January 1994, by year, type, age, region, residence, and education

Chapter VI

- 6.1 Percentage of women who have ever had problem becoming pregnant in the previous 10 years, by duration of problem, by age, region, residence, and education
- 6.2 Percentage of women with infertility problems who sought treatment, by age, region, residence, education and length of problem
- 6.3 Percentage of women/couples who went to various types of place for treatment of infertility problems, by age, residence, and education
- 6.4 Percentage of treated women/couples who received various types of treatment for infertility problems, by age, residence, and education
- 6.5 Percentage of women treated for infertility problems who eventually become pregnant and percent who had a live birth, according to characteristics, length of problem, and where treated

Chapter VII

- 7.1 Percentage of respondents who heard of specific contraceptive methods, by residence and marital status
- 7.2 Percentage of respondents who heard of specific contraceptive methods, by age and education
- 7.3 Percentage of respondents who know where to obtain specific contraceptive methods, by residence, marital status, and education
- 7.4 Percentage of respondents who have ever used specific contraceptive methods, by residence, marital status, and education
- 7.5 Percentage distribution of current contraceptive method by respondent's age, women in union.

- 7.6 Percentage distribution of current contraceptive method by number of living children, women in union
- 7.7 Percentage distribution of current contraceptive method by type of place of residence and for oversampled oblasts, women in union
- 7.8 Percentage distribution of current contraceptive method by region of residence, women in union
- 7.9 Percentage distribution of current contraceptive method by educational level, women in union
- 7.10 Percentage distribution of current contraceptive method, by marital status
- 7.11 Percent of All 15-39 Year-Old Respondents Reporting Current Use of Any Contraception or of Modern Contraception at Six Month Intervals from January 1994 to April 1999
- 7.12 Percentage distribution of source of selected contraceptive methods, by residence, among current users
- 7.13 Percentage distribution of payment for contraceptive method for current users of oral contraceptives, IUD, and condoms
- 7.14 Percentage distribution of primary reason for not using contraception, by union status
- 7.15 Percent of women with unmet need for family planning, according to two definitions by age, region, residence, and education
- 7.16 Percentage distributions of users of selected contraceptive methods according to whether they prefer to use a different method, by current and preferred method
- 7.17 Percentage distribution of reason not using preferred method of contraception, by residence
- 7.18 Percentage distribution of major problems or concerns with current contraceptive method, by current method
- 7.19 Contraceptive failure rates after one, two, and three years for selected methods of contraception, contraceptive episodes starting since January 1994
- 7.20 Contraceptive discontinuation rates (all reasons) after one, two, and three years for selected methods of contraception, contraceptive episodes starting since January 1994
- 7.21 One-year contraceptive discontinuation rates for selected reasons for selected contraceptive methods, contraceptive episodes starting since January 1994
- 7.22 Primary reason for discontinuing selected contraceptive methods, contraceptive episodes starting since January 1994 (Percentage Distributions)
- 7.23 Percent of women with poor opinion of selected aspects of various contraceptive methods, by residence, according to use of that method
- 7.24 Percentage of users of traditional methods who said that selected factors were important in their decision not to use a modern method, by residence.
- 7.25 Percentage distribution of perceived effectiveness of modern methods compared to method currently used, by residence, current users of traditional methods
- 7.26 Percentage distribution of time of most recent IUD insertion and length of time provider said IUD could be worn, by residence, women with IUD insertion since January 1994
- 7.27 Percentage distribution of the most serious problem women have had related to their IUD, by residence, women with insertion since January 1994

- 7.28 Percentage distribution of length of time physician told women they could take oral contraceptives, by residence, women who started taking oral contraceptives since January 1994
- 7.29 Percentage distribution of the most serious problem women have had related to oral contraceptives, by residence, women who did take oral contraceptives since January 1994
- 7.30 Percent of fecund in union respondents wanting no more children, who are interested in being contraceptively sterilized, by selected characteristics
- 7.31 Percentage distribution of primary reason not interested in contraceptive sterilization, by residence

Chapter VIII

- 8.1 Percent of women who received various family planning services after their most recent delivery or abortion, by residence among women who had a delivery or abortion since January 1994
- 8.2 Percent of women starting use of IUD, oral contraceptives, or injections since January 1994 with whom family planning provider discussed contraceptive options and percentage distributions of who selected method
- 8.3 Percent of women starting use of IUD, oral contraceptives, or injections since January 1994 to whom provider explained about possible side effects and percent of women receiving counseling who 1) understood most information of side effects; 2) were told when to return for follow-up, refill, etc.; and 3) were given a pelvic examination, by region, residence, and method
- 8.4 Percent of women in union who have discussed use of family planning with their partner and percentage distribution of who women think should make decisions about whether to use contraception, by selected characteristics
- 8.5 Percent of women whose partner knew about her most recent abortion and percent who discussed with her partner whether to have an abortion, by selected characteristics

Chapter IX

- 9.1 Percentage distribution of television viewing frequency, by age, region, residence, and education
- 9.2 Percentage of respondents who watch various types of television programs, by age, region, residence, and education
- 9.3 Percentage of respondents who watch television at various times, by age, region, residence, and education
- 9.4 Percentage distribution of frequency of radio listening, by age, region, residence, and education
- 9.5 Percentage of respondents who listen to various types of radio programs distribution of radio programs, by age, region, residence, and education
- 9.6 Percentage of respondents who listen to the radio at various times, by age region, residence, and education
- 9.7 Percentage distribution of frequency of reading daily newspapers by age, region, residence, and education

- 9.8 Percentage of women who read specific newspapers, by age, region, residence, and education
- 9.9 Percent of women who had seen any messages on TV or read anything in newspapers or magazines about family planning within the previous six months, and percent who think such important information should be broadcast, by selected characteristics
- 9.10 Percentage distribution of current contraceptive use, by whether women had been exposed to family planning (FP) messages in the media, among fecund sexually active women

Chapter X

- 10.1 Percent of women who had sexual intercourse before ages 16, 18, 20, by current age, by residence
- 10.2 Percentage of respondents who reported having sexual intercourse within selected periods of time, by age, residence, region, and marital status, sexually experienced women
- 10.3 Percentage distribution of frequency of sexual intercourse in the previous 30 days, by age, by current marital status, among sexually experienced women
- 10.4 Percent of respondents between the ages of 15 and 24 who have ever had sexual intercourse, by current age, by residence
- 10.5 Percent of respondents between the ages of 15 and 24 who have ever had premarital sexual intercourse, by current age, by residence
- 10.6 Percentage distribution of relationship to first sexual partner, by age at first sex, by residence 15-24 year-old sexually experienced women
- 10.7 Percentage distributions of contraceptive method used at first sexual intercourse, by age at first sex, by residence, 15-24 year-old women with premarital sexual experience
- 10.8 Percentage distribution of numbers of sexual partners in the previous 12 months, by age, residence, and marital status, sexually experienced women

Chapter XI

- 11.1 Percentage distribution of trimester of first prenatal care visit, by selected characteristics, all pregnancies ending in live birth since January 1994
- 11.2 Percentage distribution of number of prenatal care visits, by selected characteristics, all pregnancies with prenatal care ending in live births since January 1994
- 11.3 Percentage distribution of place of (most) prenatal care visits, by residence and region, all pregnancies ending in live birth since January 1994
- 11.4 Percent of respondents who had their blood pressure (BP) measured and percent who received an ultrasound during their most recent pregnancy ending in a live birth since January 1994, by selected characteristics
- 11.5 Percent of respondents who were told they had high blood pressure (BP), percent who were told they had anemia, and percent who took iron supplements during their most recent pregnancy ending in a live birth since January 1994, by selected characteristics
- 11.6 Percent of pregnancies ending in a live birth since January 1994 during which women were hospitalized, by residence, region, and year of delivery

- 11.7 Percentage distribution of length of hospitalization by residence, region, and year of delivery, pregnancies ending in a live birth since January 1994
- 11.8 Percentage distribution of type of place of delivery for pregnancies ending in a live birth since January 1994, by selected characteristics
- 11.9 Percentage of respondents who received a post partum check-up within six weeks of birth, by selected characteristics, by year of birth, for most recent birth since January 1994
- 11.10 Percentage of women who smoked cigarettes at the time they became pregnant and percentage who continued smoking during pregnancy, by selected characteristics, most recent pregnancy ending in a live birth since January 1994
- 11.11 Percent of babies born to respondents since January 1994 who were breastfed, and mean duration of breastfeeding for those ever breastfed, by selected characteristics
- 11.12 Percentage distribution of current breastfeeding status, according to type of place of residence, by age for infants under one year of age at the time of interview
- 11.13 Percentage distribution of number of cigarettes typically smoked per day, by selected characteristics
- 11.14 Percentage distribution of years since most recent (non-pregnancy related) gynecologic exam, by age, by residence, and by marital status

Chapter XII

- 12.1 Percentage of respondents who have never heard of selected STIs / conditions, by region, residence, age, education, and sexual experience
- 12.2 Percentage of respondents who have ever been diagnosed with selected STIs /conditions, by region, residence, age, and education
- 12.3 Percent of sexually active women who have experienced selected symptoms potentially related to sexually transmitted infections (STI) during the past 12 months, by age, education, and place of residence
- 12.4 Percentage distribution of place of treatment for possible symptoms of STI, by selected characteristics
- 12.5 Percentage distribution of reasons for not seeking treatment for possible STI symptoms, by place of residence, and union status
- 12.6 Percent of respondents who are aware that individuals can be infected with HIV/AIDS or have other sexually transmitted infections (STIs) without showing any signs or symptoms of disease, by selected characteristics
- 12.7 Percentage distribution of opinions regarding the protection that condoms gives against HIV and other STI's, by residence, region, age, education, and sexual experience
- 12.8 Percentage distribution of respondents' perceived risk of sexually transmitted infection, by residence, region, age, education, and number of recent sexual partners

Chapter XIII

- 13.1 Percent of respondents who recall parents/step-parents physically abusing each other and percent who were physically abused as children by someone in their household, by selected characteristics

- 13.2 Percentage of ever married respondents subjected to particular types of violence ever and during the past 12 months
- 13.3 Percentage of respondents abused in past 12 months who sustained injuries, sought treatment, or were hospitalized, by selected characteristics
- 13.4 Percentage of respondents abused in past 12 months who discussed these incidents with various types of people, by selected characteristics
- 13.5 Percent of respondents abused in past 12 months whose partner had been drinking alcohol the last time he abused her, and percent whose partner had been drinking at any time in the past year when he abused her, by selected characteristics

LIST OF FIGURES

- 3.1 Percentage distributions of current marital status, by age of respondent
- 4.1 Total fertility rates, most recent available, selected countries
- 4.2 Age-specific fertility rates, by place of residence
- 4.3 Percentage distributions of pregnancy outcomes since January 1994, by number of living children
- 4.4 Percentage distributions of planning status of pregnancies since January 1994, by number of living children
- 4.5 Percent of fecund women in union who want no more children, according to number of living children
- 5.1 Age-specific abortion rates, by place of residence
- 7.1 Percentage distribution of current contraceptive method among women in union
- 7.2 Percentage distribution of current contraceptive use among women in union, by number of living children
- 7.3 Percentage point change since January 1994 in percent of all 15-39 year-old women currently using contraception
- 7.4 One-year failure and discontinuation rates for selected contraceptive methods
- 7.5 Percent of respondents giving low ratings and for safety/health for selected contraceptive methods
- 10.1 Percent of respondents who have ever had sexual intercourse, by current age
- 10.2 Percent of respondents who ever had sexual intercourse before ages 16, 18, and 20, by current age
- 11.1 Percentage distribution of time of first prenatal visit for the last pregnancy since January 1994
- 11.2 Percentage distribution of length of hospitalization during last pregnancy since January 1994
- 11.3 Percent of respondents reporting selected behaviors/circumstances related to the health of infants/mothers for last pregnancy leading to a live birth since January 1994

1999 Ukraine Reproductive Health Survey

Summary of findings

Introduction

Since the break-up of the Soviet Union in 1991, after which it became an independent state, Ukraine, like the other countries in the region, experienced dramatic and far-reaching political, social, and economic changes. These changes have brought about major transitions in many aspects of life, some positive and some negative. One such area in which substantial change has been taking place is that of the financing and provision of health care and the structure of the Ukrainian health care system.

The transitions that Ukraine has undergone, including those in the area of health care, have no doubt contributed to dramatic changes that have occurred in reproductive health and demographic indicators in recent years. There have been major changes observed in rates of childbearing, abortion, and contraceptive use. There are also unanswered questions about the degree to which there have been changes in such areas as the utilization of maternal-child health services, the prevalence of infertility problems, and maternal behaviors that affect maternal and infant health.

Until recently, relatively little detailed and conclusive information was available about the situation in Ukraine with regard to many important reproductive health topics or the degree to which the reproductive health situation has changed over the past decade, however. The 1999 Ukraine Reproductive Health Survey (URHS) was a population-based, nationwide survey of women of reproductive age carried out from June to October of that year. It constituted the largest and most in-depth data collection effort in the area of reproductive health and related topics since Ukrainian independence.

The URHS was performed in conjunction with the Ukraine Women's Reproductive Health Initiative (UWRHI), a project sponsored by the United States Agency for International Development (USAID). This project consisted of a variety of components, designed to help to reduce maternal morbidity and mortality and improve reproductive health generally in Ukraine. The stated goals of this initiative were to improve the quality of reproductive health services, as well as access to those services, to increase the rate of modern contraceptive use, and to reduce abortion rates in service sites. It was anticipated that improved access to and quality of reproductive health services for women, expanded and improved use of effective contraception, and reduced reliance on abortion as a means of birth prevention would result in reductions in maternal morbidity and mortality.

Seven collaborating agencies took part in the design and implementation of UWRHI activities that were intended to help achieve the following objectives:

- the establishment of demonstration sites for training and delivery of family planning services
- the institutionalization of reproductive health training

- increased public information, education, and communication about family planning
- improved family planning policy environment
- increased supply and distribution of family planning methods
- improved methods of child delivery and maternal care

There were two major reasons for carrying out the 1999 URHS. First, it was intended to fill a substantial need for data regarding the current status of reproductive health of Ukrainian women. Besides providing data on the current situation overall, it also adds to what is known about reproductive health trends and differentials within the population and allows more accurate determinations to be made about the needs of the population. Secondly, the URHS was designed to provide programmatically useful results. The data collected on reproductive health can be used to help direct, modify, or develop interventions, as well as to provide information to policy makers and health care program officials and providers.

Methodology

The 1999 Ukraine Reproductive Health Survey was designed to collect information from a representative sample of all women between the ages of 15 and 44 living in households throughout Ukraine. Although some pregnancy, childbearing, and abortion occurs outside of ages 15 to 44, the relative rarity of these events at those ages in Ukraine suggested that it would be most efficient to limit the sample to women in this age range. The 1999 URHS was implemented by the Kiev International Institute of Sociology (KIIS), which was responsible for selecting the sample of households and individuals, recruiting and training interviewers, conducting field work, processing the data, and performing some of the data analysis. The United States Agency for International Development (USAID) mission in Kiev was the source of all funding for the URHS and was consulted regarding the content of the survey. Technical assistance for all phases of the survey was provided by the Division of Reproductive Health of the United States Centers for Disease Control and Prevention (CDC/DRH).

There were two survey questionnaires: a short household instrument and a much longer individual questionnaire (see Appendix A). The individual form was completed by selected women 15 to 44 years of age who agreed to be interviewed and covered a wide range of topics related to reproductive health status and needs in Ukraine. The sections of the questionnaire were: demographic characteristics; fertility, pregnancy, and abortion; contraception; information, education, and communication (IEC) concerning family planning; women's health; socioeconomic characteristics of respondents; and intimate partner violence.

The survey was designed to obtain interviews with a nationally representative sample of about 7,000 women. The sample was geographically self-weighting, with the exception of substantial over-sampling of two sites--the oblasts of Donetsk and Odessa. Three-stage cluster sampling was used to select survey respondents. Potential respondents consisted of all women between the ages of 15 and 44 years who lived in households anywhere in Ukraine. The first stage of sampling consisted of the selection of approximately 550 primary sampling units (census enumeration districts and postal zones) were selected across Ukraine. The second stage consisted of the selection of

households and women of childbearing age. Field work lasted from June through October of 1999. At least one 15-44 year-old woman was identified as living in 55% of sampled households. Of those women who were selected for interview, 85% were successfully interviewed, 8% were never found at home, and 6% refused to be interviewed.

Characteristics of respondents

Almost three-fourths of respondents were living in areas defined as urban. About half the urban population of women of childbearing age was living in cities of more than half a million people. The survey age distribution closely matched the distribution according to official national estimates. Just over six of every ten respondents were currently living in either a registered or unregistered marriage, 9% were either divorced or separated, 2% were widows, and 23% had never been married or lived with a man. As in most of eastern Europe and the former Soviet Union, Ukrainian women tend to marry at a young age compared to other developed countries. URHS data reveal that most first marriages take place when women are in their early twenties. Only about one-third of women between the ages of 20 and 24 years had never been in a registered or unregistered marriage. Survey data provide no evidence that the age at marriage has been rising in recent years.

Ukrainian women tend to be well educated, as evidenced by the fact that only 11% of respondents had not completed secondary schooling, over two-thirds of women had finished secondary school, but had not received any post-secondary schooling, and one of every five respondents had received some post-secondary education. The largest ethnic-linguistic group consisted of Ukrainians who most often spoke Ukrainian (38%), but there were also large numbers of Ukrainians who usually spoke Russian (24%), mixed language and other language Ukrainians (13%), and ethnic Russians (regardless of language) (21%).

One-half of interviewed women (50%) said they were currently employed outside the home and another 13% were on maternity leave from their jobs at the time they were interviewed. The proportion describing themselves as unemployed (i.e., not working, but wishing to do so) was 18%. Eighty-three percent of respondents' homes had a color television, 43% had a telephone, 32% had a VCR, 25% had an automatic washing machine, and 33% of respondents lived in homes with automobiles.

Pregnancy and fertility

Pregnancy and childbearing represent topics of great interest in Ukraine and neighboring countries. The fact that the fertility rate is far below the replacement level and that the population of Ukraine began decreasing in recent years, due, in large part to extremely low fertility levels, has become a major concern. The 1999 URHS questionnaire collected an extensive amount of information on pregnancy and childbearing.

Childbearing in Ukraine tends to start at a much earlier age than in other developed countries: about one-half of interviewed 20-24 year-olds had already had a child. Among respondents ages 25-29, 84% of women had already borne at least one child. Ukraine's total fertility rate (TFR) according to survey data was about 1.4 births per woman for the two years preceding the survey, similar to the

official figure for Ukraine of 1.3 births. Age-specific fertility rates conform to the typical eastern European pattern, whereby childbearing tends to start and stop at a much earlier age than in other parts of Europe. Childbearing is heavily concentrated in the early years of marriage, with 81% of the TFR accounted for by births occurring before age 30.

Forty-five percent of recent pregnancies to respondents reportedly resulted in a live birth and 7% reportedly ended in miscarriage or stillbirth. The remainder, just under half, were terminated by induced abortion. The proportion of pregnancies resulting in a live birth declined steadily with increasing age, from 59% for 15-24 year-olds to 21% among 35-44 year-olds. Only 15% of pregnancies to women with two previously born children resulted in a live birth.

Fewer than half of recent pregnancies were said by respondents to have been planned at the time they occurred (i.e., the woman wanted to become pregnant at that time). Fifty-four percent were said to unplanned: 17% were mistimed and 38% were unwanted (the woman wanted no more children at the time she became pregnant). The likelihood of a pregnancy being planned fell sharply with increasing age, from 55% for 15-24 year-olds to only 20% for 35-44 year-olds. Pregnancies classified as unwanted resulted in a live birth only 5% of the time.

Almost two-thirds of survey respondents said they wanted no more children. The proportions who wanted no more children increased sharply with the number of living children, from 11% of women with no children to 91% of women with two children and 87% of those with three or more children. Even among women with only one child, about one-half said they wanted no more.

Induced abortion

The incidence of induced abortion in Ukraine, as in most of the former Soviet republics has been very high in recent decades. However, official statistics have revealed that rates have been declining in recent years. Between 1990 and 1998, the official annual rate of abortion incidence fell by over 50%, from 77 abortions to about 36 abortions per 1,000 women of childbearing age.

Overall, 43% of respondents had ever had an induced abortion and 20% had had more than one abortion. Among women in the oldest cohorts (ages 35-39 and 40-44), about two-thirds had had at least one abortion and about one-third had had at least two abortions during their lifetime. Three percent of all women and 8% of 40-44 year-olds listed five or more abortions.

The total induced abortion rate (i.e., the mean number of lifetime abortions per woman based on current age-specific abortion rates) for Ukraine was 1.6 abortions per woman for the two years before interview. The abortion rate was .054, meaning that just over one of every 20 women reported having an abortion during a one-year period. The abortion ratio (i.e., the ratio of induced abortions to live births) was 1.10, indicating slightly more abortions than live births. All of these figures were substantially higher in urban areas than in rural areas. As with fertility, age-specific abortion rates were highest among women in their twenties (.091), followed by women 30-34 years of age (.069).

Fourteen percent of recent abortions resulted in what women described as “complications requiring

medical treatment” immediately or soon after the procedure. Conventional abortions were about 50% more likely than miniabortions (i.e., early vacuum aspiration) to result in short-term complications. Among those women experiencing complications, 37% reported rehospitalization or extended hospitalization as a result. Thirteen percent of women with recent abortions reported that they received no anesthesia in association with the procedure. The likelihood of receiving anesthesia appears to have remained unchanged in recent years.

Infertility

There has been some concern expressed in recent years that infertility may be playing an increasing role in the unprecedentedly low levels of fertility now in effect in Ukraine. Fifteen percent of respondents reported having a problem becoming pregnant in the previous 10 years, with 10% having a problem lasting at least one year and 5% at least five years. Just over half of women who reported having difficulty becoming pregnant in the previous 10 years sought treatment of some kind for this problem. The most common treatment was the prescription of anti-inflammatory drugs, followed by: hormones, physiotherapy, treatments for blockage of the Fallopian tubes, and relaxation/spa therapy. Less commonly used were more costly and invasive treatments, such as laser therapy, laparoscopy, and in vitro fertilization. About half of the women receiving treatment eventually became pregnant. Forty percent had a live birth.

Contraception

Prior to this survey relatively, little national, systematic information existed on most aspects of contraceptive use in Ukraine. The URHS collected information on a broad array of topics related to contraception, including knowledge and use of contraceptive methods, source of methods, contraceptive failure and discontinuation, side effects, and reasons for nonuse of contraception.

Virtually all women knew at least one modern contraceptive method, led by condoms (99% knowledge), the IUD (96%), and oral contraceptives (90%). Knowledge of where to obtain supplied contraceptive methods tended to be almost as high as knowledge of methods, indicating that most women who had heard of a method also knew where they could obtain that method. Overall, 74% of respondents and 87% of those currently in union reported ever using any contraceptive method. The figures for modern methods were 60% and 69%, respectively.

Sixty-eight percent of women in registered or unregistered marriages were currently using contraceptive methods at the time of interview. More women/couples were employing modern methods of contraception than traditional methods (periodic abstinence and withdrawal), though the difference was not great (38% and 30%, respectively). IUDs (used by 19% of women in union) and condoms (14%) accounted for the vast majority of modern method use in Ukraine, with oral contraceptives (OCs) (3%) a distant third. Withdrawal, the most widely practiced method (20%) and periodic abstinence (10%), were also commonly employed.

Contraceptive prevalence looks to have risen steadily in recent years, rising by about 7 percentage points in five years.

Women's consultations and pharmacies were overwhelmingly the leading sources of oral contraceptives, supplying about three of every four users. Women's consultations were the predominant source of IUDs, accounting for two-thirds of those currently used. Pharmacies supplied slightly over half of condom users.

Among survey respondents who were not currently using contraception, three-fourths cited little or no sexual activity, an inability or difficulty in becoming pregnant (subfecundity), current pregnancy, or a desire to become pregnant, as their primary reason for non-use. There was a broad assortment of reasons for non-use given by the remaining 25% of non-users, many of which could be addressed by reproductive health interventions. By the conventional definition, unmet need for contraception was 15%, very high compared with the levels in most other developed countries. Using a definition that includes users of periodic abstinence and withdrawal (methods with typically low use-effectiveness) as having unmet need more than doubles the proportion in need, to 37%.

Overall about 9% of contraceptive users became pregnant while using a method within one year of beginning use. After three years this rose to 19%. The failure rate for the IUD was 1.4% for the first year, which is in line with typical rates. The failure rate for oral contraceptives was 5.9% for the first year and 13.2% for three years. Condom failure was 7.1% for one year and 18.7% for three years. The highest failure rates for widely used methods were found for periodic abstinence and withdrawal. For periodic abstinence 15.6% of users became pregnant in the first year; for withdrawal the rate was 11.7%. Three-year rates for these methods was almost 30%. In general, discontinuation rates were very high across methods. For all methods combined, 29% of contraceptive use segments continued for no more than one year.

Respondents were asked to rate a number of birth prevention methods with regard to safety and health effects, effectiveness, and cost, as well as to give each method an overall rating. The most noteworthy result is the nearly universally highly negative overall opinions held by respondents about both conventional induced abortion (96%) and miniabortion, i.e., vacuum aspiration in the early weeks of pregnancy (95%). Opinions about abortion were equally negative regardless of whether women had ever had an abortion. Abortion was by far the worst regarded method of birth prevention, but every method asked about was rated negatively overall by at least 40% of respondents. With regard to safety and health consequences, induced abortion was again viewed the most negatively of all methods (91% for conventional abortion and 88% for miniabortion).

Despite the fact that most women want to have no more children and that most couples desire no more children long before reaching the end long before their potentially fertile years end, contraceptive sterilization is relatively rare in Ukraine; only 2% of married women of reproductive age with two or more children had been sterilized and virtually no men had undergone a vasectomy. Only 7% of fecund respondents who wanted no more children expressed an interest in sterilization.

Contraceptive counseling

An important component of the Ukraine Women's Reproductive Health Initiative has been the development and implementation of interventions intended to improve health care workers' counseling regarding contraceptive services. Only thirty-nine percent of women with a recent

induced abortion said that a health professional had spoken with them about ways of preventing an unintended pregnancy following their most recent abortion. Only 7% were referred for contraceptive services or counseling. About one out of every seven women undergoing an induced abortion left the facility where it was performed with a contraceptive method or a prescription for one. One-fourth of respondents with recent live births received counseling about contraception subsequent to their delivery.

Just over half of women who recently started using a provider-supplied method said the provider had discussed family planning options with her. About two-thirds of recent IUD, pill, and injection users recalled their provider giving information on potential side effects and what to do about them.

Information, education, and communication (IEC)

Thirty-six percent of women reported seeing family planning information on television and 39% recalled seeing such information in newspapers or magazines in the previous six months.

Despite a minority of women recalling seeing information on family planning in the mass media, 85% of women said that information on family planning should be broadcast.

Sexual experience

The URHS included a series of questions designed to study certain aspects of sexual behavior among respondents. The proportions of women who have had sex by the time they turn 16, 18 and 20 years old has increased in recent years, indicating that the age at first sex has been declining somewhat. Of respondents in the 20-24 year-old cohort, 11% first had sexual intercourse before age 16, 42% before age 18, and 74% before age 20.

It has been hypothesized by some in the region that the dramatic declines in fertility and pregnancy rates in recent years in Newly Independent States (NIS) is attributable in large part to a decline in sexual activity rate, rather than other factors. About three-fourths of all sexually experienced respondents reported having sexual intercourse in the previous thirty days, 63% in the previous week, and 16% on the previous day. Among women currently in union, the proportion sexually active was 83% for the past month and 72% for the past week. Four of every ten women in union reported having intercourse at least 10 times in the previous 30 days. The overall median coital frequency was slightly over five times per month. It is difficult to detect in these data any indication that sexual activity rates have declined to a level that would contribute significantly to declines in fertility or pregnancy.

The median age at first sexual intercourse, based on reports of whether young survey respondents had ever had sexual intercourse, was about 18.4 years. The proportion of 15 and 16 year-olds reporting that they had ever had intercourse was 6% and 11%, respectively. However, many girls became sexually active at ages 17 or 18. Just 14% of sexually experienced young women reported that their first sexual intercourse took place following marriage. About equal numbers of women said that their first sexual partner was a “fiancé” or boyfriend (each 36%). Just under half of young women who first had intercourse before marriage (47%) reported that they or their partner used

contraception during her first premarital experience. Condoms accounted for over half of this contraception (28%), with withdrawal the only other commonly used method (13%).

Maternal and child health / Women's health

Ten percent of women received no prenatal care during their last pregnancy leading to a live birth. Another 3% waited until the third trimester of pregnancy to begin prenatal care. In only 65% of recent pregnancies did prenatal care begin during the first trimester, as recommended. Among women who received prenatal care, 81% made at least ten prenatal care visits. Seventy-eight percent of women with recent live births had a diagnostic ultrasound during pregnancy. Eighty-seven percent of women said that they had their blood pressure measured during pregnancy.

Thirty-two percent of women with recent deliveries leading to a live birth reported being hospitalized for prenatal problems. In spite of economic changes and on-going health sector reform, the proportion of women hospitalized was virtually unchanged between 1994-96 and 1997-99. The median stay was between two and three weeks, with two-thirds lasting for two weeks or more. Fifty-five percent of women received a postpartum check-up within six weeks of delivery of their last baby. Overall, 9% of women were cigarette smokers at the time they became pregnant. About half of those women reported that they stopped smoking during their pregnancy.

Ninety-two percent of recently born babies were breastfed. The percentage breastfed has not changed significantly in recent years; the proportions were very similar for babies born in 1994-1995, 1996-1997, and 1998-1999. The mean duration of breastfeeding for those babies who were breastfed was 7.5 months.

Nineteen percent of respondents said that they currently smoked cigarettes. Few of the current smokers could be considered heavy smokers, though, with only 3% of women reporting that they typically smoked more than ten cigarettes per day.

Sexually transmitted infections

Knowledge of certain sexually transmitted infections and associated conditions was very widespread: syphilis (2% unaware of it), pelvic inflammatory disease (5%), gonorrhea (9%), and genital ulcers (9%). The conditions about which the most respondents were unaware were human papilloma virus (HPV) (74%), genital herpes (66%), and chlamydia (54%). A very high proportion of women reported having had pelvic inflammatory disease (PID) (38%) or genital ulcers (38%) at some time during their life. Other diseases that an appreciable number of women reported ever having been diagnosed with were: trichomoniasis (4%), syphilis (2%), gonorrhea (2%), and chlamydia (2%). Almost one-third of respondents who experienced symptoms consistent with STIs in the previous 12 months did not consult a health care provider for diagnosis or treatment.

About one-third of respondents were not aware that someone could be infected with HIV and exhibit no symptoms. About one-half of women lacked awareness that people with STIs could have no symptoms. Only 6% thought that condoms provided excellent protection against STIs. Five percent of Ukrainian women perceived themselves at high risk for acquiring an STI, with another 6% saying

they were at medium risk, 29% at low risk, and 60% at no risk.

Domestic violence

Almost one of every five respondents recalled that their parents or stepparents abused each other while the respondent was growing up and almost 30% reported that they were physically abused as a child by someone in their household. Nineteen percent of ever-married women had ever had a partner threaten to hit her, 18% had ever been pushed or slapped, 13% had been punched, kicked, or hit with an object, and 4% had been threatened with a weapon. Twenty-one percent had had any of these acts committed against her by her partner. Eight percent had experienced these types of violence within the previous 12 months. Fifty-four percent of women who reported abuse by their partner in the previous 12 months reported sustaining injuries from these incidents.

CHAPTER I

INTRODUCTION

Background

Ukraine is an eastern European country bordered by the Russian Federation to the north and east, six smaller European countries to the west and northwest, and the Black Sea to the south. Its land area is about 604,000 square kilometers (233,000 square miles), making it the largest country entirely within Europe. The population in 1999 was estimated to be about 50 million people, the second most populous of the former Soviet republics. Owing primarily to a major decline in the rate of childbearing in Ukraine, the population of Ukraine has been declining in recent years. It is now estimated to be decreasing at an annual rate of approximately 0.6 percent. The country is divided into 26 oblasts and one autonomous region, the largest administrative divisions within the republics of the former Soviet Union.

Since the break-up of the Soviet Union in 1991, after which it became an independent state, Ukraine, like the other countries in the region, experienced dramatic and far-reaching political, social, and economic changes. These changes have brought about major transitions in many aspects of life, some positive and some negative. One such area in which change has been taking place is that of the financing and provision of health care and the structure of the Ukrainian health care system. The on-going reform in the health care system has significantly affected such factors as the availability, cost, and quality of health care services of all kinds throughout the country (Barr and Field, 1996; Grischenko, 1997).

In addition, the transitions that Ukraine has undergone, including those in the area of health care, have no doubt contributed to substantial changes that have occurred in reproductive health and demographic indicators in recent years (Steshenko and Irkina, 1999; Ukraine Cabinet of Ministers, 1997). There have been major changes noted in rates of childbearing, abortion, and contraceptive use. There are also unanswered questions about the degree to which there have been changes in such areas as the utilization of maternal-child health services, the prevalence of infertility problems, and maternal behaviors that affect maternal and infant health (such as breastfeeding, prenatal care practices, and immunization coverage). However, relatively little detailed and conclusive information was available about the situation in Ukraine with regard to a number of important reproductive health topics or the degree to which the reproductive health situation has changed over the past decade.

This volume is the final report of the 1999 Ukraine Reproductive Health Survey (URHS). This nationwide survey of women of reproductive age was carried out from June to October of that year. It was a population-based survey, intended to generate results on a broad variety of reproductive health topics and to be representative of all women 15 to 44 years of age living in households in Ukraine. It constitutes the largest and most in-depth data collection effort in the area of reproductive health and related topics since Ukrainian independence. Another important national survey conducted in 1996, known as the "Health-1996 Sociological Survey", was somewhat smaller and provided a considerable amount of valuable information on reproductive health, but, since it focused

on other topics as well, did not yield as much detail as the URHS with regard to certain topics, such as family planning and pregnancy outcomes (Cabinet of Ministers of Ukraine et al, 1997) .

The Ukraine Women's Reproductive Health Initiative

The 1999 survey was performed in conjunction with the Ukraine Women's Reproductive Health Initiative (UWRHI), a project sponsored by the United States Agency for International Development (USAID) (Bergthold et al., 1998). This project consisted of a variety of components, all designed to help to reduce maternal morbidity and mortality and improve reproductive health generally in Ukraine. The stated overall goals of this initiative were to improve the quality of reproductive health services and to improve access to those services. The interim goals of the initiative were to increase the rate of modern contraceptive use and to reduce abortion rates in service sites. Those developing the initiative anticipated that improved access to and quality of reproductive health services for women, expanded and improved use of effective contraception, and reduced reliance on abortion as a means of birth prevention would result in reductions in maternal morbidity and mortality.

At the outset of the project, seven collaborating agencies were assigned specific roles in this initiative. The specific strategic objectives of the UWRHI, designed to help Ukraine meet the abovementioned goals were:

- the establishment of demonstration sites for training and delivery of family planning services: each of the sites was to provide family planning services and training
- the institutionalization of reproductive health training: at each demonstration site reproductive health was incorporated into the pre-existing program of refresher training
- increased public information, education, and communication about family planning: the project produced videos, pamphlets, and other products to public awareness and knowledge about family planning
- improved family planning policy environment: carrying out family planning/reproductive health advocacy work targeted at government agencies, legislators, and policy makers
- increased supply and distribution of family planning methods: provision of oral contraceptives, IUDs, and injectable contraceptives to the demonstration sites
- improved methods of child delivery and maternal care: this includes activities to promote breastfeeding, rooming in and other "family centered maternity care" nationwide.

The UWRHI was to a great extent a response of the donor community to a perceived need to improve reproductive health and family planning services in Ukraine. In 1995, the Cabinet of Ministers adopted a National Family Planning program for 1995-2000 that called for widespread provision and promotion of modern contraception and a reduction in the number of abortions. The four primary components of the program were: 1) preventing unwanted pregnancies, 2) providing medical-genetic counseling, 3) treating infertility, and 4) educating the public. However, funds were never made available to implement the program due to an extreme shortfall in the Ministry of Health

(MOH) budget.

The 1999 Ukraine Reproductive Health Survey

As the largest and one of the first nationwide, population-based surveys of reproductive health and related issues to take place in post-independence Ukraine, the 1999 URHS provides a substantial amount of new information on a broad assortment of reproductive health topics. There were two major reasons for carrying out this survey. First, it was intended to fill a substantial need for data regarding the current status of reproductive health of Ukrainian women. Besides providing data on the current situation overall, it also adds to what is known about reproductive health trends and differentials within the population and allows more accurate determinations to be made about the needs of the population in this area. The survey allows tabulation of many basic indicators of, for example, contraceptive use, unintended pregnancy, unmet need for family planning services, use of reproductive health services, contraceptive failure, and abortion and related factors. The data help to determine where in the country particular reproductive health needs and problems are most prevalent or severe and in which segments of the population such problems are the most or least likely to exist.

Secondly, the URHS was designed to provide programmatically useful results. The data collected on reproductive health can be used to help direct, modify, or develop interventions, as well as to provide information to policy makers and health care program officials and providers.

The information coming from the survey should prove helpful for policy makers, health care providers, program officials, international organizations, NGOs, and others working in reproductive health and related fields. Since there are great similarities between Ukraine and other formerly communist republics, this information should also prove valuable for those working in other countries within the region.

The 1999 URHS was also designed to provide important information for a broad cross-section of reproductive health and women's health topics. One of the principle issues that the survey was designed to address was the use of abortion among Ukrainian women. Like most other former Soviet republics and countries under Soviet domination, abortion has been a dominant means of birth prevention for many years. The Ukraine Women's Reproductive Health Initiative was designed to help determine effective means (and implement those means) to reduce reliance on induced abortion as a means of family planning and thereby improve reproductive health. The URHS examined the use of abortion in depth, including such factors as incidence, attitudes, cost, and complications.

A second important issue examined was the use of contraception, in order to look at levels and trends in contraceptive prevalence and method selection and at the extent to which family planning methods are being practiced effectively. The questionnaire included detailed information on many aspects of contraception. Expanded and improved contraceptive use should improve maternal health by reducing the numbers of unintended pregnancies and induced abortions. It is also important to learn more about women's opinions and attitudes regarding specific contraceptive methods and abortion, and about women's knowledge of reproductive health issues, to determine how well informed the population is and to assist in the development of information, education, and communication (IEC) messages.

A broad assortment of other issues were addressed in the survey as well. The URHS was used to learn about such things as the reproductive health services women are using and their opinions about those services, women's health behaviors, sexual activity, sexually transmitted infections, and domestic violence, among others.

The 1999 URHS was also designed to help make determinations about possible impacts of the Women's Reproductive Health Initiative. Examinations of those areas in which it was anticipated that the initiative may have an impact were performed. The survey questionnaire permits analysis of trends in some key topics, such as contraceptive use and abortion, in order to determine whether they have been changing in recent years and in particular areas of the country or segments of the population.

This Final Report describes the key initial findings from the 1999 Ukraine Reproductive Health Survey in all of the areas examined. However, it was not possible to do an exhaustive analysis of data on any single topic, because of the scope and depth of the information collected in the URHS. Data analysis will continue after the publication of this report to utilize the data as fully as possible.

CHAPTER II

SURVEY METHODOLOGY

The 1999 Ukraine Reproductive Health Survey was designed to collect information from a representative household sample of all women between the ages of 15 and 44 living throughout Ukraine, excluding those living in institutional settings. Although some pregnancy, childbearing, and abortion occurs outside the ages 15 to 44, the relative rarity of these events at those ages in Ukraine suggested that it would be most efficient to limit the sample to women in this age range.

Organizational Structure

The 1999 URHS was implemented by the Kiev International Institute of Sociology (KIIS), a professional survey organization with extensive experience in all aspects of social science survey research. KIIS was responsible for selecting the sample of households and individuals, recruiting and training interviewers, conducting field work, processing the data, and performing part of the data analysis. The United States Agency for International Development (USAID) mission in Kiev was the source of all funding for the URHS. USAID also was consulted regarding the content of the survey, to ensure that the information collected would be of maximum value for its projects. Technical assistance for all phases of the survey was provided by the Division of Reproductive Health of the United States Centers for Disease Control and Prevention (CDC/DRH). CDC/DRH was the lead agency in development of the overall survey design, questionnaire construction, coordination of survey activities, and much of the data analysis. The participation of CDC/DRH was funded through a Participating Agency Service Agreement between the USAID Office of Population and CDC/DRH. Other cooperating agencies involved in the Ukraine Women's Reproductive Health Initiative, particularly Johns Hopkins University/Population Communications Services (JHU/PCS) and The Futures Group International, contributed significantly to questionnaire development, as well as other aspects of the survey.

Questionnaire Content

The 1999 URHS consisted of two questionnaires: a short household instrument and a much longer individual questionnaire. Both questionnaires are included in Appendix A of this report. The household questionnaire was two pages long and was administered to any adult living in visited households. It consisted primarily of information regarding the individuals who lived in the household and the location of the residence.

The individual form was completed by selected women 15 to 44 years of age who agreed to be interviewed. This questionnaire covered a wide range of topics related to reproductive health status and needs in Ukraine. The sections of the questionnaire were:

- I. Social and demographic characteristics of respondents*
- II. Fertility, pregnancy, and abortion*— includes a complete pregnancy history, detailed information on abortions and live births in the preceding five years, use of maternal-

child health services, and infertility problems and treatment.

- III. *Contraception***– includes knowledge and use of specific methods, a month-by-month calendar of contraceptive use in the preceding five years, contraceptive counseling, and detailed information on many aspects of family planning.
- IV. *Information, education, and communication (IEC) and attitudes and beliefs concerning family planning***
- V. *Women’s health***– includes information on sexual behavior and sexually transmitted infection knowledge and history
- VI. *Socioeconomic characteristics of respondents***
- VII. *Intimate partner violence.***

Survey design

The survey was designed to obtain interviews with a nationally representative sample of about 7,000 women between the ages of 15 and 44 years of age living throughout Ukraine. The survey was intended to be large enough to provide stable estimates regarding the main topics of interest at the regional level (there are five regions: the North, East, Central, South, and West) and for the urban and rural sectors of Ukraine. The sample was geographically self-weighting, with the exception of substantial over-sampling of two sites--the oblasts of Donetsk and Odessa.

The two oversampled oblasts were the sites for USAID-funded reproductive health activities. Donetsk was a site of activities sponsored by the Women’s Reproductive Health Initiative that started in 1996. Odessa was the focus of USAID-sponsored activities implemented by The Policy Project (coordinated by The Futures Group International) that are designed to improve support for family planning activities. Oversampling of these areas was designed to allow local estimates to be made in regard to many of the topics addressed in the URHS.

Three-stage cluster sampling was used to select survey respondents. Potential respondents consisted of all women between the ages of 15 and 44 years who lived in households anywhere in Ukraine. The first stage of sampling consisted of the selection of primary sampling units (PSU). Approximately 550 primary sampling units were selected across Ukraine. The sample was selected proportional to population size (PPS) of each of the country’s 26 oblasts and autonomous regions (with the exception of Odessa and Donetsk). Within each oblast the sample was split proportionally into five size-of-place categories, ranging from large cities to rural areas, using software that listed the estimated population of all locations. Population estimates were based on yearly updates made to census counts. Unfortunately, the most recent census in Ukraine took place when it was still part of the Soviet Union, in 1989. Within each size of place/oblast category, PSU (census enumeration districts) were selected with probability proportionate to size. Within rural areas, post offices were selected instead of places, under the assumption that all post offices cover roughly the same population. This process guarantees a selection of households approximately proportional to the

entire population according to oblast and size of place.

The second stage of sampling consisted of the selection of dwelling units and respondents from the selected PSU. Within each selected PSU a random starting point was chosen, followed by selection of contiguous dwelling units, selected in a predetermined order. The number of dwellings visited per PSU varied from 23 to 30, depending on the size of place, since the average number of women of childbearing age per household varies according to size of place. Selection of women for interview was accomplished in the third stage by listing women in each visited household by descending age and selecting every second woman listed regardless of the household in which she was found. However, this procedure was slightly modified to ensure that no more than one woman per household was interviewed

Data Collection

The questionnaires were pretested in November 1998. Eight interviewers conducted pretest interviews with a total of about 50 women in Kiev and a nearby rural area. Based on these interviews, the survey questionnaires were modified. Data collection for the 1999 URHS was carried out by about 150 female interviewers living throughout Ukraine, most of whom were highly experienced in conducting interviews. Staff from KIIS, assisted by Dr. Petr Velebil, a Russian-speaking obstetrician-gynecologist and epidemiologist from the Czech Republic, conducted interviewer training sessions. Five such sessions were held, two in Kiev, one in Donetsk, one in Odessa, and one in L'viv. Each session had about 25-30 interviewers attending. Interviewer training sessions consisted of intensive training in field procedures and administration of the questionnaire.

Field work lasted from June through October of 1999. Each interviewer was assigned to visit a small number of PSU in the part of the country in which she lived. Interviews took place at respondents' homes and typically lasted from 60 to 75 minutes. Each interviewer forwarded her completed questionnaires to her regional supervisor, who reviewed each questionnaire and, if satisfactorily completed, sent it to the KIIS office in Kiev for data entry and final editing.

Response rates

At least one 15-44 year-old woman was identified as living in 55% of sampled households (Table 2.1, upper panel). In most of the remaining households, there were no resident females eligible for interview. Residents refused to provide interviewers with information concerning the household or its residents in fewer than 1% of the households visited. The difference between urban and rural areas was small, with urban households slightly less likely than rural ones to contain women eligible for interview.

The lower panel of Table 2.1 presents information on response rates for women who were selected for interview. Of those 15-44 year-old women who were identified as living in visited households and selected as potential respondents, 85% were interviewed. The figure was somewhat higher in rural areas (89%) than in urban areas (84%). Eight percent of women selected for interview were never found at home and 6% refused to be interviewed. Individual refusal rates were about twice as high in urban areas as in rural areas.

Table 2.1
Percentage distribution of household and individual final interview status, by residence
1999 Ukraine Reproductive Health Survey

	Total	Urban	Rural
<u>Selected households</u>			
Eligible woman identified	54.9	54.0	58.3
No eligible woman in household	43.5	44.3	40.2
Household refusal	0.6	0.7	0.2
Residents not at home	0.7	0.6	0.9
Unoccupied household	0.2	0.3	0.1
Other	0.3	0.3	0.3
Total	100.0	100.0	100.0
<i>Number of households visited</i>	<i>16,886</i>	<i>13,383</i>	<i>3,503</i>
<u>Selected eligible women</u>			
Completed interviews	85.2	84.1	89.0
Selected women refused	5.6	6.3	3.0
Selected woman absent	7.7	7.9	6.9
Woman not competent	0.4	0.5	0.2
Other	1.2	1.2	0.9
Total	100.0	100.0	100.0
<i>Number of selected women</i>	<i>8,367</i>	<i>6,587</i>	<i>1,780</i>
<i>Number of women interviewed</i>	<i>7,128</i>	<i>5,544</i>	<i>1,584</i>

CHAPTER III

CHARACTERISTICS OF RESPONDENTS

This chapter presents selected social, economic, and demographic characteristics of the women who were interviewed in the 1999 URHS. The URHS included modules that covered such topics as demographic characteristics, marriage, education, language, ethnicity, religion, employment, and other economic topics. These data provide important background information about the population of women of childbearing age in Ukraine, giving some important insights into their social and economic conditions, as well as such things as the degree of heterogeneity and diversity in the population. It also lets us know the extent to which the survey sample is representative of Ukrainian population of women of childbearing age as a whole.

Demographic and social characteristics

Table 3.1 displays percentage distributions of URHS respondent characteristics according to selected demographic and social characteristics. Almost three-fourths of respondents (73%) were living in areas defined as urban. The remaining 27% lived in rural areas, i.e., villages or farms. This distribution has less to do with Ukraine being a heavily urbanized country than it does with the fact that the definition of urban includes settlements down to quite a small size. About half the urban population of women of childbearing age was living in cities of more than half a million people.

With the exception of the Oblasts of Donetsk and Odessa, which were intentionally oversampled, the percentage distribution of the population by oblast was very similar in the sample and Ukraine's official population statistics (data not shown). Donetsk constituted 21.7% of the sample, but only 10.1% of the total population, while the figures for Odessa were 17.9% and 5.2%, respectively. During analysis, sample weights have been applied to the results to compensate for this oversampling. With the appropriate sample weights the percentage distribution by oblast is similar to the official population estimates. All results presented in this report are based on weighted data, adjusting for oversampling in Donetsk and Odessa, but unweighted totals are shown in the report's tables.

The age distribution closely matched the distribution according to official national estimates. The distribution was very flat, with very similar proportions of respondents in each five-year age group, ranging from 16.4% for ages 20-24 to 17.3% for ages 35-39. The urban and rural age distributions were also very similar to each other.

Just over six of every ten respondents were currently living in either a registered or unregistered marriage, with unregistered marriages being relatively rare (4% of women). Another 9% of respondents were either divorced or separated and 2% were widows. Twenty-three percent of women had never been in union (i.e., married or lived with a man). The percentage of women in registered marriages was considerably higher in rural areas than in urban areas (70% vs. 60%, respectively), principally a function of earlier age at marriage in rural Ukraine. The percentage divorced or separated was almost twice as high in urban areas as in rural areas.

Ukrainian women tend to be well educated, as evidenced by the fact that only 11% of respondents had not completed secondary schooling. About half of that 11% were 15-19 years old, so many of them, no doubt, were still in secondary school and will eventually complete it. Just over two-thirds of women had finished secondary school, but had not received any post-secondary schooling. Additionally one of every five respondents had received some post-secondary education. Within the “completed secondary” category there is considerable variation among women in educational attainment. A majority of women in this category obtained some additional professional technical training in addition to secondary school. The greatest difference between urban and rural women in educational attainment was that the former were about twice as likely as the latter to have received any post-secondary education (23% vs. 13%, respectively). Rural women were slightly more likely not to have completed secondary school.

Women’s stated nationality was cross-classified with the language they most often spoke. This was done because neither nationality nor language alone completely describes the extent to which people may be considered to be ethnically Russian, Ukrainian, mixed, or something else, in a society with a long history of large populations of Ukrainians and Russians and not always a clear distinction between the two. The largest ethnic-linguistic group consisted of Ukrainians who most often spoke Ukrainian (38%), but there were also large numbers of Ukrainians who usually spoke Russian (24%), mixed language and other language Ukrainians (13%), and ethnic Russians (regardless of language) (21%). In rural areas, over two-thirds of women were Ukrainian-speaking Ukrainians and only 10% classified themselves as ethnic Russians. A clear majority of Russian-speakers are urban residents.

With regard to religion, about three of every four respondents said they were Orthodox, while 11% described themselves as having no religion. Urban respondents were more likely than rural women to say they had no religion (13% vs. 7%, respectively). Although a high proportion of women reported belonging to a religion, relatively few attended religious services on a regular basis. Only 12% said they attended services at least monthly, with about half of them attending weekly. Even though the overall level of attendance was relatively low, the percent who never attended was lower in rural areas (26%) than in urban areas (35%).

Table 3.2 displays most of the same characteristics as Table 3.1 according to respondents’ region of residence. It is clear that there are major social and demographic differences between different parts of Ukraine, with the most notable differences between the Western region and the rest of the country. In The West there were roughly equal percentages of respondents in urban and rural areas, making it much more rural than the other parts of Ukraine. The East, Ukraine’s most industrialized region, on the other hand is the most urban region (86%). Age distributions were very similar across regions. Western women were the most likely to be in a registered marriage and the least likely to be in an informal or unregistered union. There were major regional differences in language and ethnicity. The proportion of women reporting themselves to be predominantly Ukrainian-speaking and ethnically Ukrainian varied tremendously, ranging from 89% in the West and 72% in the Central region to only 5% in the South. On the other hand the percentage who were ethnically Russian ranged between 35% in the East to less than 3% in the West. In all regions except the West, an overwhelming majority of respondents said they were Orthodox (78%-85%). In the West, Orthodox was also in the majority, but much less so than elsewhere (58%). Attendance at religious services

was considerably more frequent in the East than elsewhere (18% at least weekly and 17% at least monthly), but even there most women attended services infrequently. Attendance was least common in the Central and Eastern regions.

Economic characteristics

Since the break-up of the Soviet Union and the downfall of communism in this part of the world, Ukraine has been undergoing dramatic economic changes. The dissolution of the former economic system and the concurrent growth of capitalism and privatization has led to economic freedoms that did not previously exist. The transition to a market economy, however, has also had some serious negative consequences for many individuals, at least in the short run. Certain economic protections, such as employment security and controlled prices, no longer exist. The URHS included several questions related to women's employment and other economic issues.

One-half of interviewed women (50%) said they were currently employed outside the home and another 13% were on maternity leave from their jobs at the time they were interviewed (Table 3.3). The proportion describing themselves as unemployed (i.e., not working, but wishing to do so) was quite high, at 18%. The employment situation was somewhat worse in rural areas than in urban areas. The percent unemployed was 22% in the former, compared with 17% in the latter. There were substantial differences in unemployment according to region, with the proportions ranging from 14% in the heavily industrialized East to 22% in the West, the most rural region of Ukraine (data not shown).

The bottom panel of Table 3.3 displays the percentages of women who reported that they lived in households containing various possessions or amenities. Eighty-three percent of respondents' homes had a color television, 43% had a telephone, 32% had a VCR, 25% had an automatic washing machine, and 33% of respondents lived in homes with automobiles. With the exception of automatic washing machines and automobiles, urban women were substantially more likely than rural women to have each of the possessions and amenities asked about.

Marriage

As is true throughout most of eastern Europe and the former Soviet Union, women tend to marry at a very young age compared to other developed countries. URHS data reveal that most first marriages take place when women are in their early twenties. Only about one-third of women between the ages of 20 and 24 years had never been in a registered or unregistered marriage (Table 3.4 and Figure 3.1). By ages 25-29 relatively few women (8%) had never been in union. Marriage has also been much more universal than in other developed countries. In the two oldest cohorts of women interviewed, the proportions of women who had never been in union were very low, below 3%. Divorce and separation, while not at the levels found in many western countries, is not a rare occurrence. Starting with the 25-29 year-old cohort, 10% or more of each cohort was currently divorced or separated.

It also is evident that rural women tend to marry somewhat earlier than urban women. Among 20-24

year-olds, only 22% of rural respondents had never been in either a registered or unregistered marriage, compared with 41% in urban areas. In addition, the proportion of older respondents never in union was even lower in rural areas (under 2%) than it was overall. Divorce and separation was also less common in rural areas than in urban areas.

Table 3.5, showing proportions of women ever in union and currently in union, confirms the tendency to marry young among Ukrainian women. Almost two-thirds of women between the ages of 20 and 24 years had already been in a registered or unregistered marriage (i.e., in union), and by ages 25-29, 93% had ever been in union (Table 3.5). In rural areas, age at marriage tends to be even younger than overall. 78% of 20-24 year-olds and 94% of 25-29 year-olds there had ever been in union.

Table 3.6 allows one to look at recent trends in age at marriage, by showing the percentage of respondents in 5-year age various cohorts who had ever been in union before various ages. The main finding from this table is that there is no sign that age at marriage has been rising in recent years. The proportions of women who had married by 20 was relatively constant in the 20-24, 25-29 and 30-34 year-old cohorts, with slightly over four of every ten women married before age 20. This indicates little change in the age at marriage. There are also no signs of increasing age at marriage among younger cohorts. About 14-15% of 20-24 and 25-29 year-olds were already in union by age 18. By age 25, almost nine out of every ten women had been in union. These figures are typical for most of eastern Europe but are extremely high compared to other parts of the continent, where marriage most often takes place in the late twenties and thirties.

Figure 3.1
Percentage Distributions of Current Marital Status,
by Age of Respondent
1999 Ukraine Reproductive Health Survey




Table 3.1
Percentage distributions of socio-demographic characteristics of respondents, by residence
1999 Ukraine Reproductive Health Survey

<u>Characteristic</u>	<u>Total</u>	<u>Urban</u>	<u>Rural</u>
<u>Residence</u>			
Urban, Total	72.7		
City >500,000	34.0		
City 200,000-500,000	12.5		
City <200,000	17.8		
Small town	8.4		
Rural, Total	27.3		
<u>Region</u>			
North	15.6	15.6	15.7
Central	11.1	9.0	16.7
East	35.0	41.6	17.4
South	17.6	19.3	13.0
West	20.8	14.6	37.2
<u>Age</u>			
15-19	16.6	16.9	15.8
20-24	16.4	16.6	16.0
25-29	16.6	16.2	17.7
30-34	16.5	15.9	18.2
35-39	17.3	17.8	16.0
40-44	16.5	16.6	16.3
<u>Marital status</u>			
Registered marriage	62.5	59.9	69.5
Unregistered marriage	3.9	4.3	2.6
Divorced/separated	9.0	10.1	6.3
Widowed	1.8	1.7	2.1
Never married	22.8	23.9	19.6
<u>Education</u>			
< Complete secondary	11.3	10.4	13.5
< Grade 10	8.2	7.6	9.6
Technical after grades 7-8	3.1	2.8	3.9
Complete secondary	68.4	66.2	74.0
Complete secondary	27.6	25.3	33.8
Prof. Tech after grade 10-11	11.3	10.9	12.3
Complete secondary + tech.	29.5	30.0	27.9
> Complete secondary	20.4	23.3	12.5
Incomplete postsecondary	3.9	4.6	2.2
Complete postsecondary	16.4	18.7	10.3

Characteristic	Total	Urban	Rural
<u>Language spoken at home/ Nationality</u>			
Ukrainian-speaking / Ukrainian	37.8	25.8	69.7
Russian-speaking / Ukrainian	24.2	31.1	5.9
Other or mixed language / Ukrainian	12.5	13.3	10.3
Russian nationality	20.9	25.5	8.5
Other nationality	4.7	4.3	5.6
<u>Religion *</u>			
Orthodox	76.2	77.0	73.9
Roman Catholic	4.8	3.6	7.8
Other	7.9	6.6	11.2
None	11.2	12.8	7.2
<u>Frequency of church attendance</u>			
At least once/week	6.2	4.7	9.9
At least once/month	6.2	5.0	9.2
Less than once/month	55.4	55.5	55.4
Never	21.1	22.1	18.4
No religion	11.2	12.8	7.2
Total	100.0	100.0	100.0
<i>Number of respondents</i>	<i>7128</i>	<i>5544</i>	<i>1584</i>

*228 women did not state their religion.

Table 3.2
 Percentage distributions of socio-demographic characteristics of respondents, by region of residence
 1999 Ukraine Reproductive Health Survey

Characteristic	Total	North	Cent.	East	South	West
<u>Residence</u>						
Urban, Total	72.7	72.5	58.9	86.4	79.8	51.1
City >500,000	34.0	84.7	0.0	52.6	48.2	8.3
City 200,000-500,000	12.5	13.6	24.0	4.6	16.6	15.5
City <200,000	17.8	15.9	20.8	20.8	8.9	20.0
Small town	8.4	8.3	8.4	8.4	6.1	7.3
Rural, Total	27.3	27.5	41.1	13.6	20.6	48.9
<u>Age</u>						
15-19	16.6	16.7	16.2	16.9	16.9	16.1
20-24	16.4	16.0	14.9	16.4	18.0	16.3
25-29	16.6	17.0	16.0	15.2	16.6	19.1
30-34	16.5	16.1	18.4	16.1	15.3	17.5
35-39	17.3	17.3	16.8	17.4	17.4	17.4
40-44	16.5	16.9	17.8	18.1	15.8	13.6
<u>Marital status</u>						
Registered marriage	62.5	61.8	64.3	59.9	58.4	70.1
Unregistered marriage	3.9	3.8	4.3	4.5	5.3	1.5
Divorced/separated	9.0	8.2	9.6	10.9	10.2	5.3
Widowed	1.8	2.0	2.7	1.8	1.8	1.2
Never married	22.8	24.3	19.0	23.0	24.2	21.9
<u>Education</u>						
< Complete secondary	11.3	8.7	12.6	12.3	11.4	10.6
Complete secondary	68.4	66.6	70.2	69.7	63.7	70.3
> Complete secondary	20.4	24.6	17.1	18.0	24.9	19.1
<u>Language spoken at home/</u>						
<u>Nationality</u>						
Ukrainian-speaking Ukrainian	37.8	42.5	71.8	11.4	4.7	88.7
Russian-speaking Ukrainian	24.2	26.1	5.4	36.6	36.2	1.8
Other/Mixed Ukrainian	12.5	19.0	12.8	12.7	17.6	2.9
Russian nationality	20.9	10.0	6.9	35.4	32.1	2.5
Other nationality	4.7	2.5	3.0	4.0	9.4	4.2
<u>Religion *</u>						
Orthodox	76.2	85.4	84.7	77.8	80.7	58.1
Roman Catholic	4.8	2.2	2.0	0.3	0.4	19.1
Other	7.9	1.9	1.5	5.2	6.4	21.4
None	11.2	10.5	11.8	16.6	12.5	1.3
<u>Frequency of church attendance</u>						
At least once/week	6.2	3.2	1.8	2.8	3.6	18.4
At least once/month	6.2	4.5	2.0	2.1	5.0	17.3
Less than once/month	55.4	63.3	67.3	48.9	51.5	57.5
Never	21.1	18.6	17.1	29.6	27.4	5.4
No religion	11.2	10.5	11.8	16.6	12.5	1.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>7128</i>	<i>858</i>	<i>617</i>	<i>2669</i>	<i>1824</i>	<i>1160</i>

*228 women did not state their religion.

Table 3.3
 Percentage distributions of current employment status
 and percent of women who live in homes with selected possessions, by residence
 1999 Ukraine Reproductive Health Survey

Characteristic	Total	Urban	Rural
<u>Current employment</u>			
Employed	50.0	51.4	46.3
On maternity leave	12.8	11.7	15.6
Not employed*	37.3	36.9	38.1
Unemployed**	19.2	16.6	22.0
Total	100.0	100.0	100.0
<u>Possessions in home</u>			
Color television	82.9	88.6	67.9
Bathroom/Shower	79.8	86.6	61.6
Telephone	43.1	48.9	27.3
Automobile	32.8	33.0	32.2
VCR	30.2	33.5	21.3
Automatic washing machine	25.1	25.1	25.1
Personal computer	3.2	4.0	1.2
<i>Number of respondents</i>	<i>7128</i>	<i>5544</i>	<i>1584</i>

*Does not currently have a job, regardless of reason. Includes "unemployed".

**Does not have a job due to inability to find a job.

Table 3.4
 Percentage distributions of current marital status by age of respondent, by residence
 1999 Ukraine Reproductive Health Survey

Marital status	Age of respondent						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
<u>All women</u>							
Registered marriage	8.6	55.7	75.5	80.9	77.5	76.6	62.5
Unreg. marriage	2.2	5.5	5.3	4.1	3.3	2.9	3.9
Divorced/Separated	0.5	5.2	10.5	10.0	13.7	14.2	9.0
Widowed	0.1	0.2	1.2	1.8	2.9	4.5	1.8
Never married	88.6	33.4	7.5	3.2	2.6	1.8	22.8
<i>Number of women</i>	<i>1,100</i>	<i>1,182</i>	<i>1,227</i>	<i>1,195</i>	<i>1,246</i>	<i>1,178</i>	<i>7,128</i>
<u>Urban</u>							
Registered marriage	7.6	50.3	72.4	81.1	75.5	73.8	59.9
Unreg. marriage	2.8	6.1	5.5	4.7	3.8	3.3	4.3
Divorced/Separated	0.6	6.0	12.7	9.8	15.3	15.9	10.1
Widowed	0.1	0.2	0.1	1.5	2.6	4.9	1.7
Never married	88.8	41.4	8.3	2.7	2.7	2.3	25.1
<i>Number of women</i>	<i>861</i>	<i>919</i>	<i>941</i>	<i>908</i>	<i>990</i>	<i>925</i>	<i>5,544</i>
<u>Rural</u>							
Registered marriage	11.3	70.6	82.9	60.6	83.6	84.1	69.4
Unreg. marriage	0.5	4.0	4.8	2.7	1.9	1.6	2.6
Divorced/Separated	0.0	2.8	5.2	10.5	8.9	9.6	6.3
Widowed	0.0	0.4	1.5	2.5	4.1	3.6	2.0
Never married	88.2	22.2	5.7	3.7	1.6	1.2	19.6
<i>Number of women</i>	<i>239</i>	<i>263</i>	<i>286</i>	<i>287</i>	<i>256</i>	<i>253</i>	<i>1,584</i>
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 3.5
 Percent of respondents who have ever been in union and who are currently in union, by current age
 1999 Ukraine Reproductive Health Survey

Age	% Ever in union	% Currently in union
<u>All women</u>		
15-19	11.4	10.8
20-24	66.6	61.2
25-29	92.5	80.7
30-34	96.8	85.0
35-39	97.4	80.8
40-44	98.2	79.4
Total	77.3	66.4
<i>Number of women</i>	7,128	7,128
<u>Urban women</u>		
15-19	11.2	10.4
20-24	62.5	56.4
25-29	91.7	77.9
30-34	97.0	85.7
35-39	97.1	79.3
40-44	97.9	77.1
Total	76.1	64.2
<i>Number of women</i>	5,544	5,544
<u>Rural women</u>		
15-19	11.9	11.9
20-24	77.8	74.6
25-29	94.3	87.7
30-34	96.3	83.3
35-39	98.4	85.5
40-44	98.8	85.7
Total	80.4	72.1
<i>Number of women</i>	1,584	1,584

Table 3.6
 Percent of women who were married or in a consensual union before selected ages
 by current age
 1999 Ukraine Reproductive Health Survey

Married before:	Current age of respondent						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
Age 16	-	0.9	1.0	0.5	0.4	0.8	0.8
Age 18	-	14.2	14.9	9.7	9.7	7.4	11.2
Age 20	-	42.8	44.8	41.1	37.4	36.3	40.0
Age 25	-	-	88.3	89.4	83.1	85.4	84.2
Age 30	-	-	-	95.7	94.0	94.1	93.5
<i>Number of women</i>	1,100	1,182	1,227	1,195	1,246	1,178	7,128

CHAPTER IV

PREGNANCY AND CHILDBEARING

The 1999 URHS questionnaire collected an extensive amount of information on pregnancy and childbearing. The survey instrument contained a complete pregnancy history, that included, for every pregnancy: its outcome, whether the pregnancy was intended, its duration (in months), and the gender of each live-born child. In addition, for pregnancies resulting in a live birth since the beginning of 1994, more detailed information was collected on prenatal care, hospitalization during pregnancy, and breastfeeding.

Pregnancy and childbearing represent topics of considerable interest in Ukraine and nearby countries. In most countries of the region fertility has fallen to unprecedentedly low levels in recent years, with levels well below the replacement level of 2.1 births per woman. Figure 4.1 displays fertility rates for the year 2000 for selected European countries, including most of the countries formerly part of the Soviet bloc. There is considerable concern about these topics for a number of reasons. Fertility rates that have fallen to such a low level in Ukraine have led to worries about declining population size as well as related concerns, such as a rapidly aging population and an eventual shortage of working age people in the population. In fact, the population of Ukraine began decreasing in recent years, due, in large part to extremely low fertility levels. The decline is now estimated to be about 0.6%, about 300,000 people, per year. Despite the low level of fertility, there remain concerns, however, about very high rate of unintended pregnancy, which fuels the high incidence of induced abortion.

The vital statistics system in Ukraine is considered to be quite complete in the area of fertility. Thus, official estimates of fertility rates are probably reliable. However, the URHS provides information on other aspects of pregnancy and childbearing that are not generally available from other sources. In this chapter we provide survey findings such topics as: pregnancy rates, pregnancy outcomes, fertility patterns, additional children desired, and the planning status of recent pregnancies.

Fertility and pregnancy levels and patterns

As in much of eastern Europe and the former Soviet Union, childbearing in Ukraine tends to start at a much earlier age than in other developed countries. About one-half of interviewed 20-24 year-olds had already had a child and the average number of live births for women in that cohort was about 0.6 (Table 4.1). Childbearing tends to begin earlier in rural areas, where 63% of 20-24 year-olds had had a live birth, than in urban areas, where 46% had begun childbearing. Among respondents ages 25-29, 84% of women had already borne at least one child. Mean family size for the oldest cohorts (which can reasonably be viewed as completed family size, since few women have children after about age 35) was 1.8 births per woman. Childlessness, at least in the older cohorts, was still fairly uncommon, with about 6% of the oldest respondents reporting that they had had no live births. Proportions of women with any live births and mean numbers of births were slightly, but consistently, lower among urban women than among rural women across cohorts.

Table 4.2 is identical to Table 4.1, except that it shows numbers of pregnancies rather than live births. Only about 9% of 15-19 year-olds reported ever having been pregnant, but among 20-24 year-olds, the figure was 61%. Because of the high levels of induced abortion, the mean numbers of pregnancies were far higher than the number of live births, starting at ages 25-29. In the three oldest cohorts, fewer than 5% of women reported that they had never been pregnant. The proportions ever pregnant were consistently higher among rural women than among urban women.

The total fertility rate (TFR) (i.e., the mean number of children per woman based on current age-specific fertility) according to survey data was about 1.4 births per woman for the two years preceding the survey (Table 4.3). This was not substantially different from the official figure for Ukraine of 1.3 births. As in almost all of Europe, Ukraine's total fertility rate is well below the replacement level of 2.1 births per woman. Age-specific fertility rates conform to the typical eastern European pattern, whereby childbearing tends to start and stop at a much earlier age than in other parts of Europe. Fertility is heavily concentrated in the early years of marriage, with 81% of the TFR accounted for by births occurring before age 30 (Figure 4.2). By far the highest fertility is at ages 20-24, where 11.5% of women bear children in a one-year period, almost twice as high as any other age group.

The TFR in rural areas was almost 50% higher than in urban areas, 1.8 compared to 1.3 births per woman. The only age group in which there was not a substantial difference between urban and rural areas was at ages 25-29. There was considerable variation in fertility levels across Ukraine's regions. The TFR ranged from 2.0 in the western region, which is the most rural and the most ethnically Ukrainian part of the country, to 1.1 in the East, which is the most industrialized and most ethnically Russian region. The central northern, and southern regions all have fertility rates of 1.4 to 1.5. In all regions the age pattern of childbearing is similar, heavily concentrated in the twenties and extremely low after ages 30-34. Fertility was much lower among women with any post-secondary education (TFR=1.1) than among women with less formal education.

Table 4.4 shows that the total pregnancy rate (TPR) for the two years prior to the survey was 3.1 for Ukraine as a whole, meaning women average just over three pregnancies apiece during their lifetime, just over twice the total fertility rate. The age pattern of pregnancy is quite similar to that for fertility, but the age-specific pregnancy rates are highest relative to the fertility rates at the oldest ages. This is an indication that most abortions take place after women have completed childbearing, rather than to delay childbearing. Despite having the highest fertility rates, the West region has one of the lowest pregnancy rates, probably because of relatively low induced abortion incidence.

Table 4.5 demonstrates that childbearing continues to typically begin shortly after marriage for Ukrainian couples. Among women in union for less than five years, 82% had ever been pregnant and 69% had had a live birth. Only about one percent of the oldest cohort of women who were ever in union never became pregnant and never had a live birth. Even among those married as recently as 5-9 years earlier, 96% had ever been pregnant and 92% had had a live birth. Almost all figures were slightly higher among rural women than among urban women.

Table 4.6 presents percentage distributions of the number of live births to women according to age. The most interesting information in this table is the relatively small number of women with no live

births (at least at the older ages) and the similarly small numbers with large families. Women with more than three children have become quite rare and with the recent decline in fertility rates will, no doubt, become even rarer. Family sizes other than one or two children are becoming less common. Large families (more than two children) are considerably more common in rural areas than urban areas. Table 4.7 shows live birth distributions by years since first union and leads to conclusions similar to the previous table. Few women married 10 or more years report having no children.

Table 4.8 allows an examination of changes over time in the onset of childbearing. The percentage of women with births before particular ages has not changed substantially with the exception of somewhat of an increase in births before age 20, from 19% for the oldest cohort to 26% for 20-24 year-olds. Even though childbearing starts relatively early for most women, few women begin childbearing at very young ages (i.e., before about age 18), with births before age 16 almost nonexistent. Once again it is possible to see the concentration of childbearing beginning in the early twenties: about one-quarter reported a birth by age 20, but about three-quarters by age 25.

Table 4.9 displays the percentage distributions of numbers of pregnancies according to age at the time of interview. The general conclusions are similar to those seen with regard to live births. However, especially among the older cohorts, there are substantial numbers of women who have had a large number of pregnancies (about one-fourth of 40-44 year-olds had five or more pregnancies), a result of the very high abortion rates in effect, especially in previous decades.

Pregnancy Outcomes

Forty-five percent of pregnancies to respondents that ended since January 1994 resulted in a live birth and 7% reportedly ended in miscarriage or stillbirth (Table 4.10 and Figure 4.3). The remainder, just under half of pregnancies, were terminated by induced abortion. The ratio of conventional surgical abortions to miniabortions was slightly higher than 3:2 for Ukraine as a whole (with under 1% of pregnancies ending in abortions that were self-induced or otherwise performed without professional medical or surgical intervention). The proportion of pregnancies resulting in a live birth declined steadily with increasing age, from 59% for 15-25 year-olds to 21% among 35-44 year-olds. This again supports the contention that most couples still have their children shortly after marriage, while they are still young. The West has a much higher proportion of live births (65%) than elsewhere (37%-45%). Pregnancies to women with no previous live births were the most likely to result in a live birth (76%). Only 15% of pregnancies to women with two previously born children resulted in a live birth however, demonstrating the strong desire among most women to have no more than two children. The small group of women with three or more children were much more likely to have pregnancies end in a live birth. Rural pregnancies were more likely than urban pregnancies to result in a live birth. Differences in birth outcomes by educational attainment were small.

Planning status of pregnancies

Table 4.11 shows that fewer than half of pregnancies ending in 1996 or later (44%) were said by

respondents to have been planned at the time they occurred (i.e., the woman wanted to become pregnant at that time). Fifty-four percent were said to unplanned: 17% were mistimed (i.e., she wanted to wait longer to become pregnant) and 38% were unwanted (she wanted no more children at the time she became pregnant). Given such high levels of unintended pregnancy and widespread acceptance and availability of induced abortion, it is not surprising that there are high rates of abortion in Ukraine.

Since women/couples in Ukraine still tend to have a child or children shortly after marriage, it is not surprising that the likelihood of a pregnancy being planned fell sharply with increasing age, from 55% for 15-24 year-olds to only 20% for 35-44 year-olds (Figure 4.4). Likewise, pregnancies were more likely to be unintended as the number of living children rose to two. Among the relatively small number of women with three or more children the likelihood of the pregnancy being planned rose again. The proportion of pregnancies categorized as unwanted rose from 19% to 73%. Pregnancies were the most likely to be planned in the West (56%) and least likely in the East (36%). They were also more likely to be planned in rural areas than urban areas. There was little difference according to respondent's education. Of course, pregnancies ending in live birth were the most likely to be planned (80%) and those ending in abortion were the least likely (18%).

The most important point revealed in Table 4.12 (outcome according to planning status of the pregnancy) is that pregnancies classified as unwanted rarely (5% of the time) resulted in a live birth. In 92% of unwanted pregnancies, the pregnancy was terminated by abortion. In urban areas, only 3% of unwanted pregnancies resulted in a live birth. Mistimed pregnancies were much more likely to end in a live birth, 35% of the time. Planned pregnancies ended with a live birth 84% of the time, with 11% ending in miscarriage or stillbirth and 5% terminated by induced abortion.

Pregnancy Intentions

Respondents who reported that they were able to become pregnant were asked how many more children they would like to have. Overall, almost two-thirds (66%) of survey respondents said they wanted no more children (Table 4.13). As might have been expected, the proportions who wanted no more children increased sharply with the number of living children, from 11% of women with no children to 91% of women with two children and 87% of those with three or more children (Figure 4.5). Even among women with only one child, about one-half said they wanted no more, showing that one-child families have become a very acceptable option in Ukrainian society. Very few women who already had children reported that they wanted to have more than two children altogether. Among women with one living child, only 4% said they wanted at least two more. Among those with two living children, fewer than 1 % said they wanted any more. Only those without children yet were somewhat likely to report wanting at least three children (17%). In spite of the substantial differences noted in fertility levels between urban and rural areas of Ukraine, urban-rural differences in fertility preferences were very small.

Table 4.14 displays numbers of children planned at the time of interview compared with the number planned at the time respondents desired at the time of their first marriage. It is worth noting that the category of women wanting no children when they married is not included because there were only 10 such women, a dramatic demonstration of the persistence of a pronatalist attitude in Ukraine,

despite the extremely low level of childbearing in effect in recent years.

Figure 4.1
Total Fertility Rates, Most Recent Available
Selected Countries




Figure 4.2
Age-Specific Fertility, by Residence
1999 Ukraine Reproductive Health Survey




Figure 4.3
Percentage Distributions of Pregnancy Outcomes
Since January 1994, by Number of Living Children
1999 Ukraine Reproductive Health Survey

38




Figure 4.4
Percentage Distributions of Planning Status of Pregnancies
Since January 1994, by Number of Living Children
1999 Ukraine Reproductive Health Survey




Figure 4.5
Percent of Fecund Women in Union Who Want No More Children, According to Number of Living Children
1999 Ukraine Reproductive Health Survey




Table 4.1
 Percentage of respondents who have had one or more live births
 and mean number of live births, by age at interview, by residence
 1999 Ukraine Reproductive Health Survey

Age of respondent	Percent with any live births	Mean number of live births	Number of women
<u>All women</u>			
15-19	5.6	0.1	1,100
20-24	50.3	0.6	1,182
25-29	84.0	1.2	1,227
30-34	91.4	1.5	1,195
35-39	93.1	1.7	1,246
40-44	94.3	1.8	1,178
Total	69.1	1.2	7,128
<u>Urban women</u>			
15-19	5.1	0.1	861
20-24	45.7	0.5	919
25-29	81.8	1.1	941
30-34	91.2	1.4	908
35-39	92.4	1.6	990
40-44	93.4	1.7	925
Total	68.1	1.1	5,544
<u>Rural women</u>			
15-19	7.2	0.1	239
20-24	62.9	0.7	263
25-29	89.2	1.4	286
30-34	91.8	1.7	287
35-39	95.2	1.9	256
40-44	96.8	2.0	253
Total	74.7	1.2	1,584

Table 4.2
 Percentage of respondents who have had one or more pregnancies
 and mean number of pregnancies, by age at interview, by residence
 1999 Ukraine Reproductive Health Survey

Age of respondent	Percent with any pregnancies	Mean number of pregnancies	Number of women
<u>All women</u>			
15-19	9.0	0.1	1,100
20-24	61.1	1.0	1,182
25-29	90.1	2.1	1,227
30-34	95.1	2.8	1,195
35-39	96.4	3.2	1,246
40-44	96.5	3.6	1,178
Total	74.8	2.1	7,128
<u>Urban women</u>			
15-19	8.5	0.1	861
20-24	57.5	0.9	919
25-29	89.1	2.0	941
30-34	85.5	2.8	908
35-39	96.0	3.1	990
40-44	96.1	3.6	925
Total	73.6	2.1	5,544
<u>Rural women</u>			
15-19	9.8	0.1	239
20-24	70.8	1.2	263
25-29	92.5	2.3	286
30-34	95.4	2.9	287
35-39	97.6	3.3	256
40-44	97.6	3.6	253
Total	78.1	2.2	1,584

Table 4.3
Age-specific and total fertility rates*, by residence, region, and education
1999 Ukraine Reproductive Health Survey

Age of respondent	Total	Residence		Region					Education		
		Urban	Rural	North	Central	East	South	West	Low	Med.	High
15-19	.049	.044	.064	.036	.061	.043	.045	.067	.034	.057	.032
20-24	.115	.098	.158	.131	.147	.091	.102	.138	.132	.125	.077
25-29	.066	.067	.065	.073	.056	.046	.068	.091	.069	.062	.078
30-34	.036	.030	.052	.038	.023	.021	.050	.054	.042	.038	.030
35-39	.014	.012	.021	.022	.005	.009	.009	.028	.045	.014	.011
40-44	.004	.002	.008	.000	.000	.003	.002	.013	.000	.005	.001
Total Fertility Rate	1.42	1.27	1.83	1.50	1.46	1.06	1.38	1.96	1.61	1.51	1.14

*All rates are annual rates, based on reports for the two-year period preceding the date of interview.

Table 4.4
 Age-specific and total pregnancy rates*, by residence, region, and education
 1999 Ukraine Reproductive Health Survey

Age of respondent	Total	Residence		Region					Education		
		Urban	Rural	North	Central	East	South	West	Low	Med.	High
15-19	.059	.052	.078	.046	.076	.052	.056	.075	.053	.064	.042
20-24	.203	.181	.260	.225	.279	.174	.213	.187	.209	.228	.128
25-29	.165	.170	.153	.183	.153	.151	.182	.166	.205	.161	.165
30-34	.111	.113	.104	.116	.103	.104	.139	.099	.070	.122	.084
35-39	.050	.052	.045	.065	.023	.051	.061	.043	.149	.053	.031
40-44	.025	.024	.027	.018	.006	.023	.051	.026	.009	.023	.037
Total Pregnancy Rate	3.06	2.97	3.33	3.26	3.20	2.77	3.51	2.97	3.47	3.25	2.43

*All rates are annual rates, based on reports for the two-year period preceding the date of interview.

Table 4.5
 Percent of women ever in union with any live births
 and percent ever pregnant, by years since first union and residence
 1999 Ukraine Reproductive Health Survey

Years since first union	Percent ever pregnant	Percent with any live births	Number of women
<u>All women</u>			
0-4	82.4	68.7	1,051
5-9	95.7	91.6	1,268
10-14	98.2	95.7	1,220
15-19	97.6	96.2	1,117
20+	99.2	98.7	921
Total	94.7	90.2	5,577
<u>Urban</u>			
0-4	81.2	66.3	815
5-9	95.7	91.1	961
10-14	98.1	95.5	935
15-19	97.3	95.9	861
20+	99.2	98.2	724
Total	94.3	89.4	4,296
<u>Rural</u>			
0-4	85.8	74.8	236
5-9	95.7	92.6	307
10-14	98.6	96.4	285
15-19	98.4	96.8	256
20+	99.7	99.7	197
Total	95.7	92.2	1,281

Table 4.6
 Percentage distribution of number of live births, by current age and residence
 1999 Ukraine Reproductive Health Survey

Number of live births	Current age of respondent						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
<u>Total</u>							
0	94.5	50.2	16.5	8.9	7.1	5.5	29.4
1	5.5	43.2	57.1	44.9	31.2	26.2	35.1
2	0.0	6.0	22.9	39.2	53.1	56.7	30.2
3	0.0	0.5	3.0	5.6	6.6	9.6	4.3
4+	0.0	0.1	0.6	1.4	2.1	1.9	1.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	1,100	1,182	1,227	1,195	1,246	1,178	7,128
<u>Urban</u>							
0	94.9	54.4	17.7	8.4	7.4	6.1	31.6
1	5.1	40.8	59.8	47.4	32.5	26.0	34.9
2	0.0	4.4	20.4	38.9	53.5	58.7	29.5
3	0.0	0.2	1.8	4.0	5.3	8.7	3.4
4+	0.0	0.2	0.3	1.3	1.4	0.5	0.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	861	919	941	908	990	925	5,544
<u>Rural</u>							
0	93.3	37.1	10.4	8.9	4.8	2.8	25.4
1	6.7	50.8	43.5	29.7	20.1	18.3	28.5
2	0.0	10.8	37.5	47.7	60.3	55.9	35.8
3	0.0	1.3	7.3	10.7	9.9	15.8	7.6
4+	0.0	0.0	1.2	2.9	4.9	7.2	2.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	239	263	286	287	256	253	1,584

Table 4.7
 Percentage distribution of number of live births, by number of years since first union, by residence
 1999 Ukraine Reproductive Health Survey

Number of live births	Number of years since first union					Total
	0-4	5-9	10-14	15-19	20+	
<u>All women</u>						
0	31.3	8.4	4.3	3.9	1.9	9.8
1	62.6	57.2	42.6	24.3	20.1	42.1
2	5.7	30.3	44.7	60.8	60.2	40.0
3+	0.4	4.1	8.5	11.1	17.8	8.1
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>1,051</i>	<i>1,268</i>	<i>1,220</i>	<i>1,117</i>	<i>921</i>	<i>5,577</i>
<u>Urban</u>						
0	33.7	8.9	4.5	4.1	2.5	10.6
1	61.9	62.3	48.9	27.2	21.8	45.2
2	4.3	26.6	40.2	60.4	61.9	38.3
3+	0.2	2.2	6.4	8.4	13.8	5.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>815</i>	<i>961</i>	<i>935</i>	<i>861</i>	<i>724</i>	<i>4,296</i>
<u>Rural</u>						
0	25.2	7.4	3.6	3.2	0.5	7.9
1	64.6	45.1	27.5	16.9	15.6	34.4
2	9.3	38.9	55.5	61.7	55.3	44.4
3+	0.9	8.6	13.4	18.2	28.6	13.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>236</i>	<i>307</i>	<i>285</i>	<i>256</i>	<i>197</i>	<i>1,281</i>

Table 4.8
 Percentage of women who had a live birth before selected ages, by current age
 1999 Ukraine Reproductive Health Survey

Live birth before:	Current age of respondent						
	15-19	20-24	25-29	30-34	35-39	40-44	15-44*
Age 16	0.1	0.0	0.1	0.0	0.2	0.1	0.1
Age 18	4.2	5.5	5.3	2.8	3.7	2.9	4.0
Age 20	--	25.7	26.0	22.8	20.9	19.0	22.9
Age 25	--	--	76.4	77.9	74.0	73.8	75.5
Age 30	--	--	--	89.7	89.2	88.1	89.0
Median age at first birth	NA	NA	21.8	21.8	22.0	22.1	--
<i>Number of women</i>	1100	1182	1227	1194	1246	1176	7,125**

*Table only includes women who have reached the age in question.

**Data missing for 3 women.

Table 4.9
 Percentage distribution of number of pregnancies, by current age, by residence
 1999 Ukraine Reproductive Health Survey

Number of pregnancies	Current age of respondent						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
<u>All women</u>							
0	90.9	38.6	9.7	4.9	3.6	3.3	25.1
1	7.5	34.8	28.6	15.5	10.6	8.0	17.4
2	1.4	18.2	29.5	28.9	23.8	20.2	20.4
3	0.1	5.8	18.7	22.6	26.1	24.5	16.4
4	0.0	1.5	7.8	14.6	15.9	18.9	9.8
5	0.0	0.6	2.9	6.8	10.8	8.2	4.9
6	0.0	0.2	1.9	3.0	3.7	6.0	2.5
7	0.0	0.3	0.2	1.7	1.9	4.7	1.5
8+	0.0	0.0	0.7	1.9	3.5	6.2	2.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>1,100</i>	<i>1,182</i>	<i>1,227</i>	<i>1,195</i>	<i>1,246</i>	<i>1,178</i>	<i>7,128</i>
<u>Urban</u>							
0	91.2	42.1	10.6	5.0	4.0	3.6	26.3
1	7.3	33.4	31.1	17.0	11.4	8.9	18.0
2	1.3	16.6	28.5	28.4	23.9	18.3	19.4
3	0.2	5.4	17.0	22.9	25.9	24.2	15.9
4	0.0	1.3	7.3	13.1	15.1	19.6	9.4
5	0.0	0.8	2.7	6.7	10.3	7.9	4.8
6	0.0	0.3	1.8	2.9	3.8	6.2	2.5
7	0.0	0.2	0.2	1.5	2.3	4.6	1.5
8+	0.0	0.0	0.7	2.3	3.1	6.7	2.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>861</i>	<i>919</i>	<i>941</i>	<i>908</i>	<i>990</i>	<i>925</i>	<i>5,544</i>
<u>Rural</u>							
0	90.3	28.8	7.5	4.7	2.4	2.4	21.9
1	8.1	38.7	22.3	12.2	8.3	5.5	15.9
2	1.6	22.8	31.8	30.4	23.4	25.1	22.9
3	0.0	7.0	22.8	21.7	26.7	25.4	17.5
4	0.0	2.1	8.9	17.9	18.4	16.9	10.9
5	0.0	0.2	3.7	6.7	12.3	8.9	5.3
6	0.0	0.0	2.3	3.3	3.4	5.7	2.5
7	0.0	0.4	0.0	2.2	0.5	5.2	1.4
8+	0.0	0.0	0.7	1.1	4.6	4.8	1.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>239</i>	<i>263</i>	<i>286</i>	<i>287</i>	<i>256</i>	<i>253</i>	<i>1,584</i>

Table 4.10
 Percentage distributions of outcomes of pregnancies ending since January 1994, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Pregnancy outcome							Total	Pregs.
	Live birth	Stillbirth	Miscarriage	Induced abortion					
				Total	Conventional	Miniabortion	Self-induced		
Total	45.3	0.4	7.1	47.2	28.7	17.8	0.7	100.0	2,802
Age									
15-24	59.3	0.5	7.5	32.7	20.5	11.0	1.2	100.0	913
25-34	43.9	0.3	6.9	48.8	29.1	19.2	0.5	100.0	1,443
35-44	20.7	0.3	6.8	72.2	44.4	27.5	0.3	100.0	446
Region									
North	44.9	0.3	5.3	49.5	22.5	24.9	2.1	100.0	336
Central	44.4	1.2	7.0	47.3	25.1	21.8	0.4	100.0	243
East	36.7	0.5	6.8	56.0	38.5	17.1	0.4	100.0	1,029
South	39.8	0.1	8.2	52.0	33.5	17.8	0.7	100.0	754
West	65.3	0.0	8.1	26.6	14.8	11.6	0.2	100.0	440
Residence									
Urban	41.5	0.5	6.4	51.7	30.5	20.4	0.8	100.0	2,162
Rural	54.8	0.2	8.9	36.1	24.2	11.4	0.5	100.0	640
Living children*									
0	75.9	0.7	8.3	15.1	8.9	5.1	1.1	100.0	985
1	35.0	0.2	7.8	57.0	34.7	21.7	0.6	100.0	1,126
2	14.8	0.2	3.5	81.5	47.4	33.8	0.3	100.0	561
3+	42.9	0.0	8.2	48.9	40.3	8.6	0.0	100.0	130
Education									
< Comp. secondary	47.8	1.0	10.5	40.7	26.9	13.3	0.5	100.0	247
Comp. secondary	45.7	0.3	6.9	47.2	29.6	16.8	0.8	100.0	2,019
> Comp. secondary	42.8	0.4	6.3	50.6	26.3	23.8	0.5	100.0	536

*Living children at the time of the pregnancy

Table 4.11
 Percentage distribution of planning status of pregnancies ending since January 1994,
 by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Planning status of pregnancy				Total	Number of pregs.
	Planned	Mistimed	Unwanted	Unsure		
Total	43.6	16.6	37.5	2.3	100.0	2,802
Age						
15-24	55.4	23.6	18.8	2.2	100.0	913
25-34	43.3	15.8	38.5	2.5	100.0	1,443
35-44	19.9	5.0	72.9	2.2	100.0	446
Region						
North	42.3	16.9	39.4	1.5	100.0	336
Central	46.5	13.6	38.7	1.2	100.0	243
East	36.9	17.5	43.2	2.4	100.0	1,029
South	41.2	19.8	36.6	2.4	100.0	754
West	56.2	13.7	26.9	3.4	100.0	440
Residence						
Urban	39.7	17.8	39.9	2.5	100.0	2,162
Rural	53.2	13.6	31.4	1.8	100.0	640
Living children*						
0	74.3	19.3	3.6	2.8	100.0	985
1	34.6	22.6	40.6	2.1	100.0	1,126
2	12.2	3.8	82.5	1.6	100.0	561
3+	32.8	2.8	60.6	3.8	100.0	130
Education						
< Comp. secondary	45.7	18.9	32.3	3.1	100.0	247
Comp. secondary	43.3	16.1	38.3	2.3	100.0	2,019
> Comp. secondary	43.5	17.5	36.9	2.2	100.0	536
Pregnancy outcomes						
Live birth	80.2	12.7	4.4	2.7	100.0	1,251
Stillbirth/miscarriage	75.8	12.3	10.3	1.6	100.0	204
Abortions	18.0	21.1	59.7	1.2	100.0	1,347

* Living children at the time that pregnancy occurred

Table 4.12
 Percentage distribution of outcomes of pregnancies ending since January 1994,
 by planning status and by residence
 1999 Ukraine Reproductive Health Survey

Planning status / Residence	Outcome of pregnancy			Total	Number of pregnancies
	Live birth	Stillbirth / Miscarriage	Induced abortion		
<u>Total</u>					
Planned	83.5	11.5	5.0	100.0	1,236
Mistimed	34.6	6.0	59.5	100.0	459
Unwanted	5.3	3.2	91.5	100.0	1,045
Unsure	52.7	9.9	37.4	100.0	62
<u>Urban</u>					
Planned	83.6	11.1	5.2	100.0	893
Mistimed	32.8	5.7	61.4	100.0	373
Unwanted	3.1	3.0	93.9	100.0	846
Unsure	50.0	7.7	42.3	100.0	50
<u>Rural</u>					
Planned	83.2	12.2	4.5	100.0	343
Mistimed	40.4	6.5	53.1	100.0	86
Unwanted	12.5	4.1	83.3	100.0	199
Unsure	*	*	*	100.0	12

*Fewer than 25 pregnancies in category

Table 4.13
 Number of additional children desired, by number of living children, by residence
 among fecund women currently in union
 1996 Ukraine Reproductive Health Survey

Living children	Additional children desired					Total	Number of women
	0	1	2	3+	Unsure		
<u>Total</u>							
0	11.1	32.4	30.6	16.6	9.4	100.0	459
1	49.3	28.9	1.6	2.2	18.1	100.0	1,952
2	92.1	1.8	0.1	0.7	5.3	100.0	1,799
3+	86.8	1.9	0.0	5.5	5.8	100.0	300
Total	66.3	15.9	3.6	3.2	11.0	100.0	4,510
<u>Urban</u>							
0	11.9	33.3	29.5	15.9	9.4	100.0	373
1	50.7	27.9	1.4	1.9	18.1	100.0	1,567
2	92.1	2.1	0.1	0.7	4.9	100.0	1,305
3+	90.6	0.2	0.0	3.9	5.2	100.0	172
Total	65.5	16.7	3.7	2.9	11.2	100.0	3,417
<u>Rural</u>							
0	8.6	29.6	33.8	18.7	9.3	100.0	86
1	44.9	31.6	2.2	3.3	18.0	100.0	385
2	92.2	1.0	0.2	0.6	5.9	100.0	494
3+	83.0	3.6	0.0	7.1	6.3	100.0	128
Total	68.4	14.1	3.4	3.7	10.4	100.0	1,093

NOTE: Currently pregnant women are considered to have one more living child than they actually do.

Table 4.14
 Percentage distribution of total number of children desired,
 by number desired at time of first marriage, by residence, fecund women currently in union
 1996 Ukraine Reproductive Health Survey

Number of children desired at time of first union*	Total number of children currently desired, including currently living children					Total	Number of women
	0	1	2	3+	Unsure		
<u>Total</u>							
1	3.1	62.9	29.8	2.3	1.9	100.0	853
2	1.1	21.9	69.7	5.9	1.4	100.0	2,189
3+	0.3	16.1	51.1	29.3	3.2	100.0	268
Unsure	2.9	29.3	47.9	13.4	6.5	100.0	1,122
<u>Urban</u>							
1	3.4	65.5	27.2	1.7	2.2	100.0	672
2	1.5	23.6	68.7	5.0	1.2	100.0	1,632
3+	0.4	17.3	51.6	28.7	2.0	100.0	195
Unsure	2.8	34.1	48.7	9.0	5.4	100.0	857
<u>Rural</u>							
1	2.2	55.9	36.8	3.9	1.1	100.0	181
2	0.0	18.1	72.0	8.2	1.7	100.0	557
3+	0.0	13.7	49.9	30.7	5.8	100.0	73
Unsure	3.5	18.2	46.0	23.5	8.8	100.0	265

*Only ten respondents desired to have no children when married. These respondents are not included in the table.

CHAPTER V

INDUCED ABORTION

The incidence of induced abortion in Ukraine, as in most of the former Soviet republics, has been very high in recent decades (Popov, 1991; Blayo, 1993). However, official statistics have revealed that rates have been declining in recent years. Figure 5.1 shows the trend in induced abortion rates in recent years in Ukraine, according to official Ministry of Health statistics. (These figures have been adjusted to eliminate miscarriages and stillbirths, which are usually combined with induced abortions when official statistics are compiled.) It is readily seen that the trend in abortion rates has been steadily downward since at least the middle of the 1980s. Between 1990 and 1998, the official rate of abortion incidence fell by over 50%, from 77 abortions to about 36 abortions per 1,000 women of childbearing age. Figure 5.2 shows abortion rates based on recent survey data for several countries in eastern Europe and the former Soviet Union. Abortion rates in these countries have been among the highest in the world for several decades. Although the rate for Ukraine is considerably lower than in some other countries in the region, it is still quite high by international standards. As we will see below, abortion remains perhaps the most important means of birth prevention in Ukraine, despite the recent decline in incidence.

Proportions of women with any abortions

By any measure, induced abortion is a common procedure in Ukraine. The simplest way to look at the incidence of abortion is to examine the proportion of women who undergo the procedure. Overall, 43% of respondents had ever had an induced abortion of any type (Table 5.1). Twenty percent of respondents had had more than one abortion. These figures tend to understate the incidence of abortion, however, since many younger women have not yet been at risk or have been at risk for a relatively short time for unintended pregnancy or, therefore, abortion. Among women in the oldest cohorts (ages 35-39 and 40-44), about two-thirds had had at least one abortion and about one-third had had at least two abortions during their lifetime (Table 5.1). Women between 40 and 44 years of age averaged 1.6 abortions. However, since most of the abortions to women in the oldest cohorts did not take place recently, these figures may not be good indicators of recent abortion activity. Few females under the age of 20 reported having had any induced abortions. Percentages of women with abortions, as well as mean numbers of abortions, were consistently higher in urban areas than in rural areas.

Table 5.2 shows that the percentages of women with any abortions differ relatively little between the North, Central, South, and East regions of the country, but the percentage is far lower in the West, the part of the country that is the most ethnically Ukrainian and the most rural; only 28% of women reported any abortions (compared to 43% nationally) and 9% reported having more than one abortion (compared to 20% nationally). Women with the least education were less likely than other women to have had abortions. Abortions were the least common among ethnically Ukrainian women who spoke primarily Ukrainian and were highest among women describing themselves as ethnically Russian.

Although the largest numbers of women reported having had only zero or one abortions, many respondents reported having multiple abortions (Table 5.3). Three percent of all women and 8% of 40-44 year-olds listed 5 or more lifetime abortions.

Current incidence of abortion

The total induced abortion rate (i.e., the mean number of lifetime abortions per woman based on current age-specific abortion rates) for Ukraine as a whole was about 1.6 abortions per woman for the two years before interview (Table 5.4). The abortion rate (i.e., the probability that a woman reported having an abortion during the previous 12 months) was .054, meaning that just over one of every 20 women have an abortion during a one-year period. The abortion ratio (i.e., the ratio of induced abortions to live births) was 1.10, indicating about equal numbers of abortions and live births. All of these figures were substantially higher in urban areas than in rural areas. As with fertility, age-specific abortion rates were highest among women in their twenties (.091), followed by women 30-34 years of age (.069) (Figure 5.2).

It should be noted that the induced abortion rate according to the survey (.054) was about 30 percent higher than the official rate published by the Ministry of Health (about .039 for the same period). This is an indication that the official statistics miss a significant proportion of abortions that are occurring, possibly because many abortions are being performed by providers who are not reporting them for any number of reasons. It also is an indication that reporting of abortion experience among URHS respondents tended to be relatively complete. If, in fact, the official rates of abortion are somewhat underreported, it makes it difficult to draw conclusions on the actual trends in abortion rates in recent years.

Table 5.5 shows abortion indicators according to region and respondents' educational attainment. Abortion levels were slightly higher in the South than elsewhere (TAR=2.05, compared to 1.57 nationally) and much lower in the West, where the rate was only 0.77 abortions per woman. Abortion rates tended to be slightly lower among the best educated women than among other Ukrainian women.

Table 5.6 displays total pregnancy, fertility, and abortion rates according to selected characteristics of respondents. During the two years prior to interview, the total pregnancy rate was barely above three pregnancies per woman, with slightly higher rates of abortions than live births. The total abortion rate in Donetsk Oblast, one of the UWRHI project sites, was slightly above the national rate. The rate in Odessa was about the same as the national rate.

Types of abortions, complications, and cost

Until fairly recently, almost all induced abortions in Ukraine consisted of procedures that would be considered "conventional abortions" in the West. However, many abortions now performed are what are commonly referred to as "miniabortions". This procedure, also sometimes referred to as "menstrual regulation", is performed using vacuum aspiration early in pregnancy. It tends to be a simpler, more easily performed procedure than those employed for conventional abortions. Of all abortions undergone by survey respondents since the beginning of 1994, 61% were reported to be conventional abortions and 38% were miniabortions (Table 5.7). An additional 1% were reported as self-induced or otherwise performed outside of a medical setting. The distribution of types of abortion seems to be stable, with similar distributions in 1994-1996 and 1997-1999. Miniabortions were most common among 25-34 year-old women (39%), in the North region (54%), in urban areas (39%), and among the best educated women (49%). They were least common relative to conventional procedures in the East region (30%).

Women were asked about complications and health problems associated with each of their recent induced abortions (including miniabortions) since the beginning of 1994, both “soon after” and at least six months after the procedure. Fourteen percent of abortions performed since January 1994 resulted in what women described as “complications requiring medical treatment” immediately or soon after the procedure (Table 5.8, left-hand panel). As might have been expected, conventional abortions were about 50% more likely than miniabortions to result in short-term complications (16% vs. 11%). Although the differences were not statistically significant, complication rates were slightly higher among urban women than rural women for each type of abortion. When interpreting these data, it should be kept in mind that what constitutes a “complication” is subjectively defined by the respondent and not by medical personnel.

Among those women experiencing complications, 37% (5% of all women undergoing abortions) reported rehospitalization or extended hospitalization as a result, with a higher likelihood of hospitalization following conventional abortions (39% vs. 32%) (Table 5.8, center panel). Six percent of abortions were reported to have resulted in long-term problems, with the likelihood of problems twice as high from conventional abortions as from miniabortions (Table 5.8, right-hand panel). There were no notable differences in long-term complication rates between urban and rural areas.

When asked why they decided to have an abortion, the overwhelming majority of women gave as their primary reason that they wanted no more children (60%) or gave what were considered to be “social/economic reasons”, i.e., such reasons as inadequate resources or inadequate housing (25%) (Table 5.9). Smaller numbers of abortions were attributed to such factors as: continuing the pregnancy presented a health risk for the woman (4%), not being married (4%), the woman’s partner did not want a child (2%), and a risk of birth defects (2%). The percentage wanting no more children, not surprisingly, rose with age. It was also the lowest in the South region. Social/economic reasons were most often given by the least well educated women, whom it is assumed could least afford larger families.

Thirteen percent of abortions occurring in 1997 or later were free of charge to women (Table 5.10). An additional 14% were paid for with goods or services, rather than with money. Thus, about 7 of every 10 abortions were paid for with money. The vast majority of these (63% of all abortions) were reported to have cost the equivalent of less than US \$30. Only 4% cost more than US \$50. There were noteworthy differences in payment according to women’s residence. Rural women were more likely than urban women not to pay for their abortions and tended to pay less when they were not free. Abortions in the Central and East regions were more likely than in other regions to be free. As education rose, so did the amount women paid for abortions.

Thirteen percent of women with abortions since January 1994 reported that they received no anesthesia in association with their most recent abortion (Table 5.11). The likelihood of receiving anesthesia appears to have remained unchanged in recent years. Women having miniabortions were about twice as likely as those having conventional abortions not to have had anesthesia. Older women and urban women were less likely than others not to receive anesthesia. Women in the East region were the most likely to be anesthetized (91%), while women in the Central region were the least likely (77%).

Figure 5.1

**Induced Abortion Rates per 1,000 Women 15-44 Years,
1985-1998, Official Statistics, Ukraine**




Figure 5.2
Age-Specific Abortion Rates, by Place of Residence
1999 Ukraine Reproductive Health Survey

59




Table 5.1
 Percentage of respondents who have had any induced abortions or two or more induced abortions
 and mean number of induced abortions, by age at interview, by residence
 1999 Ukraine Reproductive Health Survey

Age of woman	Percent with 1 or more abortions	Percent with 2 or more abortions	Mean number of abortions	<i>Number of women</i>
<u>All women</u>				
15-19	1.8	0.2	0.0	1,100
20-24	22.8	5.5	0.3	1,182
25-29	44.3	16.7	0.7	1,227
30-34	57.6	27.8	1.1	1,195
35-39	65.5	32.0	1.3	1,246
40-44	67.2	37.2	1.6	1,178
Total	43.3	20.0	0.8	7,128
<u>Urban women</u>				
15-19	2.2	0.3	0.0	861
20-24	23.7	6.0	0.3	919
25-29	46.9	18.3	0.8	941
30-34	59.6	29.5	1.2	908
35-39	67.3	33.7	1.4	990
40-44	70.7	41.0	1.7	925
Total	45.1	21.5	0.8	5,544
<u>Rural women</u>				
15-19	0.8	0.0	0.0	239
20-24	20.4	4.2	0.3	263
25-29	37.9	12.7	0.6	286
30-34	52.8	24.0	0.9	287
35-39	60.1	27.0	1.2	256
40-44	57.6	26.7	1.2	253
Total	38.7	15.9	0.6	1,584

Table 5.2
 Percentage of respondents who have had any induced abortions or two or more induced abortions
 and mean number of induced abortions, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Age of woman	Percent with 1 or more abortions	Percent with 2 or more abortions	Mean number of abortions	<i>Number of women</i>
<u>All women</u>	43.3	20.0	0.8	7,128
<u>Residence</u>				
Urban	45.1	21.5	0.8	5,544
Rural	38.7	15.9	0.6	1,584
<u>Region</u>				
North	44.0	16.8	0.8	858
Central	48.8	22.9	0.9	617
East	47.6	23.3	1.0	2,669
South	49.2	27.0	1.1	1,824
West	27.7	9.2	0.4	1,160
<u>Education</u>				
<Complete secondary	20.8	10.8	0.4	778
Complete Secondary	46.7	22.1	0.9	4,828
>Complete secondary	44.6	18.0	0.8	1,522
<u>Language at home/Ethnicity</u>				
Ukrainian.-speaking/Ukrainian	38.4	15.2	0.7	2,157
Russian-speaking/Ukrainian	41.9	19.5	0.8	2,071
Other, Mixed/Ukrainian	46.5	23.2	1.0	839
Russian nationality	51.9	27.3	1.2	1,688
Other nationality	43.9	20.2	0.9	373

Table 5.3
 Percentage distribution of number of abortions, by current age, by residence
 1999 Ukraine Reproductive Health Survey

Number of abortions	Current age of respondent						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
<u>All Women</u>							
0	98.2	77.2	55.8	42.5	34.3	32.5	56.6
1	1.7	17.3	27.6	29.8	33.7	30.1	23.4
2	0.2	4.1	10.5	15.1	16.8	17.2	10.7
3	0.0	0.9	4.1	6.9	7.4	7.8	4.5
4	0.0	0.4	1.1	3.4	3.5	4.9	2.2
5-9	0.0	0.1	0.8	2.3	4.0	6.9	2.4
10+	0.0	0.0	0.3	0.2	0.4	0.8	0.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>1,100</i>	<i>1,182</i>	<i>1,227</i>	<i>1,195</i>	<i>1,246</i>	<i>1,178</i>	<i>7,128</i>
<u>Urban</u>							
0	97.8	76.3	53.2	40.2	32.5	28.8	54.8
1	1.9	17.7	28.5	30.1	33.8	29.7	23.6
2	0.3	4.2	11.9	15.1	17.6	18.4	11.3
3	0.0	1.0	3.9	7.9	7.7	8.7	4.9
4	0.0	0.6	1.4	3.4	3.8	6.1	2.5
5-9	0.0	0.2	0.9	2.8	4.4	7.5	2.6
10+	0.0	0.0	0.2	0.2	0.3	0.8	0.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>861</i>	<i>919</i>	<i>941</i>	<i>908</i>	<i>990</i>	<i>925</i>	<i>5,544</i>
<u>Rural</u>							
0	99.2	79.7	62.1	47.2	39.5	42.4	61.3
1	0.8	16.2	25.2	28.9	33.5	30.9	22.8
2	0.0	3.6	7.0	14.8	14.5	13.8	9.1
3	0.0	0.6	4.5	4.9	6.5	5.2	3.7
4	0.0	0.0	0.5	3.2	2.7	1.7	1.4
5-9	0.0	0.0	0.4	1.1	2.9	5.2	1.6
10+	0.0	0.0	0.4	0.0	0.4	0.8	0.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>239</i>	<i>263</i>	<i>286</i>	<i>287</i>	<i>256</i>	<i>253</i>	<i>1,584</i>

Table 5.4
Age-specific abortion rates and other measures of induced abortion*, by residence
1999 Ukraine Reproductive Health Survey

Age of respondent	Total	Urban	Rural
15-19	.013	.014	.012
20-24	.091	.090	.092
25-29	.091	.105	.070
30-34	.069	.080	.049
35-39	.033	.039	.019
40-44	.018	.020	.016
Total Abortion Rate	1.57	1.74	1.29
Abortion Rate**	.054	.058	.045
Abortion Ratio***	1.10	1.42	0.72

*All rates are annual rates, based on reports for the two-year period preceding the date of interview.

**Proportion of women 15-44 years of age having induced abortions in one year.

***Ratio of induced abortions to live births

Table 5.5
 Selected measures of induced abortion*, by region and education
 1999 Ukraine Reproductive Health Survey

Region/ Education	Total Abortion Rate	Abortion Rate**	Abortion Ratio***
<u>All women</u>	1.57	.054	1.10
<u>Region</u>			
North	1.76	.060	1.16
Central	1.60	.054	1.12
East	1.74	.058	1.63
South	2.05	.070	1.45
West	0.77	.027	0.39
<u>Education</u>			
<Comp secondary	1.82	.052	0.91
Comp. Secondary	1.62	.055	1.09
>Comp secondary	1.40	.048	1.22

*All rates are annual rates, based on reports for the two-year period preceding the date of interview.

**Proportion of women 15-44 years of age having induced abortions in one year.

***Ratio of induced abortions to live births

Table 5.6
Total fertility, pregnancy, and induced abortion rates*, by selected characteristics
1999 Ukraine Reproductive Health Survey

Characteristic	Total Rate of:		
	Pregnancy	Fertility	Induced Abortion
<u>All women</u>	3.06	1.42	1.57
<u>Residence</u>			
Urban	2.97	1.27	1.74
Rural	3.33	1.83	1.29
<u>Region</u>			
North	3.26	1.50	1.76
Central	3.20	1.46	1.60
East	2.77	1.06	1.74
South	3.51	1.38	2.05
West	2.97	1.96	0.77
<u>Education</u>			
<Comp secondary	3.47	1.61	1.82
Comp. secondary	3.25	1.51	1.62
>Comp secondary	2.43	1.14	1.40
<u>Oversampled oblasts</u>			
Donetsk (East)	2.66	1.06	1.71
Odessa (South)	2.88	1.30	1.59

*For the period 2 years prior to interview.

Table 5.7
 Percentage distribution of types of abortions, by selected characteristics,
 all abortions since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	Type of abortion			Total	<i>Number of abortions</i>
	Conventional	Miniabortion	Self-induced*		
Total	61.1	37.8	1.1	100.0	2,032
<u>Year</u>					
1994- 1996	62.9	36.2	0.8	100.0	1,054
1997 -1999	59.1	39.4	1.5	100.0	978
<u>Age</u>					
15-24	66.9	31.4	1.7	100.0	710
25-34	60.1	39.1	0.9	100.0	1,005
35-44	63.9	35.8	0.3	100.0	317
<u>Region</u>					
North	43.2	53.9	2.9	100.0	235
Central	56.9	42.5	0.6	100.0	181
East	69.6	29.8	0.6	100.0	855
South	61.5	37.4	1.1	100.0	583
West	58.2	40.8	1.1	100.0	178
<u>Residence</u>					
Urban	59.6	39.1	1.3	100.0	1,684
Rural	66.4	33.0	0.6	100.0	348
<u>Education</u>					
< Comp. secondary	69.2	29.0	1.8	100.0	149
Comp. secondary	63.3	35.6	1.1	100.0	1,474
> Comp. secondary	50.6	48.5	0.9	100.0	409

* Includes all abortions for which medical intervention was not used.

Table 5.8
 Percentage of induced abortions since January 1994 resulting in complications requiring medical treatment,
 and percentage of complications resulting in hospitalization, by type of abortion, by residence
 1999 Ukraine Reproductive Health Survey

Type of abortion	Complications requiring medical treatment "Soon after abortion"		Received additional hospitalization after abortion		Long-term health problems after abortion ***	
	%	N	%	N	%	N
<u>Total</u>						
All abortions*	14.1	2,032	37.4	272	5.7	2,004
Regular abortions	15.9	1,285	38.6	194	7.1	1,281
Miniabortions	10.9	723	31.9	73	3.3	717
<u>Urban</u>						
All abortions*	14.8	1,684	39.1	230	5.9	1,679
Regular abortions	16.9	973	40.3	164	7.4	1,050
Miniabortions	11.4	554	34.4	62	3.4	607
<u>Rural</u>						
All abortions*	12.1	348	30.0	42	5.2	343
Regular abortions	13.0	312	31.3	30	6.3	231
Miniabortions	9.5	169	**	11	3.0	110

NOTE: Abortions for which women did not remember whether they had complications have been excluded.

*Total does not equal sum of regular abortions and miniabortions because it includes self-induced abortions as well.

** Fewer than 25 abortions with problems requiring medical treatment

***At least six months after abortion

Table 5.9
 Percentage distribution of primary reason for abortion since January 1994,
 by age at abortion, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Reason for abortion							Total	Number of abortions
	Want no more children	Social/economic reason	Pregnancy dangerous to woman	Not married/no partner	Partner did not want child	Risk of fetal defects	Other		
Total	59.8	25.1	4.3	3.7	2.2	1.7	3.2	100.0	2,032
<u>Age</u>									
15-24	50.6	28.9	3.4	7.9	2.9	1.7	4.6	100.0	710
25-34	62.0	24.4	4.8	1.8	2.2	1.9	2.9	100.0	1,005
35-44	73.2	18.5	4.6	0.4	0.8	1.1	1.4	100.0	317
<u>Region</u>									
North	62.9	16.6	5.9	5.1	3.4	1.3	4.7	100.0	235
Central	63.5	24.9	3.9	2.8	1.7	0.6	2.8	100.0	181
East	61.9	26.2	3.1	3.2	1.4	1.3	2.8	100.0	855
South	50.4	29.7	5.8	3.7	4.3	2.9	3.1	100.0	583
West	61.2	24.0	3.8	4.4	0.6	2.7	3.3	100.0	178
<u>Residence</u>									
Urban	59.9	25.2	4.1	4.0	2.3	1.5	2.9	100.0	1,684
Rural	59.3	24.5	5.0	2.4	1.9	2.4	4.5	100.0	348
<u>Education</u>									
< Comp. secondary	48.7	32.4	7.3	5.6	3.9	0.0	2.2	100.0	149
Comp. secondary	62.9	24.5	3.8	3.1	1.6	1.0	3.0	100.0	1,474
> Comp. secondary	52.8	24.4	4.8	4.9	3.9	4.9	4.4	100.0	409

Table 5.10
 Percentage distribution of payment for most recent abortion since January 1997, by age at abortion, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Payment					Total	Number of abortions
	None	Monetary			Non monetary, goods/other		
		< \$30	\$ 31-50	> \$ 50			
Total	13.1	62.5	7.0	3.6	13.8	100.0	789
<u>Age</u>							
15-24	12.7	66.3	8.4	3.6	9.0	100.0	257
25-34	14.9	58.5	6.3	3.8	16.6	100.0	406
35-44	8.2	67.9	6.4	2.9	14.7	100.0	126
<u>Region</u>							
North	9.6	70.2	1.9	7.7	11.6	100.0	104
Central	23.2	62.3	1.5	0.0	13.0	100.0	69
East	16.9	58.3	6.4	1.7	16.7	100.0	329
South	6.5	63.2	14.6	6.1	9.6	100.0	221
West	7.1	64.3	8.6	2.9	17.1	100.0	66
<u>Residence</u>							
Urban	11.8	61.7	8.6	4.0	14.0	100.0	646
Rural	17.7	65.2	1.7	1.9	13.5	100.0	143
<u>Education</u>							
< Comp. secondary	28.3	60.6	3.8	1.8	5.6	100.0	63
Comp. secondary	12.9	61.7	6.4	3.3	15.7	100.0	556
> Comp. secondary	7.9	65.8	10.2	5.2	10.9	100.0	170

Table 5.11
 Percentage of women who received local anesthesia for their most recent abortion since January 1994,
 by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Anesthesia	No anesthesia	Don't remember	Total	Abortions
<u>Total</u>	86.1	13.2	0.7	100.0	1,430
<u>Year</u>					
1994	86.2	12.8	0.9	100.0	200
1995	86.0	13.7	0.3	100.0	204
1996	87.7	11.8	0.5	100.0	238
1997	85.1	14.5	0.5	100.0	266
1998	84.8	14.0	1.2	100.0	323
1999	87.7	12.0	0.3	100.0	199
<u>Type of abortion</u>					
Regular abortions	90.4	8.7	0.9	100.0	805
Miniabortions	80.9	18.6	0.5	100.0	491
<u>Age</u>					
15-24	85.2	14.6	0.3	100.0	492
25-34	85.7	13.4	0.9	100.0	710
35-44	89.3	9.9	0.8	100.0	228
<u>Region</u>					
North	85.9	13.5	0.6	100.0	178
Central	76.8	23.2	0.0	100.0	125
East	91.4	7.7	0.9	100.0	594
South	84.7	15.3	0.0	100.0	395
West	80.4	18.2	1.4	100.0	138
<u>Residence</u>					
Urban	88.3	11.2	0.5	100.0	1,164
Rural	78.9	19.9	1.2	100.0	266
<u>Education</u>					
< Comp. secondary	88.2	10.5	1.3	100.0	100
Comp. secondary	84.2	14.5	0.7	100.0	1,016
> Comp. secondary	89.6	10.0	0.4	100.0	314

CHAPTER VI

INFERTILITY

There can be little doubt that the typical desired family size has declined to very low levels and that abortion rates remain high. However, there has been some concern expressed in recent years that infertility may be playing an increasing role in the unprecedentedly low levels of fertility now in effect in Ukraine and other countries in the region. There are several factors that have suggested to researchers that there has been a growing inability among the population to have as many children as they would like. Among these factors are the increasing spread of sexually transmitted infections, high levels of chronic alcohol abuse (especially in males), and the high incidence of repeat abortions. All of these factors can potentially lead to a reduction in the ability to become pregnant or bear a child. There have also been anecdotal reports of increases in the numbers of women/couples seeking infertility treatment. However, to our knowledge, no definitive research has been conducted to determine the extent to which infertility has been suppressing rates of childbearing or to estimate the numbers of couples who are being affected by infertility.

The 1999 URHS included a module designed to collect information from respondents about their difficulties they have experienced in having children and any treatment they might have undergone in response to difficulties becoming pregnant or bearing a child. Although the survey data does not allow a thorough analysis of the extent and impact of infertility in Ukraine, it does provide the ability to examine certain aspects of the infertility situation, such as the proportion of couples who report problems becoming pregnant and the probability that couples receive treatments of various types.

Prevalence of infertility

Table 6.1 presents the proportion, among women who had ever tried to become pregnant, who said they had experienced problems becoming pregnant in the last 10 years, according to how long the problem lasted. Overall, almost one of every six respondents (16%) reported ever having such a problem. When the definition of an infertility problem is narrowed to longer durations, we still find that 14% had a problem lasting at least one year, 12% lasting at least two years and 7% had a problem lasting at least five years, seemingly relatively large percentages. The percentages reporting difficulty becoming pregnant were slightly higher than overall in urban areas and among the least well educated. Problems were less common in the West region.

Treatment

Overall, just over half of women who reported having difficulty becoming pregnant in the previous 10 years (52%) sought treatment for the problem (Table 6.2). In about two-thirds of those cases where treatment was sought, only the woman sought treatment. The proportion seeking treatment tended to increase with age, education of the respondent, and the length of the problem. The likelihood that both partners would seek treatment increased sharply with respondent's education.

The initial source of treatment for infertility problems overwhelmingly tended to be women's consultation clinics (72%) (Table 6.3). The remaining 28% sought treatment at an assortment of types of facilities, including 6% who went to private clinics or offices. Of those women who were treated,

29% went to more than one place. The secondary sources also consisted of a wide array of types of places, including 7% private clinics or offices. The best educated, the oldest, and urban women were the most likely to go to more than one type of facility and to utilize a private second source.

A broad assortment of infertility treatments were used by survey respondents or their partners, with the most common being the prescription of anti-inflammatory drugs, reported by 48% of such women (Table 6.4). Other treatments commonly mentioned were: hormones (24%), physiotherapy (19%), treatments for blockage of the Fallopian tubes (16%), and relaxation/spa therapy (15%). Less commonly used were more technologically advanced and more costly treatments requiring greater intervention, such as: laser therapy (5%), laparoscopy (5%), and in vitro fertilization (IVF) (2%). Every treatment listed, except for relaxation therapy was more commonly used by urban women than rural women and by the best educated than less well educated women. The differences are especially large for such treatments as IVF and laser therapy. Use of physiotherapy and relaxation therapy increased sharply with the age of the respondent.

About half of the women receiving treatment eventually became pregnant (Table 6.5). Forty percent had a live birth. In spite of the different types and places of treatment, there were only small, and statistically insignificant, differences in the likelihood of becoming pregnant or having a live birth by age, residence, education, and the duration of the problem. It is difficult to interpret this result. It could mean that many of the infertility problems tend to disappear themselves or are alleviated by relatively little intervention. It also may indicate that modern, more expensive, technologically advanced treatments are not yet in widespread enough use to have affected rates of treatment success.

Table 6.1
 Percentage of women who have ever had problem becoming pregnant in the previous 10 years,
 by duration of problem, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Duration of problem becoming pregnant				Number of women
	Ever	1+ year	2 + years	5+ years	
Total	15.9	13.9	12.3	6.7	2,528
<u>Age</u>					
15-19	32.2	14.2	6.3	0.0	45
20-24	30.6	24.3	19.9	3.7	322
25-29	26.6	23.5	20.9	11.8	507
30-34	15.9	15.4	14.5	9.4	533
35-39	8.0	7.7	7.1	5.5	577
40-44	3.3	3.3	3.3	2.9	544
<u>Region</u>					
North	17.3	15.6	13.5	6.6	289
Central	15.7	15.2	14.1	9.4	192
East	17.9	15.3	13.3	7.1	803
South	15.4	12.9	11.9	5.7	739
West	13.1	11.5	9.9	5.9	505
<u>Residence</u>					
Urban	16.4	14.3	12.9	6.9	1,886
Rural	14.7	12.9	10.9	6.0	642
<u>Education</u>					
< Comp. secondary	23.8	18.0	16.2	8.7	142
Comp. secondary	15.2	13.6	12.0	6.2	1,844
> Comp. secondary	16.1	13.7	12.1	8.0	542

Table 6.2
 Percentage of women with infertility problems who sought treatment,
 by age, region, residence, education and length of problem
 1999 Ukraine Reproductive Health Survey

Characteristic	Did not seek treatment	Person who sought treatment			Number of women
		Woman	Partner	Both	
Total	47.7	34.0	0.3	18.0	415
<u>Age</u>					
15-19	*	*	*	*	14
20-24	55.1	31.4	0.0	13.5	100
25-29	47.4	33.2	0.0	19.3	135
30-34	39.6	38.0	0.0	22.4	93
35-39	41.5	39.1	0.0	19.4	52
40-44	*	*	*	*	21
<u>Region</u>					
North	58.0	30.0	0.0	12.0	50
Central	40.0	26.7	0.0	33.3	30
East	52.6	30.7	0.9	15.8	151
South	44.8	35.9	0.0	19.3	118
West	37.9	43.9	0.0	18.2	66
<u>Residence</u>					
Urban	47.1	34.8	0.4	17.6	319
Rural	49.1	31.9	0.0	18.9	96
<u>Education</u>					
< Comp. secondary	54.2	36.5	0.0	9.3	33
Comp. secondary	48.1	34.9	0.4	16.5	286
> Comp. secondary	43.8	29.7	0.0	26.5	96
<u>Length of problem**</u>					
1 + years	42.6	36.2	0.0	21.2	361
2 + years	41.4	36.3	0.0	22.4	326
5 + years	37.6	35.5	0.0	26.9	179

*Fewer than 25 respondents in category.

**Does not include those who did not remember. Categories are not mutually exclusive, so total does not equal 415.

Table 6.3
 Percentage distribution of women/couples who sought for treatment of infertility problems,
 by initial and second place of treatment, by age, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	First place of treatment					Second place of treatment						Total	No. of women
	Women consult.	MCH center	FP center	Private clinic	Other	Women consult.	MCH center	FP center	Private clinic	Other	None		
Total	72.4	4.7	6.6	5.9	10.4	1.5	2.6	4.2	7.3	13.6	70.9	100.0	266
<u>Age</u>													
15-24	76.8	3.5	9.5	7.8	2.4	1.1	4.5	2.4	2.4	15.0	74.7	100.0	56
25-34	72.2	3.4	6.0	4.6	13.8	1.0	1.9	4.8	8.6	11.9	71.7	100.0	130
35-44	66.5	10.9	4.4	7.4	10.8	3.6	2.1	4.7	9.9	17.2	62.6	100.0	40
<u>Residence</u>													
Urban	77.4	3.6	5.5	7.4	6.1	0.8	2.8	3.3	9.7	14.5	68.9	100.0	176
Rural	59.6	7.4	9.5	2.1	21.4	3.1	2.1	6.4	0.9	11.3	76.0	100.0	50
<u>Education</u>													
< Comp. sec.	*	*	*	*	*	*	*	*	*	*	*	100.0	16
Comp. sec.	68.9	4.9	6.0	6.4	13.8	2.1	2.3	4.7	5.3	12.3	73.3	100.0	159
> Comp. sec.	83.6	5.3	5.3	2.3	3.5	0.0	4.2	3.9	14.7	21.2	56.1	100.0	51

Table 6.4
 Percentage of treated women/couples who received various types of treatment for infertility problems,
 by age, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Type of infertility treatment									Number of women
	Hormones	Anti-inflammatory drugs	Fallopian blockage treatment	Physiotherapy	Laser therapy	Laparoscopy/ Microsurgery	Relaxation/ Spa therapy	IVF	Other	
Total	24.2	48.0	15.5	18.9	4.7	4.7	14.9	2.2	18.8	226
<u>Age</u>										
15-24	21.4	62.3	12.5	14.1	5.2	1.8	7.2	0.0	17.7	56
25-34	26.8	46.8	16.2	19.1	6.1	5.3	15.3	3.1	16.3	130
35-44	20.0	32.6	17.5	25.0	0.0	7.4	24.9	2.4	29.9	40
<u>Residence</u>										
Urban	26.7	50.0	17.5	21.5	5.5	5.1	15.9	2.3	18.1	176
Rural	16.0	42.0	8.0	10.0	2.0	4.0	12.0	2.0	21.9	50
<u>Education</u>										
< Comp. sec	*	*	*	*	*	*	*	*	*	16
Comp. sec	22.0	44.0	14.4	16.5	3.8	3.9	12.6	0.6	21.0	159
> Comp. sec	31.4	58.7	21.5	25.4	7.6	9.6	7.8	7.7	15.6	51

NOTE: Row percentages add to more than 100 percent, since some women received more than one type of treatment.
 *Fewer than 25 women in category.

Table 6.5
 Percentage of women treated for infertility problems who eventually become pregnant and
 percent who had a live birth, according to selected characteristics and length of problem
 1999 Ukraine Reproductive Health Survey

Characteristic	Percent who become pregnant	Percent who had a live birth	<i>Number of women</i>
Total	52.8	40.4	226
<u>Age</u>			
15-24	53.7	37.3	56
25-34	51.3	41.7	130
35-44	56.7	40.6	40
<u>Residence</u>			
Urban	52.1	40.9	176
Rural	54.5	39.1	50
<u>Education</u>			
< Comp. secondary	*	*	16
Comp. secondary	50.5	40.5	159
> Comp. secondary	56.7	42.5	51
<u>Length of problem</u>			
1 + year	53.5	43.3	207
2 + years	54.5	45.6	188
5 + years	54.0	46.1	108

*Fewer than 25 respondents in category

CHAPTER VII

CONTRACEPTION

One of the principal reasons for carrying out the 1999 Ukraine Reproductive Health Survey was to perform an in-depth examination of contraceptive knowledge, attitudes, and practices among Ukrainian women. Prior to this survey, relatively little national systematic information existed on most aspects of contraceptive use in Ukraine (Vovk, 1997). Until recently, the conventional wisdom had been that the prevalence of use of modern contraception in eastern Europe and the former Soviet Union was quite low, leading to high levels of unintended pregnancy and induced abortion there. Recent surveys in other countries in the region, however, have shown that overall contraceptive use, as well as the use of effective modern contraceptive methods, is often very widespread, despite high rates of induced abortion. Thus, it was not surprising to find in the URHS that the use of family planning methods in Ukraine has reached a very high level. This survey collected information on a broad array of topics related to contraception, including knowledge and use of contraceptive methods, source of methods, contraceptive failure and discontinuation, side effects, and reasons for nonuse of contraception, among others. Virtually no respondents refused to provide information on any of the topics related to contraception about which they were asked.

Knowledge and ever use of contraceptive methods

In general, knowledge of the most readily available methods was widespread. Table 7.1 shows that virtually all women knew at least one modern contraceptive method (99.6%). Nearly all respondents said that they had heard of condoms (99% knowledge) and the IUD (96%), followed by oral contraceptives (90%). A majority of respondents also were familiar with female sterilization (67%), spermicides (60%), and the diaphragm (58%). The only modern method asked about for which knowledge remained very low was contraceptive implants, which remain unavailable to most Ukrainians, known by only 18% of respondents. Among non-supplied methods, both periodic abstinence and withdrawal were known by between 80% and 90% of women.

Knowledge of every method was higher in urban than in rural areas, but, for most methods these differences were relatively small, the differences being greatest for sterilization (female and male), spermicides, diaphragms, and implants. Not surprisingly, most contraceptive methods tended to be much more widely known by women currently or previously in union than those never in union. For most methods, knowledge rose with age up to ages 25-29 and remained relatively constant from 25-29 to 40-44 (Table 7.2). However, all age groups and education level groups had at least 97% knowledge of condoms. With the exception of condoms, knowledge increased sharply with educational attainment for every method asked about.

Knowledge of where to obtain supplied contraceptive methods tended to be almost as high as knowledge of methods, indicating that most women who had heard of a method also reported that they knew where they could obtain that method. Knowledge of method source ranged from 97% for condoms to 13% for implants. Differentials paralleled those for knowledge of methods: higher in urban women, better educated women, and women ever in union.

Overall, 74% of all respondents and 87% of those currently in union reported ever using any contraceptive method (Table 7.4). The figures for modern methods were 60% and 69%,

respectively. The methods most likely to have ever been used were withdrawal (47%), condoms (44%), periodic abstinence (38%), and the IUD (30%). With the exception of oral contraceptives (14%), all other methods had been practiced by very few women. Except for withdrawal, all other widely used methods were more likely to have been used by urban than by rural women. In general, there was very little difference between currently in union and previously in union women in regard to ever use. Ever use increased substantially with educational attainment, except for some rarely used methods.

Current Contraceptive Prevalence

About two-thirds (68%) of women in registered or unregistered marriages were currently using contraceptive methods at the time of interview (Figure 7.1 and Table 7.5). In addition, more women/couples were employing modern methods of contraception than traditional methods (periodic abstinence and withdrawal), though the difference was not great (38% and 30%, respectively). It should be noted that users of methods considered to be of very poor or no effectiveness, particularly douching and folk methods, were not considered to be users of contraception in these tabulations. Including such women would have raised the contraceptive prevalence rate by approximately 3 percentage points. IUDs (used by 19% of women in union) and condoms (14%) accounted for the vast majority of modern method use in Ukraine, with oral contraceptives (OCs) (3%) a distant third. Withdrawal (at 20%, the most widely practiced method overall) and periodic abstinence, i.e., rhythm, calendar, and related methods (10%), were also widely used. Only about 1% of women had been contraceptively sterilized, despite the fact that most respondents wanted to have no more children.

Contraceptive prevalence was highest between ages 25-29 and 35-39, at just over 70%. Prevalence was by far the lowest at ages 15-19 (47%). In the youngest cohort, most modern contraceptors were using condoms. At the older ages, IUDs, withdrawal, and periodic abstinence were the most commonly used methods. Overall contraceptive prevalence was only 34% among women with no living children, another indication that most couples still want to have a child soon after marriage (Figure 7.2 and Table 7.6). Prevalence peaked at 75% among those with two living children, before falling back to 63% for those with three or more living children. Unlike most other contraceptive methods, use of condoms and oral contraceptives fell steadily as the number of children increased. Use of the IUD was rare among women with no children. As expected, contraceptive sterilization was very rare except among women with at least two children. Even among that group only 3% had been sterilized.

The greatest differences in contraceptive use between urban and rural areas were not so much in prevalence (69% vs. 63%, respectively), but in the types of methods selected (Table 7.7). In urban areas, 42% of couples were using a modern method, while 27% were using a traditional method.. In rural areas, however, this was almost reversed; use of traditional methods exceeded modern methods, 36% to 27%. Condom and OC use was much less prevalent in rural areas than in urban areas, while the reverse was true for withdrawal. Table 7.7 also shows prevalence figures for the Donetsk and Odessa Oblasts, the two oversampled oblasts in the URHS. Contraceptive prevalence there was 65% and 72%, respectively. Most noteworthy, however, is that women in both oblasts relied more heavily on modern methods (45% and 47%) than other Ukrainian women, even those in urban areas. Table 7.8 reveals that, despite other reproductive differences that have been noted between Ukraine's regions, there is almost no variation in contraceptive prevalence by region. There

is however, a correlation with the types of method employed. The western region relies much more heavily than others on traditional contraception (42% traditional and 25% modern). In all other regions the figures were more or less reversed. Most of the traditional use in the West is withdrawal. The Central and East regions are areas of particularly high IUD use.

Contraceptive use among women in union was also strongly correlated with educational attainment (Table 7.9). Among the relatively small number of women who had not completed secondary school, only about 57% were using contraception, less than the prevalence among better educated women (66% for those who completed secondary school and 76% for those who received any post-secondary education. The prevalence of condom, oral contraceptive, and periodic abstinence use, showed great proportional increase with educational attainment. Reliance on withdrawal decreased somewhat with increasing education.

Table 7.10 shows current contraceptive use according to union status. Thirty-five percent of women previously in union and 22% of those never in union were using contraception. The majority of users never in union, were relying on condoms.

Recent trends in contraceptive prevalence

Table 7.11 and Figure 7.3 show the trend in contraceptive prevalence in Ukraine from the beginning of 1994 to the date of the survey, based on data from the URHS contraceptive calendar. In the calendar, women reported, to the best of their recollection, their month-by-month contraceptive history for the previous five years. Because the survey questionnaire included no marriage or sexual activity history, these calculations include all interviewed women, rather than just women in union or sexually active women. Overall prevalence looks to have risen steadily among 15 to 39 year-old women in Ukraine during that time period. In just over five years prevalence rose by about 7 percentage points, more than 1 percentage point per year. Recent trends in contraceptive prevalence have been similar in urban and rural areas. Very notably, during this time there were greater increases in the use of modern contraception (5 percentage points) than in the use of traditional methods (2 percentage points).

Source of contraceptive methods

Percentage distributions of sources of oral contraceptives (OCs), IUDs and condoms (the most widely used supplied methods) are displayed in Table 7.12. Women's consultation centers and pharmacies were overwhelmingly the leading sources of OCs for respondents, supplying about three of every four users. Women's consultation centers were the predominant source of IUDs, accounting for two-thirds of those currently used. Most of the remainder were supplied by hospitals. Pharmacies supplied slightly over half of condom users. Substantial numbers were also provided by women consultation centers and drug kiosks. Major sources of supply for these methods were quite similar for urban and rural women.

Information collected in the survey about the cost of contraceptive methods proved to be problematic because of women's difficulty in remembering the amount they paid as well as substantial changes in the costs of goods and services and in the value of Ukrainian currency. However, it is informative to examine the types of payments women or couples made for their current contraceptive method. Table 7.13 shows that among OC users, 82% paid for their pills. A slightly higher percentage of

IUD users received their supplies without paying—about one-fourth said they did not pay. About three-fourths of condom users said they paid for their supplies, but an additional 11% did not know (probably because their partner obtained their condoms). There were very small differences between urban and rural areas with regard to whether respondents paid for their contraceptive method.

Reasons for not using contraception

Among survey respondents who were not currently using contraception, two-thirds cited lack of sexual activity, an inability or difficulty in becoming pregnant (subfecundity), current pregnancy, or a desire to become pregnant, as their primary reason for non-use (Table 7.14). These women are not likely to be targeted as potential users of contraception in a family planning program. There was a broad assortment of reasons for non-use given by the remaining 34% of non-users, many of which could be addressed by reproductive health interventions. The most commonly given reasons were “occasional sex only” (9%) and “don’t know” (8%). Fear of health consequences or side effects (3%), partner objections (3%), and cost/access issues (3%), were the next most commonly cited reasons. It is important to note that certain reasons, particularly a preference for abortion and religion, were almost never mentioned. Not surprisingly, there were large differences in reasons for non-use according to marital/union status. Among those not currently in union, lack of sexual activity and infrequent sexual activity were, of course, the most important factors. Among those currently in union, factors such as subfecundity, pregnancy, wanting to become pregnant, and fear of side effects were more often cited than among other respondents.

Unmet need for contraception

Table 7.15 presents estimates of the percentage of women in need of family planning services according to two definitions. By the first definition (the conventional definition), women who are sexually active, not pregnant, able to become pregnant, do not want to become pregnant, and are not using any contraceptive method are considered to have unmet need for contraception. By this definition, unmet need was 15%, very high compared with the levels in most other developed countries. The second definition additionally includes users of periodic abstinence and withdrawal (methods with typically low use-effectiveness) as having unmet need. This definition more than doubles the proportion with unmet need to 37%. This is an extremely high level of unmet need, fitting well with Ukraine’s high abortion rate. Women with no living children were substantially less likely to be in need than those with children by both definitions I and II. Respondents who had completed secondary school but had no post-secondary education were more likely to have unmet need than others were. There were virtually no differences across regions of the country. Rural women were more likely to have unmet need than urban women. It should be kept in mind, however, that these indicators only take into account whether people are using a method, but do not include such factors as consistency of use and method effectiveness.

Preference for other methods/Problems with current method

Current users of contraception were asked if they would prefer to use a method of pregnancy prevention other than the one they were currently using. A relatively low 25% of women said they preferred a different method, but the percentages varied considerably according to the method currently used (Table 7.16). The two methods that women were the most likely to want to switch from were withdrawal (37%) and condoms (30%), both male controlled methods with relatively low

use-effectiveness. Users of the IUD and tubal ligation were the most likely to be satisfied with their present method, with only 10% and 11%, respectively, preferring a different method. The IUD was also by far the method most commonly mentioned as being the one women preferred to use, among women using a different method (mentioned by about 40%). Oral contraceptives were the only other method frequently mentioned by respondents as one they would like to adopt.

Table 7.17 shows that about one-half of women who reported that they preferred to be using a different method, said that the major reason for their failure to use that method was either cost (27%) or fear of health consequences or side effects associated with the method (20%). The only other reasons frequently cited were that the physician would not prescribe their preferred method (15%) and that she did not know enough about the method or how to obtain it (14%). The most notable difference between urban and rural areas was that cost was much more likely to be a major factor in rural areas than in urban areas. It is noteworthy that all four of the most common reasons given for not using preferred methods all could be addressed by interventions designed to improve contraceptive use.

Overall, about 80% of contraceptive users said they were having no major problems or concerns with their current method (Table 7.18). The proportion with no concerns ranged from 73% among those using a traditional method to 84% for condom users, 85% for IUD users, and 87% for sterilized women. The problem/concern most often mentioned, especially for traditional methods and condoms, was low effectiveness of the method. Among women using an IUD or who had been sterilized, health concerns were the leading concern. Among users of OCs, side effects were the predominant concern.

Contraceptive failure and discontinuation

Data from the questionnaire's contraceptive/pregnancy calendar were used to calculate rates of contraceptive failure (the probability of becoming pregnant while using a particular method) and discontinuation (the probability of stopping use of a particular method for any reason) for the most widely used methods. Table 7.19 presents rates of failure after one, two, and three years for all methods combined and for five specific methods. Overall about 9% of contraceptive users became pregnant while using a method within one year of beginning use. After three years this rose to 19%. Of course, there were substantial differences between methods. There were quite small differences in failure rates between urban and rural areas.

The failure rate for the IUD, based on calendar data, was 1.4% for the first year, which is in line with typical rates (Figure 7.4). Failure rates after three years rose to 3.5%. Urban-rural differences were small. The failure rate for oral contraceptives was 5.9% for the first year and 13.2% for three years, somewhat higher than the rates typically seen (Hatcher et al., 1998). Condom failure was 7.1% for one year and 18.7% for three years. The highest failure rates for widely used methods were found for periodic abstinence and withdrawal. For periodic abstinence 15.6% of users became pregnant in the first year; for withdrawal the rate was 11.7%. Three-year rates for these two methods was almost 30%. There were no consistent differences between urban and rural areas in contraceptive failure or use-effectiveness, as the inverse is known.

Anecdotal reports have suggested that there is a considerable amount of method switching, as well as frequent starting and stopping of contraception in the much of the former Soviet Union. The

URHS data support this belief. In general, contraceptive discontinuation rates were very high across methods. For all methods combined, 29% of contraceptive use segments continued for no more than one year (Table 7.20). After three years, about half of women had discontinued use. Of the five methods most widely used, all except the IUD exhibited extremely high rates of discontinuation, from 29% (withdrawal) to 54% (OCs) in the first year and from 53% (withdrawal) to 74% (OCs) after three years (Figure 7.4). Only 6% of IUD segments ended within one year.

Table 7.21 displays some reason-specific one year discontinuation rates for the five leading methods of contraception in Ukraine. For oral contraceptive users there were a wide variety of reasons for discontinuation, including: to give the body a rest (15% of segments terminating in less than one year), side effects (9%), health concerns (8%), cost/supply problems (7%), and physician's decision (7%). Especially of concern is the high rate of discontinuation to "give the body a rest", which has no medical justification. Side effects were somewhat of a problem for IUD users, causing 10% discontinuation in the first year. Concerns about health were responsible for 2% discontinuation. Condoms tended to be discontinued for a more effective method (7%), because of cost (4%), or inconvenience of the method (3%). The major reason for discontinuing periodic abstinence and withdrawal was to use a more reliable method.

Table 7.22 shows percentage distributions of reasons for discontinuing the five most widely used methods. It should be kept in mind that these distributions are not the same as discontinuation rates and that percentages should only be compared within methods, not between methods. Oral contraceptive users cited a wide assortment of reasons for stopping. The most commonly given reasons were "giving her body a rest", side effects, health concerns, pregnancy (i.e., failure), and desire to get pregnant. Physician's decision and cost/supply issues were also mentioned relatively often. Among IUD users the principal reasons for discontinuation were to give the body a rest, physician's decision/recommendation, health concerns, and pregnancy while using. Condom use was discontinued for a broad assortment of reasons, led by pregnancy and want of a more reliable method. Periodic abstinence and withdrawal use tended to be terminated because of pregnancy (over 40%), desire for a more reliable method, or inconvenience of the method. In the case of withdrawal, partner's objections were commonly cited.

Opinions about fertility control methods

Respondents were asked to rate a number of birth prevention methods with regard to safety and health effects, effectiveness, and cost, as well as to give each method an overall rating. For each characteristic, women rated each method between 1 (extremely negative) and 10 (extremely positive). Table 7.23 and Figure 7.5 show the percentages of women who gave very low ratings (3 or less) for each of seven methods. (Not included in these tabulations are women who did not have an opinion about particular characteristics for a given method. For some methods, particularly injectables and tubal ligation, the proportions of women with no opinion were very high.)

Survey findings regarding feelings about abortion run counter to the opinion sometimes expressed that women in this region of the world tend to prefer abortion to contraception. Probably the most noteworthy result is the nearly universally negative overall opinions held by respondents about both conventional induced abortion (96%) and miniabortion (95%). Opinions about abortion were equally negative regardless of whether women had ever had an abortion (top panel, Table 7.23). Despite the fact that abortion was by far the most poorly regarded method of birth prevention, every

method asked about was rated negatively overall by at least 40% of respondents who had an opinion about the method. Condoms and the IUD were the only methods not viewed very negatively by a majority of respondents, 41% and 49% negative, respectively. After conventional abortion and miniabortion, female sterilization (81%) and injectables (73%), methods that were not widely used in Ukraine, were the methods most often viewed negatively. Except for abortion, users of a particular method were far less likely than non-users to view that method negatively. IUD users (3%) and oral contraceptive users (8%) were particularly unlikely to view their method negatively.

With regard to safety and health consequences, induced abortion was again viewed the most negatively (91% for conventional abortion and 88% for miniabortion). Slightly over one-half of respondents considered female sterilization (57%) and injectables (53%) to be unsafe. Condoms (9% negative) were considered the safest method by a wide margin, followed by the IUD (27%). As with overall opinions, users of a particular method were less likely than others to consider that method unsafe.

None of the methods about which women were asked in the survey were widely considered to be of low effectiveness. However, even the methods known to be of extremely high reliability, such as sterilization and the IUD were considered by at least 7% of respondents to be of low effectiveness. Interestingly, sterilized women were no less likely than others to believe that sterilization was of low effectiveness.

Only condoms were viewed by more than a small proportion of women (58%) as very costly. Because condoms play such an important role in preventing the spread of HIV/AIDS and other STIs, the fact that so much of the population views them as very costly is essential information for policy makers and the reproductive health community. The IUD was viewed as expensive by 15% of women. Cost was the only aspect of contraception which was generally viewed more negatively by users than by non-users. This probably reflects greater awareness of the actual costs of obtaining methods among those women who have actually used particular methods.

Use of non-supplied methods

Because non-supplied methods of contraception tend to have higher probabilities of failure than modern, supplied methods, the URHS examined the reasons that women and couples chose methods such as periodic abstinence and withdrawal. The survey asked every respondent who was currently using any non-supplied method (mainly periodic abstinence or withdrawal) whether a number of factors were at least “somewhat important” in their method selection. These factors included: health/side effects of supplied methods; the naturalness of the method; partner preference; lack of knowledge/ information about other methods; cost of other methods; difficulty in obtaining other methods; and religious beliefs. All but religion were cited by a substantial proportion (i.e., at least 47% or more) of users of non-supplied methods as being at least “somewhat important” in influencing their method choice (Table 7.24). The possible health and side effects of supplied methods (81%) and the naturalness of non-supplied methods (82%) were by far the most important factors cited in choosing withdrawal and periodic abstinence. About two of every three women said that the cost of other methods played an important role in method selection, about the same as the proportion mentioning their partner’s preferences. Several factors that family planning/reproductive health programs could affect seem to play a significant role in the decision-making process. In addition to the cost of other methods, difficulty in obtaining other methods and a lack of knowledge

about other methods influenced the choices of 52% and 47%, respectively, of users of less effective, non-supplied methods. Rural women were slightly more likely to cite the cost and availability of other methods as a reason.

Table 7.25 reveals non-supplied method users' opinions about the effectiveness of their current method relative to "methods received from a doctor or pharmacy, like the IUD or pills". Only about one of every five such women were aware that methods such as the IUD prevented pregnancy more effectively than the method they were using. Another one-third felt that their current method was actually more effective than modern methods. Twelve percent admitted that they did not know the relative effectiveness of the methods.

By publicizing the relative effectiveness of various types of contraception, disseminating accurate information on health effects, and improving knowledge of and access to other methods, reproductive health programs in Ukraine can contribute to increased use of highly effective methods and, hence, reduced reliance on induced abortion to prevent unwanted births.

IUD use

The IUD has been the most popular form of modern contraception in much of eastern Europe and the former Soviet Union for many years. However, little information has been collected on a national basis in Ukraine on side effects or other aspects of IUD provision and use. The URHS included a series of questions for respondents who had an IUD inserted since the beginning of 1994, relating to the timing of insertions, information given by the IUD provider, and problems encountered related to the IUD. Tables 7.26 and 7.27 provide some findings from these questions.

Most IUD insertions (71%) took place neither following a delivery nor an abortion (top panel, Table 7.26). There were, however, substantial numbers of insertions after induced abortions (23%). Eighty-five percent of recent Ukrainian IUD users reported that their provider told them how long their IUD could be left in place (Table 7.26, bottom panel). About one in ten IUD users were told that the device could be left in place for six or more years, about half reported being told that it could be left in place for 4-5 years, and 23% said they were told that the IUD could only be left in place for less than four years. Unfortunately, because we do not have information on the type of IUD that women used (this question was not asked because women were unlikely to know the type) we cannot assess the adequacy of these recommendations, since there are considerable differences between different types of IUDs as to how long they remain effective. Six percent of IUD users reported that their provider did not give them any information regarding how long the device could be left in place and another 9% could not remember whether their physician told them anything about how long her IUD should remain.

About three of every ten women with an IUD inserted since January 1994 reported that they experienced physical problems associated with the device (Table 7.27). The proportion was very similar in urban and rural areas of Ukraine. About half of those reporting problems mentioned either heavy bleeding during menstrual periods or bleeding or spotting between periods. Inflammation/discharge/infection was the only other specific problem frequently mentioned.

Oral contraceptive use

The questionnaire included a module on oral contraceptive use for respondents who reported any segments of OC use beginning since January 1994. This module was similar to the one used for IUDs, including questions on information given by providers, problems related to pill use, and related topics. Although oral contraceptives are not widely used in Ukraine currently, they seem to be growing in popularity and are being promoted as a safe and effective form of pregnancy prevention that is appropriate for many women in the population.

Despite the relatively small number of current OC users interviewed (210), there were a considerable number of brands of OCs being used, with 23 different brands mentioned (data not shown). The most widely used brands among current users were Tri-Regol (26%), Regividon (22%), Marvelon (19%), and Triqvilar (17%). Twelve percent of OC users reported that they were using Postinor, a very high dose pill also used as a morning-after pill.

Slightly fewer than one-half (43%) of recent OC users in the URHS reported that their physician did not tell them how long they could continue to take OCs (Table 7.28). Unlike the IUD, however, this is not of great concern, since most women can take OCs for many years with no ill effects. Fourteen percent reported that their physician told them they should only take pills for less than two years. As was shown previously, many physicians continue to tell patients that they should discontinue OCs after various lengths of time to “give the body a rest”, a recommendation that is not supported by medical evidence. About one in four women were told they could take OCs as long as they wanted. Rural women were less likely to be told this and more likely to be told they should take OCs only for a short period of time.

Forty-one percent of recent and current OC users reported having had physical problems related to their use of this contraceptive method (Table 7.29). Nausea and weight gain/bloating were the most commonly reported problems, each reported by 8% of users. Additionally, 6% reported headaches and 5% cited bleeding between menstrual periods.

Contraceptive sterilization

Despite the fact that most women want to have no more children and that most couples desire no more children long before reaching their potentially fertile years end, contraceptive sterilization is rarely employed as a method of pregnancy prevention in Ukraine. The URHS found that only 2% of married women of reproductive age with two or more children had been sterilized and that virtually no men had undergone a vasectomy. This low reliance on sterilization seems to be a phenomenon observable throughout the region. Recent surveys in Russia, the Czech Republic, and Romania, for instance, also revealed similarly low prevalence of sterilization (Czech Statistical Office et al. 1995, Romanian Ministry of Health 1995, VCIOM 1998), as does anecdotal information from much of eastern Europe and the former Soviet Union. A number of factors probably contribute to the low prevalence of sterilization, one of which is legislative. Except for medical reasons, it is only legally permissible for women with three or more children or, if a woman is over 30 years of age, with two children. This law, however, actually represents a liberalization of sterilization practices. Until after the break-up of the Soviet Union, only women with a medical condition contraindicating pregnancy or childbirth could legally be sterilized.

Regardless of the regulations on when sterilizations are permitted, the survey results show that few women claim to be interested in tubal ligation. Overall, among fecund respondents who wanted to have no more children, only 7% expressed interest in sterilization (Table 7.30). An additional 5% said they were not sure if they would be interested, leaving an overwhelming 88% who were not interested. There were a few noteworthy differentials in sterilization interest according to women's characteristics. Respondents in the West were the least likely to be interested. Women with the largest families and those with the highest levels of education, were more likely to express interest than others. Non-users of contraception were less likely than users to be interested, but there was little difference between users of various methods of contraception.

When asked the reason for not being interested in sterilization, the factor most commonly given (respondents could only give one reason) was that women simply "had not thought about it", cited by 34% of respondents (Table 7.31). This indicates that sterilization is not even a contraceptive option that many women ever consider. The next most common reason was a fear of health risks associated with sterilization (25%). Other reasons that were given with some frequency were lack of information about sterilization (10%), fear of operation (9%), and might eventually want another child (8%). Women rarely mentioned religion, cost of the procedure, or partner's objections as major factors.

Although, at first appearance, increasing the incidence of contraceptive sterilization seems an obvious way to decrease levels of unintended pregnancy and induced abortion and, hence, reduce maternal morbidity and mortality, there are a number of major obstacles to overcome before such a change can actually occur. Legal barriers would need to be removed. There is currently not adequate equipment available to provide sterilization to large numbers of women (or men). Maybe, most importantly, the procedures are not currently ones that are even considered to be acceptable contraceptive options for all but a small minority of Ukrainians. Any movement directed at trying to increase the use of sterilization as a family planning method will have to address all of these problems.

Figure 7.1
Percentage Distributions of Current Contraceptive Method
Among Women in Union
1999 Ukraine Reproductive Health Survey




Figure 7.2
Percentage Distributions of Current Contraceptive Use Among
Women in Union, by Number of Living Children
1999 Ukraine Reproductive Health Survey




Figure 7.3
Percentage Point Change Since January 1994 in Percent of
All 15-39 Year-Old Women Currently Using Contraception
1999 Ukraine Reproductive Health Survey




Figure 7.4
One-Year Failure and Discontinuation Rates
for Selected Contraceptive Methods
1999 Ukraine Reproductive Health Survey





Figure 7.5
Percent of Respondents Giving Low Ratings Overall and
for Safety/Health for Selected Contraceptive Methods
1999 Ukraine Reproductive Health Survey





Table 7.1
 Percentage of respondents who heard of specific contraceptive methods,
 by residence and marital status
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Total	Residence		Marital status		
		Urban	Rural	Currently in union	Previously in union	Never in union
Condoms	98.7	98.9	97.9	98.7	98.9	98.3
IUD	95.9	96.2	95.0	98.5	98.3	87.1
Oral contraceptives	90.1	91.9	85.4	91.1	92.8	86.1
Female sterilization	67.2	70.5	58.2	71.4	76.9	50.1
Spermicide	60.2	64.4	49.1	62.2	65.4	52.1
Diaphragm	57.8	62.5	45.3	61.1	66.3	44.2
Injectables	48.0	50.6	41.2	50.5	52.1	38.9
Vasectomy	45.4	49.2	35.4	48.0	52.8	34.6
Implants	18.0	20.5	11.5	18.5	21.0	15.3
Withdrawal	88.2	88.7	86.9	93.6	92.2	70.5
Periodic abstinence	81.9	84.1	76.0	86.3	86.7	66.8
Any method	99.6	99.7	99.6	99.9	99.8	99.8
Any modern method	99.6	99.6	99.5	99.8	99.8	98.8
Any traditional method	92.3	92.9	90.7	96.3	95.3	79.3
<i>Number of women</i>	<i>7128</i>	<i>5544</i>	<i>1584</i>	<i>4796</i>	<i>800</i>	<i>1532</i>

Table 7.2
 Percentage of respondents who heard of specific contraceptive methods, by age and education
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Total	Age						Education		
		15-19	20-24	25-29	30-34	35-39	40-44	<Comp. Sec.	Comp. Sec.	>Comp. Sec.
Condoms	98.7	98.2	99.0	98.7	99.5	98.4	98.1	97.3	98.6	99.7
IUD	95.9	83.5	97.7	98.5	98.6	98.7	98.1	83.5	97.0	98.7
Oral contraceptives	90.1	82.2	89.8	92.4	92.4	93.5	90.5	80.2	90.1	95.8
Female sterilization	67.2	39.6	62.9	73.6	76.4	78.8	71.4	42.1	66.2	84.3
Spermicide	60.2	46.1	58.3	62.8	66.3	66.2	61.4	38.9	58.8	76.8
Diaphragm	57.8	35.0	52.7	60.5	65.2	68.6	64.4	29.3	56.8	77.0
Injectables	48.0	33.0	45.4	50.4	56.8	53.2	49.0	28.8	47.2	61.4
Vasectomy	45.4	23.9	44.0	50.7	52.1	53.4	48.3	21.8	42.4	68.8
Implants	18.0	10.3	17.6	17.4	22.5	20.5	19.9	8.6	16.5	28.3
Withdrawal	88.2	65.5	88.8	94.2	95.0	94.0	91.6	68.9	89.5	94.6
Periodic abstinence	81.9	61.3	80.4	85.7	88.1	88.3	87.3	60.6	82.2	92.4
Any method	99.6	98.7	99.8	99.9	99.8	99.9	99.7	98.6	99.7	100.0
Any modern method	99.6	98.7	99.8	99.8	99.8	99.9	99.5	98.3	99.7	100.0
Any traditional method	92.3	76.2	93.2	96.2	97.5	96.0	95.0	77.9	93.4	96.7
<i>Number of women</i>	<i>7128</i>	<i>1100</i>	<i>1182</i>	<i>1227</i>	<i>1195</i>	<i>1246</i>	<i>1178</i>	<i>778</i>	<i>4828</i>	<i>1522</i>

Table 7.3
 Percentage of respondents who know where to obtain specific contraceptive methods, by residence, marital status, and education
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Total	Residence		Marital status			Education		
		Urban	Rural	Currently in union	Previously in union	Never in union	<Comp. Sec.	Comp. Sec.	>Comp. Sec.
Condoms	96.9	97.3	95.7	97.3	97.6	95.3	94.4	96.8	98.3
IUD	90.7	91.3	89.1	95.1	95.6	75.6	72.4	92.3	95.5
Oral contraceptives	85.1	87.2	79.7	86.8	89.4	78.4	71.8	85.2	92.3
Spermicide	55.5	59.5	44.8	57.4	61.4	47.1	33.9	54.2	71.9
Diaphragm	51.4	55.3	41.2	54.8	59.5	37.6	24.0	50.5	69.8
Female sterilization	49.8	51.8	44.3	53.3	58.8	35.2	28.7	49.4	62.9
Injectables	40.2	42.0	35.6	42.5	45.0	31.5	24.0	39.6	51.3
Vasectomy	31.9	34.1	26.3	33.8	38.6	23.2	14.1	30.5	49.7
Implants	13.0	14.6	8.9	13.5	15.9	10.5	5.1	12.2	20.2
Periodic abstinence*	74.3	76.7	67.9	79.3	80.0	56.9	49.5	74.5	87.2
<i>Number of women</i>	<i>7128</i>	<i>5544</i>	<i>1584</i>	<i>4796</i>	<i>800</i>	<i>1532</i>	<i>778</i>	<i>4828</i>	<i>1532</i>

*Knows where to get information about using periodic abstinence

Table 7.4
 Percentage of respondents who have ever used specific contraceptive methods, by residence, marital status, and education
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Total	Residence		Marital status			Education		
		Urban	Rural	Curr. in union	Prev. in union	Never in union	<Comp. Sec.	Comp. Sec.	>Comp. Sec.
Condoms	44.1	49.1	30.7	49.2	51.8	25.6	24.2	43.0	58.9
IUD	29.7	30.7	27.0	38.0	39.1	1.1	11.7	31.8	32.7
Oral contraceptives	13.8	15.8	8.5	16.0	16.5	6.0	5.2	13.4	20.0
Spermicide	4.2	5.1	2.1	5.1	4.6	1.5	0.8	3.7	7.8
Injectables	1.2	1.4	0.8	1.5	1.2	0.4	0.8	1.2	1.6
Female sterilization	1.0	1.0	1.0	1.4	0.7	0.0	1.0	1.1	0.8
Diaphragm	0.6	0.8	0.3	0.7	1.4	0.1	0.2	0.6	1.1
Vasectomy	0.1	0.1	0.1	0.1	0.3	0.0	0.0	0.1	0.0
Implants	0.1	0.2	0.1	0.2	0.0	0.1	0.0	0.1	0.2
Withdrawal	47.1	46.6	48.4	55.9	55.8	17.2	24.7	48.3	55.4
Periodic abstinence	37.6	39.7	31.8	45.1	45.0	12.0	14.3	37.1	52.2
Any method	74.2	76.0	69.5	87.2	83.9	31.6	40.8	76.4	85.5
Any modern method	59.8	64.3	48.0	69.4	70.4	27.0	32.0	60.5	73.0
Any traditional method	57.3	57.9	55.8	68.6	67.3	19.9	28.4	59.0	68.0
<i>Number of women</i>	<i>7,128</i>	<i>5,544</i>	<i>1,584</i>	<i>4,796</i>	<i>800</i>	<i>1,532</i>	<i>778</i>	<i>4,828</i>	<i>1,532</i>

Table 7.5
 Percentage distribution of current contraceptive use, by method, by respondent's age, women in union
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Age					
		15-19	20-24	25-29	30-34	35-39	40-44
Using any method	67.5	47.4	63.0	71.6	71.7	70.9	61.1
Using a modern method	37.6	27.2	33.9	42.0	41.9	40.1	30.2
IUD	18.6	3.4	12.9	19.3	21.5	23.1	16.6
Condoms	13.5	20.8	15.8	16.0	15.0	10.7	9.6
Oral contraceptives	3.0	1.2	4.3	3.3	3.0	2.9	1.9
Female sterilization	1.4	0.0	0.1	1.1	1.6	2.3	1.7
Spermicide	0.8	1.9	0.8	1.6	0.4	0.7	0.4
Other methods	0.3	0.0	0.1	0.7	0.3	0.4	0.1
Using a traditional method	29.9	20.2	29.2	29.7	29.9	30.8	30.9
Withdrawal	19.5	13.8	24.2	20.6	19.9	17.0	17.8
Periodic abstinence	10.4	6.4	4.9	9.1	10.0	13.9	13.1
Using no method*	32.5	52.6	37.0	28.4	28.3	29.1	38.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>4,794**</i>	<i>121</i>	<i>737</i>	<i>993</i>	<i>1,009</i>	<i>1,000</i>	<i>934</i>

*Includes users of douche and folk methods

**Data missing for 2 women.

Table 7.6
 Percentage distribution of current contraceptive use, by method, by number of living children,
 women in union
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Living children			
		0	1	2	3+
Using any method	67.5	33.7	68.9	74.8	62.8
Using a modern method	37.6	21.2	41.4	39.5	27.6
IUD	18.6	2.6	18.4	22.8	17.5
Condoms	13.5	14.6	17.2	10.8	7.1
Oral contraceptives	3.0	2.5	3.7	2.8	0.6
Female sterilization	1.4	0.5	0.4	2.5	1.8
Spermicide	0.8	1.0	1.2	0.5	0.4
Other methods	0.3	0.0	0.5	0.2	0.2
Using a traditional method	29.9	12.5	27.5	35.3	35.3
Withdrawal	19.5	9.3	18.0	22.5	24.6
Periodic abstinence	10.4	3.2	9.5	12.8	10.7
Using no method*	32.5	66.3	31.1	25.2	37.2
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>4794</i>	<i>448</i>	<i>2009</i>	<i>1906</i>	<i>322</i>

*Includes users of douche and folk methods

Table 7.7
 Percentage distribution of current contraceptive use, by method, by type of place of residence
 and for oversampled oblasts, women in union
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Place of residence		Oversampled oblasts	
		Urban	Rural	Donetsk	Odessa
Using any method	67.5	69.2	63.3	65.4	71.7
Using a modern method	37.6	41.9	27.2	44.8	46.5
IUD	18.6	19.0	17.8	20.5	15.4
Condoms	13.5	16.5	6.4	17.8	22.5
Oral contraceptives	3.0	3.6	1.4	4.4	4.6
Female sterilization	1.4	1.4	1.4	1.1	1.4
Spermicide	0.8	1.0	0.3	0.7	1.9
Other methods	0.3	0.4	0.0	0.3	0.7
Using a traditional method	29.9	27.3	36.1	20.6	25.2
Withdrawal	19.5	16.2	27.4	10.1	15.5
Periodic abstinence	10.4	11.1	8.7	10.5	9.7
Using no method*	32.5	30.8	36.7	34.6	28.3
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>4,794</i>	<i>3,641</i>	<i>1,153</i>	<i>1,038</i>	<i>842</i>

*Includes users of douche and folk methods

Table 7.8
 Percentage distribution of current contraceptive use, by method, by region of residence, women in union
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Region				
		North	Central	East	South	West
Using any method	67.5	68.3	68.1	66.3	68.9	67.1
Using a modern method	37.6	37.8	37.7	43.1	42.9	24.8
IUD	18.6	15.9	25.2	21.9	17.1	12.7
Condoms	13.5	16.1	8.9	15.9	16.4	8.2
Oral contraceptives	3.0	3.1	2.3	3.0	4.8	1.8
Female sterilization	1.4	1.4	1.4	1.3	1.2	1.6
Spermicide	0.8	0.9	0.0	0.6	2.1	0.5
Other methods	0.3	0.4	0.0	0.4	0.5	0.1
Using a traditional method	29.9	30.5	30.4	23.2	26.0	42.3
Withdrawal	19.5	21.5	20.1	12.7	14.9	31.7
Periodic abstinence	10.4	9.0	10.3	10.5	11.1	10.6
Using no method*	32.5	31.7	31.9	33.7	31.1	32.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>4,794</i>	<i>566</i>	<i>426</i>	<i>1,777</i>	<i>1,194</i>	<i>831</i>

*Includes users of douche and folk methods

Table 7.9
 Percentage distribution of current contraceptive use, by method, by educational level, women in union
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Educational level		
		<Complete secondary	Complete secondary	>Complete secondary
Using any method	67.5	56.5	65.9	75.9
Using a modern method	37.6	28.6	35.7	46.6
IUD	18.6	15.2	19.1	17.9
Condoms	13.5	8.0	11.6	21.6
Oral contraceptives	3.0	2.1	2.7	4.3
Female sterilization	1.4	2.3	1.4	1.1
Spermicide	0.8	0.7	0.6	1.6
Other methods	0.3	0.5	0.3	0.2
Using a traditional method	29.9	27.9	30.2	29.3
Withdrawal	19.5	23.6	20.1	16.1
Periodic abstinence	10.4	4.3	10.1	13.2
Using no method*	32.5	43.5	34.1	24.1
Total	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>4,794</i>	<i>290</i>	<i>3,456</i>	<i>1,048</i>

*Includes users of douche and folk methods

Table 7.10
 Percentage distribution of current contraceptive use, by method, by marital status
 1999 Ukraine Reproductive Health Survey

Current contraceptive method	Total	Marital status		
		Currently in union	Previously in union	Never in union
Using any method	53.5	67.5	34.9	21.7
Using a modern method	31.2	37.6	22.9	16.6
IUD	13.8	18.6	12.4	0.5
Condoms	12.9	13.5	7.5	13.8
Oral contraceptives	2.5	3.0	1.6	1.6
Female sterilization	1.0	1.4	0.7	0.0
Spermicide	0.7	0.8	0.3	0.4
Other methods	0.3	0.3	0.4	0.2
Using a traditional method	22.3	29.9	12.0	5.1
Withdrawal	14.5	19.5	6.6	3.7
Periodic abstinence	7.8	10.4	5.4	1.4
Using no method*	46.5	32.5	65.1	78.3
Total	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	7,128	4,794	800	1,532

*Includes users of douche and folk methods

Table 7.11
 Percent of All 15-39 Year-Old Respondents Reporting Current Use of Any Contraception
 or of Modern Contraception at Six Month Intervals from January 1994 to April 1999
 1999 Ukraine Reproductive Health Survey

Date	Total			Urban			Rural		
	% Using Any Method	% Using Modern Method	% Using Traditional Method	% Using Any Method	% Using Modern Method	% Using Traditional Method	% Using Any Method	% Using Modern Method	% Using Traditional Method
1/1994	48.8	27.7	21.1	49.5	30.4	19.1	46.9	20.9	26.0
7/1994	50.1	29.1	21.0	51.2	32.1	19.1	47.0	21.3	25.7
1/1995	50.7	29.4	21.3	51.6	32.0	19.6	48.3	22.7	25.6
7/1995	51.1	30.0	21.1	52.5	32.9	19.6	47.7	22.3	25.4
1/1996	51.6	30.1	21.5	52.6	33.0	19.6	49.0	22.5	26.5
7/1996	51.7	30.5	21.2	52.9	33.4	19.5	48.8	22.9	25.9
1/1997	52.4	30.6	21.8	53.1	33.4	19.7	50.5	23.5	27.0
7/1997	53.1	31.3	21.8	53.8	34.1	19.7	51.0	23.9	27.1
1/1998	53.6	31.6	22.0	54.7	34.9	19.8	50.6	22.9	27.7
7/1998	53.9	32.0	21.9	55.4	35.4	20.0	50.1	23.1	27.0
1/1999	54.2	32.1	22.1	55.4	35.6	19.8	51.1	23.1	28.0
4/1999	55.9	32.9	23.0	56.9	36.3	20.6	52.4	23.9	28.5

Table 7.12
 Percentage distribution of source of selected contraceptive methods, by residence, among current users
 1999 Ukraine Reproductive Health Survey

Source of method	Contraceptive method								
	Total			Urban			Rural		
	OCs	IUD	Condoms	OCs	IUD	Condoms	OCs	IUD	Condoms
Women's consultation centers	39.3	66.0	10.0	41.1	69.9	9.6	*	55.8	12.2
Pharmacy	34.6	3.7	56.6	34.8	2.8	56.2	*	6.0	59.1
Hospital	6.6	20.4	2.1	5.9	17.8	2.3	*	27.1	0.5
Drug kiosk	1.7	0.1	7.3	1.2	0.0	7.1	*	0.5	9.1
Maternity house	2.1	5.3	0.5	2.5	5.0	0.6	*	6.0	0.0
Private clinic/physician	1.0	0.4	0.3	1.2	0.3	0.3	*	0.9	0.0
MCH center	0.4	0.1	0.4	0.5	0.2	0.4	*	0.0	0.0
Other source	15.7	4.4	22.8	12.8	4.0	23.5	*	3.7	30.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>198</i>	<i>1,005</i>	<i>973</i>	<i>179</i>	<i>780</i>	<i>867</i>	<i>19</i>	<i>225</i>	<i>106</i>

*Fewer than 25 respondents

Percentage distribution of method of payment for contraceptive method for current users of oral contraceptives, IUD, and condoms
1999 Ukraine Reproductive Health Survey

Type of payment	Contraceptive method								
	Total			Urban			Rural		
	OCs	IUD	Condoms	OCs	IUD	Condoms	OCs	IUD	Condoms
Cash	82.3	72.4	76.0	80.8	72.3	75.9	*	72.6	76.8
Payment in goods or services	0.2	1.4	0.3	0.2	1.4	0.4	*	1.4	0.0
No payment	16.8	26.1	10.8	18.2	26.1	10.5	*	26.0	12.6
Don't remember/Don't know	0.7	0.1	12.7	0.8	0.2	13.2	*	0.0	10.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>198</i>	<i>1,005</i>	<i>1,016</i>	<i>179</i>	<i>780</i>	<i>906</i>	<i>19</i>	<i>225</i>	<i>110</i>

*Fewer than 25 respondents in category

Table 7.14
 Percentage distribution of primary reason for not using contraception, by union status
 1999 Ukraine Reproductive Health Survey

Reason for not using contraception	Total	Union status		
		Currently in union	Previously in union	Never in union
<u>Reasons related to pregnancy, fecundity, or sexual activity</u>	66.7	62.7	80.8	59.3
Not sexually active	25.4	3.9	64.8	48.6
Subfecund	13.5	18.4	8.7	0.8
Pregnant	12.6	18.9	1.4	5.5
Want pregnancy	8.2	11.7	2.1	3.6
Difficult to get pregnant	7.0	9.8	3.8	0.8
<u>Other reasons</u>	33.3	37.3	19.2	40.7
Occasional sex only	8.8	4.7	10.1	24.7
Partner objections	2.7	3.2	0.6	4.3
Fear of side / health effects	2.7	3.9	1.2	0.0
Cost / availability / difficult to use	2.6	3.2	0.6	3.2
Breastfeeding / postpartum	1.9	3.0	0.0	0.0
Doctor will not prescribe	1.6	2.2	1.3	0.0
Haven't bothered	1.0	1.5	0.4	0.0
Previous side effects	0.6	0.9	0.0	0.0
Religion	0.6	1.0	0.1	0.0
Prefer abortion	0.1	0.2	0.0	0.0
Other	2.6	3.4	0.8	2.2
Don't know	8.2	10.3	4.2	6.3
Total	100.0	100.0	100.0	100.0
<i>Number of women</i>	<i>1,391</i>	<i>877</i>	<i>372</i>	<i>142</i>

Table 7.15
Percent of women with unmet need for family planning, according to two definitions
by age, region, residence, and education
1999 Ukraine Reproductive Health Survey

<u>Characteristic</u>	Definition I*	Definition II**
Total	14.9	37.0
<u>Age</u>		
15-24	10.6	23.7
25-34	16.8	43.8
35-44	17.3	43.5
<u>Region</u>		
North	14.6	36.9
Central	15.5	38.9
East	15.8	33.5
South	13.2	31.8
West	14.8	46.5
<u>Residence</u>		
Urban	13.9	33.8
Rural	17.5	45.6
<u>Living children</u>		
0	7.4	14.1
1	19.0	43.2
2+	17.4	50.1
<u>Education</u>		
< Complete secondary	11.7	22.9
Complete secondary	16.7	40.2
> Complete secondary	10.7	34.3

***Definition I:** Women are considered to be in need if they are fecund, sexually active or in union, not pregnant, did not want to get pregnant at the time of interview, and are not using any type of contraception.

****Definition II** is the same as definition I, except that it also includes women using typically less effective methods of contraception (withdrawal and periodic abstinence).

Table 7.16
 Percentage distribution of users of selected contraceptive methods according to whether they prefer to use a different method, by current and preferred method
 1999 Ukraine Reproductive Health Survey

Respondent's current method	Contraceptive method preferred by women											<i>No. of users</i>
	Prefers current method	Prefers other method	IUD	Pills	Steriliz. (f. or m.)	AM- after pills	Spermicide	Condom	Natural methods	Other	Not Sure	
IUD	89.9	10.1	0.0	2.5	1.4	0.9	0.8	0.3	0.6	2.1	1.5	997
Tubal ligation	88.6	11.4	8.0	0.0	0.0	0.9	0.0	0.6	0.0	0.0	1.9	68
Periodic abst.	77.9	22.1	9.5	3.9	2.0	0.6	0.8	2.1	0.2	1.7	1.4	556
Pills	76.9	23.1	10.8	0.0	1.0	0.2	2.4	0.0	3.5	2.3	2.8	198
Spermicide	75.6	24.4	10.1	1.6	1.2	0.8	0.0	0.0	3.5	3.9	3.5	53
Condoms	69.9	30.1	12.2	6.5	1.9	2.0	1.9	0.4	1.3	3.2	0.7	1,014
Withdrawal	62.7	37.3	16.0	5.7	1.6	2.9	2.3	2.5	1.5	1.7	3.1	970
All methods	74.8	25.2	9.7	4.4	1.7	1.7	1.5	1.2	1.1	2.1	1.8	4,037 *

*Includes methods with fewer than 50 users, which are not listed separately.

Table 7.17
 Percentage distribution of reason not using preferred method of contraception, by residence
 1999 Ukraine Reproductive Health Survey

Reason not using preferred method	Total	Residence	
		Urban	Rural
Cost	27.2	23.1	38.0
Fear of side / health effects	20.2	19.8	20.3
Doctor will not prescribe	15.3	16.6	11.9
Do not know enough about / how to obtain it	13.6	14.9	10.0
Husband objects to it	4.2	4.4	3.6
Poor effectiveness	2.9	3.1	2.7
Current method is permanent / long term	1.9	2.5	0.5
Difficult to get	1.3	1.5	0.7
Method is not natural	0.9	1.2	0.2
Religious reasons	0.4	0.4	0.5
Other	8.0	8.8	6.6
Don't know	4.1	3.7	5.0
Total	100.0	100.0	100.0
<i>Number of respondents</i>	<i>881</i>	<i>674</i>	<i>207</i>

Table 7.18
 Percentage distribution of major problems or concerns with current contraceptive method, by current method
 1999 Ukraine Reproductive Health Survey

Current method	Major problems or concerns								Total	No. of users
	None	Side effects	Health concerns	Cost/availability	Effective-ness	Incon-venient	Partner disapproval	Other		
IUD	84.9	4.5	6.2	0.5	2.0	0.4	0.1	1.4	100.0	993
Condoms	84.3	0.9	1.8	0.7	7.4	3.9	0.7	0.2	100.0	971
Oral contraceptives	76.9	9.6	2.7	5.8	3.1	0.0	0.7	0.0	100.0	197
Sterilization	86.8	1.9	6.3	0.0	0.0	0.0	0.0	5.0	100.0	69
Vaginal methods	81.8	0.8	4.4	8.3	4.8	0.0	0.0	0.0	100.0	52
Traditional methods	73.1	0.9	2.7	0.2	16.3	3.0	3.6	0.1	100.0	1,690
Combined	89.4	2.9	0.0	0.9	3.8	2.9	0.0	0.0	100.0	42
All methods *	79.2	2.2	3.4	0.8	9.6	2.3	1.8	0.5	100.0	4,037

* Includes method with fewer than 25 users, which are not listed separately.

Table 7.19
 Contraceptive failure rates after one, two, and three years for selected methods of contraception,
 contraceptive episodes starting since January 1994
 1999 Ukraine Reproductive Health Survey

Duration	Contraceptive Method					
	All methods	IUD	Oral contracept.	Condoms	Periodic abstinence	Withdrawal
<u>All Women</u>						
1 year	.086	.014	.059	.071	.156	.117
2 years	.145	.029	.106	.135	.240	.203
3 years	.188	.035	.132	.187	.298	.272
<i>Contraceptive episodes</i>	9065	1784	706	2361	1408	2179
<u>Urban</u>						
1 year	.089	.015	.067	.070	.160	.128
2 years	.148	.031	.105	.130	.235	.226
3 years	.189	.038	.135	.180	.294	.290
<i>Contraceptive episodes</i>	7293	1390	611	2096	1125	1527
<u>Rural</u>						
1 year	.076	.012	*	.083	.142	.096
2 years	.137	.022	*	.171	.259	.167
3 years	.183	.026	*	.233	.313	.242
<i>Contraceptive episodes</i>	1772	394	95	265	283	652

*Number of contraceptive episodes too small to estimate a reliable failure rate

Table 7.20
 Contraceptive discontinuation rates (all reasons) after one, two, and three years
 for selected methods of contraception, contraceptive episodes starting since January 1994
 1999 Ukraine Reproductive Health Survey

Duration	Contraceptive Method					
	All methods	IUD	Oral contracept.	Condoms	Periodic abstinence	Withdrawal
<u>All Women</u>						
1 year	.285	.064	.543	.361	.328	.293
2 years	.412	.134	.693	.513	.474	.437
3 years	.494	.213	.740	.604	.553	.529
<i>Contraceptive episodes</i>	9065	1784	706	2361	1408	2179
<u>Urban</u>						
1 year	.298	.067	.532	.352	.328	.340
2 years	.429	.143	.698	.504	.476	.493
3 years	.511	.225	.729	.599	.557	.577
<i>Contraceptive episodes</i>	7293	1390	611	2096	1125	1527
<u>Rural</u>						
1 year	.245	.056	*	.415	.329	.206
2 years	.358	.108	*	.564	.469	.335
3 years	.438	.178	*	.639	.540	.442
<i>Contraceptive episodes</i>	1772	394	95	265	283	652

*Number of contraceptive episodes too small to estimate a reliable failure rate

Table 7.21
 One-year contraceptive discontinuation rates for selected reasons for selected contraceptives methods,
 contraceptive episodes starting since January 1994
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Primary reason for discontinuing method						
	Side effects	Health concerns	Physician's decision	Give body a rest	For a better method	Cost/Supply problems	Inconvenient method
Oral Cont.	.094	.083	.067	.150	.030	.072	.028
IUD	.102	.021	*	*	*	*	*
Condoms	*	*	*	*	.067	.035	.033
Periodic abstinence	*	*	*	*	.097	*	.011
Withdrawal	*	*	*	*	.063	*	.018

*Discontinuation rate for this reason was less than .01

Table 7.22
 Percentage distribution of primary reason for discontinuing selected contraceptive methods,
 contraceptive episodes starting since January 1994
 1999 Ukraine Reproductive Health Survey

Reason for discontinuing method	Contraceptive method				
	Oral Contraceptives	IUD	Condoms	Periodic abstinence	Withdrawal
Pregnant while using	10.7	9.4	21.0	43.1	41.4
No sex/ Relationship ended/ Can't get pregnant	5.5	2.8	11.3	7.9	8.3
Wanted better method	3.5	0.9	14.7	21.1	16.4
Wanted to get pregnant	9.8	8.2	12.9	10.9	10.3
Inconvenient method	3.2	1.3	5.9	2.8	3.3
Side effects	12.5	7.2	1.6	0.0	0.6
Physician's decision	8.6	21.4	3.0	2.7	3.5
Health concerns	11.9	15.6	1.6	1.8	1.1
Partner objected	0.9	0.4	11.8	2.0	10.2
Supply/ Cost	7.9	0.2	6.6	0.7	0.5
Give body a rest	21.2	25.8	3.6	1.2	0.7
Other	2.3	6.4	3.3	2.0	1.3
Don't remember	2.1	0.6	2.8	3.8	2.5
Total	100.0	100.0	100.0	100.0	100.0
<i>No. of contraceptive episodes</i>	<i>517</i>	<i>829</i>	<i>1423</i>	<i>878</i>	<i>1251</i>

Table 7.23
 Percent of women with poor opinion of selected aspects of various contraceptive methods, by residence,
 according to use of that method
 1999 Ukraine Reproductive Health Survey

Characteristic	Contraceptive method						
	Oral contracep.	IUD	Injectable	Condoms	Female steriliz.	Induced abortion*	Mini- abortion*
<u>Overall opinion</u>							
Uses the method	8.4	3.1	25.3	13.4	38.4	96.0	95.3
Uses another	60.7	51.1	72.6	47.9	80.3	96.2	95.3
Uses no method	56.7	52.3	74.8	42.7	82.4	NA	NA
Total	57.3	49.7	73.4	40.5	80.5	96.2	95.3
<u>Safety and health</u>							
Uses the method	8.8	7.6	25.3	5.5	19.6	89.7	83.9
Uses another	36.5	29.7	51.9	9.0	56.8	92.0	88.4
Uses no method	36.5	32.3	54.0	9.7	59.4	NA	NA
Total	35.7	27.1	52.7	8.8	57.3	91.1	87.7
<u>Effectiveness</u>							
Uses the method	2.0	1.7	0.0	3.8	6.9	NA	NA
Uses another	11.2	10.3	11.9	11.3	6.2	NA	NA
Uses no method	12.5	9.1	15.1	10.7	7.7	NA	NA
Total	11.4	8.4	13.1	9.9	6.8	NA	NA
<u>Cost</u>							
Uses the method	14.8	23.3	0.0	60.9	12.9	8.4	10.7
Uses another	5.8	13.7	4.4	55.8	4.2	7.3	8.0
Uses no method	5.7	11.7	3.9	58.2	4.4	NA	NA
Total	6.1	14.8	4.2	57.5	4.4	7.7	8.5

NOTE: Respondents with no opinion have been deleted from the estimates for corresponding cells.
 *For induced abortion and miniabortion, "uses the method" includes women, who ever had an abortion of that type. "Uses another" includes women who have never had an abortion of that type.

Table 7.24
 Percentage of users of withdrawal or periodic abstinence who said that selected factors were important
 in their decision not to use a modern contraceptive method, by residence.
 1999 Ukraine Reproductive Health Survey

Selected factors	Total		Urban		Rural	
	Very important	Somewhat important	Very important	Somewhat important	Very important	Somewhat important
Difficulty of getting other methods	21.2	30.6	19.4	29.9	24.8	32.1
Cost of other methods	39.9	26.6	36.6	27.8	46.4	24.1
Lack of information about other methods	20.7	26.7	20.7	27.9	20.6	24.3
Health / Side effects of other methods	64.4	16.3	65.6	16.4	62.0	16.2
Husband / partner preference	31.9	33.3	32.1	32.2	31.4	35.4
Religious beliefs	8.1	12.5	6.8	10.9	10.7	15.7
Naturalness of the method	61.0	20.9	61.1	19.9	60.7	22.9
<i>Number of users</i>	1,689	1,689	1,214	1,214	457	457

Table 7.25
 Percentage distribution of perceived effectiveness of modern methods compared to method currently used,
 by residence, current users of withdrawal or periodic abstinence
 1999 Ukraine Reproductive Health Survey

Perceived effectiveness	Total	Urban	Rural
Current method more effective	33.9	32.2	37.5
About equally effective	35.2	37.8	29.8
Current method less effective	19.1	19.8	17.7
Don't know	11.8	10.2	14.9
Total	100.0	100.0	100.0
<i>Number of women</i>	1,688	1,213	475

Table 7.26
 Percentage distribution of time of most recent IUD insertion and length of time provider said
 IUD could be worn, by residence, women with insertion since January 1994
 1999 Ukraine Reproductive Health Survey

	Total	Urban	Rural
Time of IUD insertion			
Immediately after delivery	6.6	7.4	4.4
Immediately after abortion	22.5	23.4	19.9
Neither	70.9	69.2	75.7
Length of time provider said IUD could be worn			
Less than 4 years	23.3	23.5	22.9
4-5 years	50.6	50.5	50.7
6-10 years	6.0	6.0	5.9
More than 10 years	0.1	0.2	0.0
As long as she wants	4.0	3.8	4.8
Other	0.7	0.7	0.6
Physician did not say	6.4	5.8	8.1
Don't remember	8.9	9.5	7.0
Total	100.0	100.0	100.0
<i>Number of women</i>	1,552	1,211	341

Table 7.27
 Percentage distribution of the most serious problem women have had related to their IUD,
 by residence, women with insertion since January 1994
 1999 Ukraine Reproductive Health Survey

Problem	Total	Urban	Rural
No problems	69.1	68.5	70.5
Heavy bleeding during menstrual period	8.4	9.9	4.4
Bleeding/Spotting between menstrual periods	6.3	6.1	6.8
Inflammation/Infection/Discharge	6.9	6.0	9.5
Cramping	2.0	1.9	2.2
Husband/Partner complained about string	0.3	0.2	0.6
Other	3.5	3.3	4.1
Do not remember	3.5	4.0	2.1
Total	100.0	100.0	100.0
<i>Number of women</i>	<i>1,552</i>	<i>1,211</i>	<i>341</i>

Table 7.28
 Percentage distribution of length of time physician told women they could take oral contraceptives,
 by residence, women who started taking oral contraceptives since January 1994
 1999 Ukraine Reproductive Health Survey

Length of time provider said woman could take oral contraceptives	Total	Urban	Rural
Less than 2 years	13.9	12.7	20.5
2 - 5 years	3.9	4.3	1.5
6 - 15 years	0.6	0.6	0.0
As long as woman wants	25.1	26.2	18.8
Other	3.8	3.7	4.6
Physician did not say	42.8	42.0	47.7
Don't remember	9.9	10.5	6.9
Total	100.0	100.0	100.0
<i>Number of women</i>	<i>1,552</i>	<i>1,211</i>	<i>341</i>

Table 7.29
 Percentage distribution of the most serious problem women have had related to oral contraceptives,
 by residence, women who started taking oral contraceptives since January 1994
 1999 Ukraine Reproductive Health Survey

Problem	Total	Urban	Rural
No problems	58.8	59.6	54.8
Nausea	8.4	7.7	12.3
Bloating /weight gain	8.3	7.8	11.2
Headaches	5.6	4.9	9.2
Bleeding between menstrual periods	5.3	6.3	0.0
Vision problems	0.0	0.0	0.0
Other	8.3	8.1	9.6
Do not remember	5.2	5.6	2.9
Total	100.0	100.0	100.0
<i>Number of women</i>	<i>551</i>	<i>481</i>	<i>70</i>

Table 7.30
 Percent distribution of interest in being contraceptively sterilized, by selected characteristics,
 fecund respondents in union who want no more children
 1999 Ukraine Reproductive Health Survey

Characteristic	Interested	Not interested	Not sure	Total	N
Total	6.6	87.9	5.5	100.0	2,268
<u>Region</u>					
North	5.9	89.2	4.9	100.0	266
Central	6.9	85.3	7.8	100.0	216
East	7.3	88.9	3.8	100.0	857
South	7.6	85.1	7.2	100.0	535
West	4.8	88.9	6.3	100.0	394
<u>Residence</u>					
Urban	7.0	88.3	4.7	100.0	1,722
Rural	5.4	87.0	7.6	100.0	546
<u>Age</u>					
15-19	--	--	--	100.0	12*
20-24	11.1	79.9	8.8	100.0	147
25-29	5.5	89.3	5.2	100.0	337
30-34	8.5	85.3	6.2	100.0	540
35-39	5.6	88.3	6.1	100.0	662
40-44	5.4	90.9	3.8	100.0	570
<u>Living Children</u>					
0	2.3	88.7	8.9	100.0	29
1	6.6	88.9	4.5	100.0	727
2	6.2	87.4	6.4	100.0	1,323
3+	9.7	87.4	2.9	100.0	189
<u>Education</u>					
< Complete secondary	1.3	94.5	4.2	100.0	104
Complete secondary	6.5	87.7	5.7	100.0	1,628
> Complete secondary	7.6	87.2	5.2	100.0	536
<u>Current contraceptive method</u>					
IUD	5.5	89.1	5.5	100.0	660
Condoms	8.1	87.9	3.9	100.0	414
Oral contraceptives	7.4	87.7	4.9	100.0	92
Periodic abstinence	6.5	91.1	2.5	100.0	360
Withdrawal	6.9	85.1	8.1	100.0	589
No contraception	3.0	88.9	8.1	100.0	106

*Fewer than 25 women in category.

Table 7.31
 Percentage distribution of primary reason not interested in contraceptive sterilization, by residence
 1999 Ukraine Reproductive Health Survey

Reason not interested in sterilization	Total	Urban	Rural
Haven't thought about it	33.7	34.4	31.9
Health risks	25.2	26.1	23.0
Don't know enough about sterilization	9.6	8.9	11.0
Fear of operation	9.3	7.9	12.7
Might want another child	8.2	8.9	6.4
Cost of an operation	3.5	3.3	4.0
Not culturally / socially acceptable	2.2	1.9	2.9
Husband / partner would object	2.1	2.0	2.2
Religious reasons	0.8	0.7	0.9
Not sexually active	0.3	0.3	0.2
Other	4.3	4.7	4.4
Don't know	0.8	0.9	0.4
Total	100.0	100.0	100.0
<i>Number of women</i>	<i>1,996</i>	<i>1,521</i>	<i>475</i>

CHAPTER VIII

CONTRACEPTIVE COUNSELING

An important component of the Ukraine Women's Reproductive Health Initiative has been the development and implementation of interventions intended to improve health care workers' counseling regarding contraceptive services. The URHS questionnaire included a series of questions designed to determine the kinds of interaction women in the survey sites typically have with family planning providers regarding issues such as whether to use contraception, the information given to women about contraceptive methods, method selection, and satisfaction with the services received. It also examined various aspects of communications between partners regarding such issues as contraceptive use and abortion.

Post-abortion/Post-partum counseling

The top panel of Table 8.1 presents results regarding post-abortion contraceptive counseling. Only thirty-nine percent of women who had an induced abortion since the beginning of 1994 said that a health professional had spoken with them about ways of preventing an unintended pregnancy following their most recent abortion. The proportion was virtually identical in urban and rural areas. The likelihood that a woman would be referred for contraceptive services or counseling was very low, only 7%. Only 16% of women undergoing an induced abortion left the facility where it was performed with a contraceptive method or a prescription for one.

The bottom panel of Table 8.1 shows information on contraceptive counseling following recent deliveries. Only one-fourth of respondents with recent live births reported that a doctor or nurse talked to them (or offered to talk to them) about contraception subsequent to their delivery, a considerably lower proportion than after abortions. The proportion who were referred for counseling or services was only 5%. Only 4% actually left the delivery facility with a contraceptive method or a prescription for one.

These figures show a clear need for improved and expanded post-abortion and post-delivery counseling, referrals, and provision of contraceptive supplies.

Content of Counseling/Method selection

Although a survey like the URHS cannot capture all the important interactions between family planning providers and clients, women were asked a number of questions to try to determine the extent to which health care workers provided some basic information and services. Table 8.2 describes information from women who had used oral contraceptives, the IUD, or injectable contraceptives since January of 1994, and refers to the last time they started using any of these methods.

High quality contraceptive counseling should stress that women/couples are able ultimately to select their own contraceptive method, rather than the provider making the decision unilaterally. The provider should discuss the various available methods, giving the client as much useful information as possible, in order that she/they can make a well informed decision. Each respondent first

reported whether her family planning provider had discussed the various family planning options available to her. Just over half (54%) of women who recently started using a provider supplied method responded positively (Table 8.2). Differences according to most characteristics in the percentages of women with whom providers discussed contraceptive options tended to be small. Percentages were lowest for the youngest women and highest for the oldest. They also increased with education and were higher for IUD users than pill users.

Following method selection, the provider should give information on how to use the method, possible side effects associated with the method, and when the client should return for follow-up. Table 8.3 (first column) shows that about two-thirds of recent IUD, pill, and injection users recalled their provider giving information on potential side effects and what to do about them. Of those respondents who received any contraceptive counseling, large majorities said that they: understood most or all of the information provided on the method selected (87%); were told when to return for follow-up (82%); and were given a pelvic exam during their visit (87%). IUD users were considerably more likely than pill users to have received more complete counseling. Among pill users, only 50% had possible side effects explained to them, 52% were told when to return for follow-up, and 63% received a pelvic exam.

Partner involvement in reproductive health decisions

It is preferable that both partners be involved in such decisions as whether to use contraception and what type of method to use. Table 8.4 reveals that 82% of married respondents had had discussions about family planning with their partner. Although differentials in this table are not great, the percentages who had discussions were lowest in the South and West, in rural areas, among the youngest and oldest respondents, and among the least well educated. A large majority of respondents (82%) said that the decision regarding what contraceptive method to use should be made jointly by the man and woman. Only 13% felt this decision should be up to the woman alone.

In the vast majority of instances (97%), women reported that their partner was aware of their most recent abortion. Almost 9 of every 10 women who opted for an abortion discussed with her partner whether she should have the procedure (Table 8.5). Although the difference was not significant, this percentage was slightly lower in the West (86%) and among the oldest cohort of women (86%).

Table 8.1
 Percent of women who received various family planning services
 after their most recent delivery or abortion, by residence
 among women who had a delivery or abortion since January 1994
 1999 Ukraine Reproductive Health Survey

Type of Service	Residence		
	Total	Urban	Rural
<u>Post-Abortion</u>			
Talked to About Ways to Prevent Pregnancy	39.4	39.4	39.2
Referred for Contraceptive Services or Counseling	6.6	6.9	5.7
Left Facility with Contraceptive Method or Prescription	15.6	16.0	14.4
<i>Number of Women</i>	<i>1,435</i>	<i>1,168</i>	<i>267</i>
<u>Post-Delivery</u>			
Doctor or Nurse Offered to Discuss Contraception	25.5	26.3	23.9
Referred for Contraceptive Services or Counseling	4.5	4.8	4.0
Left Facility with Contraceptive Method or Prescription	3.8	4.2	3.2
<i>Number of Women</i>	<i>1,724</i>	<i>1,269</i>	<i>455</i>

Table 8.2
 Percent of women starting use of IUD, oral contraceptives, or injections since January 1994
 with whom family planning provider discussed contraceptive options and
 percentage distributions of who selected method
 1999 Ukraine Reproductive Health Survey

Characteristic	% discussed with provider	Who selected method				Total	No. of Women
		Woman	Provider	Both	Not sure		
<u>All women</u>	53.5	57.7	22.5	18.4	1.4	100.0	1,978
<u>Region</u>							
North	46.4	62.3	22.1	15.2	0.4	100.0	235
Central	53.9	50.0	22.9	24.5	2.7	100.0	190
East	56.0	58.0	23.7	17.5	0.8	100.0	803
South	50.8	59.1	18.8	19.1	3.0	100.0	523
West	57.3	57.0	24.4	18.1	0.5	100.0	227
<u>Residence</u>							
Urban	53.0	58.2	22.7	17.8	1.3	100.0	1,581
Rural	54.7	56.3	21.8	20.4	1.6	100.0	397
<u>Age</u>							
15-19	31.3	71.0	14.3	14.7	0.0	100.0	28
20-24	50.5	61.5	19.1	17.4	2.0	100.0	262
25-29	52.6	57.1	23.0	18.3	1.6	100.0	447
30-34	53.9	55.8	21.7	21.2	1.3	100.0	469
35-39	53.0	57.0	23.9	17.6	1.5	100.0	462
40-44	59.0	58.5	24.3	16.8	0.4	100.0	310
<u>Education</u>							
<Comp secondary	46.8	54.6	22.4	21.2	1.8	100.0	98
Comp. Secondary	53.4	57.9	22.5	18.3	1.3	100.0	1,387
>Comp secondary	57.8	57.8	22.6	18.1	1.5	100.0	493
<u>Method used*</u>							
IUD	56.3	57.7	21.1	20.0	1.2	100.0	1,484
Oral contraceptives	44.8	58.5	26.7	13.0	1.8	100.0	482

*12 users of injections, not shown separately

Table 8.3
 Percent of women starting use of IUD, oral contraceptives, or injections since January 1994
 to whom provider explained about possible side effects* and percent of women receiving counseling who
 1) understood most of information on side effects; 2) were told when to return for follow-up, refill, etc.;
 and 3) were given a pelvic examination, by region, residence, and method
 1999 Ukraine Reproductive Health Survey

Characteristic	Of women receiving method from professional		Of women receiving counseling			N
	% possible side effects explained*	N	% understood most or all of explanation	% told when to return, etc	% given pelvic exam	
All women	65.3	1,920	86.5	82.0	86.9	1,643
<u>Region</u>						
North	62.3	231	88.0	83.3	84.4	192
Central	66.0	187	81.2	84.9	87.9	164
East	67.5	762	87.6	81.6	87.2	651
South	62.3	519	88.0	80.2	86.5	444
West	66.5	221	84.4	81.8	88.5	192
<u>Residence</u>						
Urban	64.7	1,529	86.5	81.3	86.1	1,315
Rural	67.1	391	86.5	84.2	89.6	328
<u>Method used</u>						
IUD	70.1	1,446	87.1	90.1	93.5	1,285
Oral contraceptives	50.3	463	83.9	52.2	62.8	349

*Only women who were provided contraception by a physician, nurse or midwife.

Table 8.4
Percent of women in union who have discussed use of family planning with their partner and percentage distribution of who women think should make decisions about whether to use contraception, by selected characteristics
1999 Ukraine Reproductive Health Survey

<i>Characteristic</i>	<i>% discussed with partner</i>	<i>Who should select method</i>				<i>Total</i>	<i>Women</i>
		<i>Woman</i>	<i>Partner</i>	<i>Both</i>	<i>Not sure</i>		
<u>All women</u>	81.6	13.4	2.5	81.5	2.1	100.0	4,409
<u>Region</u>							
North	84.8	12.9	1.9	82.7	2.5	100.0	516
Central	84.1	8.6	3.7	84.9	2.9	100.0	382
East	83.2	14.7	2.6	81.4	1.4	100.0	1,627
South	79.7	13.7	1.6	82.1	2.6	100.0	1,115
West	77.0	13.8	3.0	78.6	4.7	100.0	769
<u>Residence</u>							
Urban	82.3	14.0	2.1	81.9	2.1	100.0	3,340
Rural	79.7	11.9	3.6	80.6	4.0	100.0	792
<u>Age</u>							
15-19	76.9	8.6	3.4	80.6	7.4	100.0	120
20-24	85.4	12.0	1.9	83.6	2.5	100.0	724
25-29	83.5	11.6	2.8	83.8	1.8	100.0	955
30-34	82.2	13.5	2.2	81.7	2.6	100.0	947
35-39	82.1	15.5	2.3	79.2	3.1	100.0	886
40-44	75.1	15.0	3.2	79.2	2.7	100.0	777
<u>Education</u>							
<Comp secondary	77.7	10.8	3.1	78.2	8.0	100.0	273
Comp. Secondary	80.9	13.6	2.7	81.5	2.3	100.0	3,162
>Comp secondary	84.9	13.3	1.9	82.4	2.4	100.0	974

Table 8.5
Percent of women whose partner knew about her most recent abortion and
percent who discussed with her partner whether to have an abortion, by selected characteristics
1999 Ukraine Reproductive Health Survey

Characteristic	% whose partner knew about most recent abortion	% who discussed with partner whether to have abortion	<i>Number of Women*</i>
<u>All women</u>	97.1	89.2	2,369
<u>Region</u>			
North	97.2	93.0	281
Central	98.2	90.6	221
East	97.6	89.0	951
South	97.9	88.4	659
West	93.9	85.5	257
<u>Residence</u>			
Urban	96.9	90.1	1,883
Rural	97.9	88.9	486
<u>Age</u>			
15-19	*	*	11
20-24	96.9	88.7	210
25-29	97.6	91.8	438
30-34	98.0	91.8	556
35-39	96.7	88.0	602
40-44	96.5	86.3	552
<u>Education</u>			
<Complete secondary	98.3	89.4	124
Complete Secondary	97.2	89.6	1,733
>Complete secondary	96.5	87.8	512

*Excludes women who did not have a partner at the time of abortion.

CHAPTER IX

INFORMATION, EDUCATION, AND COMMUNICATION (IEC)

The 1999 URHS included a series of questions regarding respondents' mass media habits, preferences, and attitudes. This included information on television viewing, radio listening, and newspaper reading. In addition, the survey assessed exposure to reproductive health information on television and in print in the six months prior to the interview, as well as opinions about the acceptability of placing family planning information on television and radio. These findings have programmatic implications for the development and placement of IEC messages in future interventions to improve reproductive health and the utilization of services.

Television viewing habits

Eighty percent of respondents said that they watch television virtually every day (Table 9.1). On the other hand, only about 8% of respondents said they generally watched television less than once per week. The greatest difference in viewing frequency appears to be between urban and rural women, 83% and 73% respectively watching television daily. Women 35-44 years were slightly less likely to be infrequent viewers. There was no consistent relationship apparent between education and television viewing. Such widespread overall viewership indicates that television is a promising means for reaching women with health and family planning information, even though commercial time is expensive.

The most popular types of television programs were soap operas, frequently watched by 60% of women (Table 9.2). Soap operas were most popular in rural areas (66%), in the West region (67%) and among women who did not complete secondary school (67%). Other types of television shows that were popular included entertainment program (watched by 55%), news (54%) and music programs (52%), the only other types of programs regularly watched by a majority of respondents. Just over one-third of respondents regularly watched women's programs, while 27% watched health programs. Few respondents said that they regularly watched religious programs, business programs, or sports. It was not surprising to find that entertainment and music programs were most popular among the youngest cohort of women, while women at older ages were about twice as likely as younger respondents to watch news and political programs. Women's programs and health programs, which are the most typical place to air health information messages, were most popular among women over 25 years old, who lived in urban areas, and had higher education. However, to reach the large proportions of women who do not watch those programs, especially the less educated and younger women, it might be beneficial to place information on soap operas and music and entertainment programs

The heaviest viewing times for television were in the evening, with 67% of viewers reporting that they watched television between 6 p.m. and 10 p.m. (Table 9.3). The only other time of high viewership was after 10 p.m. (35%). Television viewing after 6 p.m. was lower among the youngest respondents and those with incomplete secondary education, although still much heavier than at other times of the day even for those groups. About 11% of respondents watched television between noon and 6 PM. The youngest respondents, under 24 years old, and those who did not complete secondary school were the most likely to watch television during the morning or afternoon. About 20% of these women watched TV between noon and 6PM.. At this time of day,

health messages might be included in entertainment or music program, which are most popular among younger, less educated women.

Radio listening habits

Half of all respondents said that they generally listened to the radio daily (Table 9.4). Another 8% listened to the radio at least once a week. About four of every ten respondents said that they rarely or never listen to the radio. Urban women, 15-24 year-olds, and women from the East region, and those with more than secondary education tend to listen to the radio more often than other groups.

Most radio listeners, listened to music and news programs (50% and 34%, respectively, of the 60% who listened) (Table 9.5). About one in every ten women reported listening to women's programs and health programs. Almost about one-half as many women listened to health and women programs in the South region as elsewhere. Both health and women's programs were most popular among the best educated women and those over 35-44 years old and least popular among the youngest and the least well educated.

Radio listening times were much less concentrated than television watching times, as might have been expected (Table 9.6). The largest percentages of radio listeners said they had no regular listening times (26% of all women) or listen in the morning (25%). Respondents were much less likely to listen to radio regularly in the afternoon (16%) or in the evening (12%) than in the morning. Only about 4% of women said they listened to the radio after 10PM.

Newspaper readership

URHS respondents could be split into three roughly equal-sized groups in regard to frequency of reading newspapers. Just under one-third of all women said they almost never read newspapers, another one-third read newspapers at least three times per week, and a slightly larger group (38%) read them occasionally (twice per week or less)(Table 9.7). Daily newspaper reading, 18% overall, was most common in the North and Central regions of Ukraine, where respectively 31% and 21% of the women said they read a newspaper daily. compared with about half as many in South and West regions (13% and 14%, respectively). Frequency of newspaper reading increased with the age and especially with the level of education of respondents. There was little difference, however, between urban and rural women.

The national newspaper that was most frequently read included *Facti* (Facts), read by 27% of women, most commonly in urban areas, in the North region, and among the most educated women (Table 9.8). In second place was *Telenedelya* (*Television Week*), read by 15% of women and highest in urban areas, the East and South regions, and among the youngest women. Seventeen percent of the respondents said they read local newspapers. These papers were most frequently read by women over 35 years of age, rural residents, and those with higher levels of education.

Exposure to and attitudes about health messages in the media

About one-third of women reported seeing family planning information on television (36%) and 39% recalled seeing such information in newspapers or magazines within six months of interview (Table 9.9). Exposure to such information both on TV and in print decreased with increasing age,

was higher in urban areas than in rural areas, and increased with educational attainment. It also tended to be lower in the West region than it was elsewhere. This indicates that health promotional campaigns may be more likely to reach younger people, often a key group in these types of activities, through television and appropriate print media.

Despite a minority of women recalling seeing information on family planning in the mass media, the vast majority of women felt that such information should be available through the media (Table 9.9, third column). Eighty-five percent of women, with only slight difference among subgroups, said that information on family planning should be broadcast. Women in the West (80%) and 35-44 year-old women (82%) were less likely than others to favor broadcasting such information.

Table 9.10 examines the correlation between exposure to family planning messages in the media correlated and current contraceptive use among fecund sexually active women. It shows that the use of modern contraceptive methods was more common among women who had been exposed to family planning messages in the media than among women who have not been exposed (46% vs. 37%). Use of traditional methods, though, was slightly higher among those who did not report seeing such messages. Without more information, though, it is not possible to draw conclusions about whether exposure to media messages was a cause of greater modern contraceptive use or simply a correlate.

Table 9.1
 Percentage distribution of TV viewing frequency, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Every day	At least once per week	At least once per month	< once per month / Never	Other	Total	<i>Number of women</i>
Total	79.6	12.1	1.4	6.3	0.2	100.0	7,128
<u>Residence</u>							
Urban	82.5	11.3	1.2	4.8	0.2	100.0	5,544
Rural	73.2	14.3	1.8	10.4	0.3	100.0	1,584
<u>Region</u>							
North	82.1	12.4	1.3	4.2	0.0	100.0	858
Central	77.6	13.9	0.9	7.1	0.5	100.0	617
East	82.5	10.7	1.5	3.1	0.2	100.0	2,669
South	78.5	11.9	1.5	7.9	0.2	100.0	1,824
West	76.6	13.6	1.5	8.1	0.2	100.0	1,160
<u>Age</u>							
15-24	82.0	11.4	1.3	5.3	0.0	100.0	2,282
25-34	81.4	10.7	1.5	6.2	0.2	100.0	2,422
35-44	76.5	14.2	1.3	7.6	0.4	100.0	2,424
<u>Education</u>							
< Complete secondary	80.6	9.7	0.9	8.5	0.3	100.0	778
Complete secondary	78.9	12.8	1.5	6.6	0.2	100.0	4,828
> Complete secondary	83.1	11.2	1.3	4.3	0.1	100.0	1,522

Table 9.2
 Percentage of respondents who watch various types of TV programs, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Types of programs watched														N
	Soap operas	Entertainment	News	Music	Women	Health	Politics	Children	Sports	Busi-ness	Religion	Plays/drama	Other	Do not watch	
Total	60.3	55.3	53.6	52.0	35.8	27.1	11.8	8.8	4.2	4.0	2.9	2.6	1.1	6.3	7,128
<u>Residence</u>															
Urban	58.1	58.4	54.8	54.2	37.6	28.2	13.4	9.5	4.5	4.6	2.9	3.1	1.3	4.8	5,544
Rural	66.1	47.1	50.4	46.3	31.1	24.3	7.4	7.1	3.4	2.3	3.1	1.3	0.7	10.4	1,584
<u>Region</u>															
North	58.7	52.5	57.5	52.5	38.7	27.1	13.4	8.2	5.1	3.5	2.8	3.3	1.1	4.2	858
Central	62.7	54.7	52.0	44.6	28.3	23.7	7.7	8.0	4.9	2.1	2.4	1.8	1.4	7.1	617
East	58.5	62.3	54.6	57.7	39.1	29.3	11.8	9.5	3.8	4.1	2.0	3.6	0.6	3.1	2,669
South	55.7	51.4	51.5	50.7	33.8	24.2	11.8	9.4	4.3	4.1	3.1	2.1	2.1	7.9	1,824
West	66.8	49.2	51.6	47.1	33.9	27.9	12.9	8.1	3.7	5.1	4.3	1.3	0.9	8.1	1,160
<u>Age</u>															
15-24	61.6	62.6	35.9	65.7	30.9	20.9	6.7	7.5	5.2	4.6	1.4	1.7	1.1	5.3	2,282
25-34	58.9	54.8	57.4	50.6	38.6	29.2	11.7	12.1	4.2	3.6	3.2	2.6	1.1	6.2	2,422
35-44	60.3	48.6	67.2	39.9	37.9	31.2	16.9	6.8	3.2	3.8	3.9	3.5	1.2	7.6	2,424
<u>Education</u>															
< Comp. secondary	66.5	58.8	31.4	61.1	20.8	17.5	4.8	7.9	3.8	2.2	1.9	2.0	0.4	8.5	778
Comp. secondary	62.4	53.8	52.9	50.8	37.1	27.9	10.2	9.2	3.9	3.2	2.9	2.9	1.0	6.6	4,828
> Comp. secondary	49.6	58.4	68.3	51.2	39.8	30.0	21.1	8.1	5.2	7.7	3.3	2.6	1.7	4.3	1,522

Table 9.3
 Percentage of respondents who watch TV at various times, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Times likely to watch television						Number of Women
	Rarely / Never watch TV	6:00-12:00	12:00-18:00	18:00-22:00	After 22:00	No regular time	
Total	6.3	13.8	11.4	67.0	35.1	15.9	7,128
<u>Residence</u>							
Urban	4.8	14.2	12.6	67.8	33.6	17.9	5,544
Rural	10.4	12.5	8.2	64.9	39.3	10.8	1,584
<u>Region</u>							
North	4.2	13.9	9.2	72.6	37.4	12.6	858
Central	7.1	17.9	7.8	76.5	44.5	4.2	617
East	3.1	14.3	13.9	64.5	28.1	20.4	2,669
South	7.9	12.5	12.9	59.7	32.7	22.2	1,824
West	8.1	11.6	9.3	68.2	41.9	11.9	1,160
<u>Age</u>							
15-24	5.3	17.1	18.9	64.7	32.9	17.2	2,282
25-34	6.2	14.2	8.5	66.9	39.1	15.1	2,422
35-44	7.6	9.9	6.9	69.4	33.2	15.6	2,424
<u>Education</u>							
< Comp. secondary	8.5	18.7	22.7	59.6	28.1	18.6	778
Comp. secondary	6.6	13.4	10.4	66.7	35.3	16.0	4,828
> Comp. secondary	4.3	12.2	8.4	72.3	37.9	14.3	1,522

Table 9.4
 Percentage distribution of frequency of radio listening, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Frequency of listening to radio					Total	Number of women
	Every day	At least once per week	At least once per month	< once per month/Never	Other		
Total	50.0	8.2	1.2	40.4	0.1	100.0	7,128
<u>Residence</u>							
Urban	53.2	8.8	1.2	36.7	0.2	100.0	5,544
Rural	41.5	6.8	1.2	50.4	0.1	100.0	1,584
<u>Region</u>							
North	50.5	10.4	1.1	38.0	0.0	100.0	858
Central	43.8	6.9	0.5	48.8	0.0	100.0	617
East	55.2	8.7	1.3	34.6	0.2	100.0	2,669
South	49.1	6.5	1.1	43.0	0.2	100.0	1,824
West	44.9	8.0	1.5	45.4	0.2	100.0	1,160
<u>Age</u>							
15-24	53.1	9.3	1.2	36.2	0.1	100.0	2,282
25-34	47.3	7.7	1.5	43.4	0.2	100.0	2,422
35-44	49.6	7.8	0.9	41.6	0.1	100.0	2,424
<u>Education</u>							
< Complete secondary	47.6	8.5	1.3	42.6	0.0	100.0	778
Complete secondary	48.4	8.1	1.2	42.1	0.2	100.0	4,828
> Complete secondary	56.7	8.6	1.2	33.4	0.2	100.0	1,522

Table 9.5
 Percentage of respondents who listen to various types of radio programs distribution of radio programs, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Types of programs listened to														Number of women
	Music	News	Health	Women	Commer- cials	Pers. ann- ouncemnts	Politics	Children	Religion	Sports	Busi- ness	Plays/ drama	Other	Don't listen	
Total	50.2	34.1	10.8	10.3	9.1	8.8	7.0	2.4	2.2	2.0	1.7	1.4	2.5	40.4	7,128
<u>Residence</u>															
Urban	53.8	34.6	10.9	10.4	9.6	9.1	7.3	2.4	2.0	2.1	1.9	1.4	2.9	36.7	5,544
Rural	40.4	32.9	10.5	9.9	7.8	7.7	6.3	2.3	2.7	1.7	1.1	1.3	1.7	50.4	1,584
<u>Region</u>															
North	47.6	34.7	12.1	13.5	8.1	6.0	8.2	2.5	2.7	2.3	2.4	2.3	4.9	38.0	858
Central	40.8	32.1	10.7	11.2	5.4	6.4	5.6	2.2	2.1	1.6	0.9	1.8	1.6	48.8	617
East	59.2	35.0	10.9	10.1	9.7	10.1	7.9	2.5	1.4	2.1	1.8	1.2	1.8	34.6	2,669
South	51.1	32.8	6.6	6.4	10.9	8.2	5.5	1.5	1.2	1.9	1.4	0.6	1.9	43.0	1,824
West	41.0	34.3	12.9	11.1	9.3	10.2	6.7	2.8	4.1	1.9	1.5	1.5	3.0	45.4	1,160
<u>Age</u>															
15-24	58.0	26.1	7.7	8.2	7.8	6.3	3.9	1.6	0.9	2.0	1.6	0.8	2.0	36.2	2,282
25-34	47.4	34.7	11.8	10.5	10.0	10.1	9.8	3.0	1.2	1.9	1.8	1.4	2.8	43.4	2,422
35-44	45.2	41.3	12.7	12.1	9.5	9.8	7.4	2.5	3.6	2.0	1.6	2.0	2.8	41.6	2,424
<u>Education</u>															
< Comp. secondary	50.9	21.9	5.7	6.8	6.0	4.9	2.6	2.0	0.6	1.1	1.0	0.7	1.1	42.6	778
Comp. secondary	48.2	33.9	11.2	10.7	8.7	8.9	6.7	2.6	2.7	2.0	1.5	1.5	2.8	42.1	4,828
> Comp. secondary	56.3	41.6	12.4	10.9	12.2	10.2	10.7	1.7	1.7	2.5	2.8	1.5	2.4	33.4	1,522

Table 9.6
 Percentage of respondents who listen to the radio at various times, by age region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Times likely to listen to radio						<i>Number of Women</i>
	Rarely / Never listen	6:00-12:00	12:00-18:00	18:00-22:00	After 22:00	No regular time	
Total	40.4	24.7	16.3	11.5	4.1	26.0	7128
<u>Residence</u>							
Urban	36.7	26.1	16.8	11.3	4.1	28.2	5,544
Rural	50.4	21.1	15.0	12.1	3.8	20.1	1,584
<u>Region</u>							
North	38.0	29.4	13.4	12.6	5.2	24.7	858
Central	48.8	28.2	12.9	9.1	2.2	15.0	617
East	34.6	23.5	19.2	10.6	3.6	32.8	2,669
South	43.0	19.9	14.6	10.6	3.6	30.2	1,824
West	45.4	25.6	16.9	14.3	5.3	17.8	1,160
<u>Age</u>							
15-24	36.2	22.2	19.9	12.7	4.9	28.0	2,282
25-34	43.4	24.5	15.2	10.8	3.9	24.7	2,422
35-44	41.6	27.4	13.9	10.9	3.4	25.3	2,424
<u>Education</u>							
< Comp. secondary	42.6	18.3	17.9	11.5	3.5	26.0	778
Comp. secondary	42.1	23.8	15.2	11.3	3.9	25.8	4,828
> Comp. secondary	33.4	31.3	19.2	12.1	4.7	26.3	1,522

Table 9.7
 Percentage distribution of frequency of reading daily newspapers by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Frequency of reading newspapers					Total	Number of women
	Every day	3 -4 times week	1 -2 times a week	< once per week	Never		
Total	17.6	13.7	26.2	11.7	30.8	100.0	7,128
<u>Residence</u>							
Urban	18.1	13.9	25.8	11.3	30.8	100.0	5,544
Rural	16.1	12.9	27.5	12.7	30.8	100.0	1,584
<u>Region</u>							
North	30.7	17.1	27.1	10.4	14.6	100.0	858
Central	21.3	15.5	15.4	13.3	24.5	100.0	617
East	15.3	12.9	16.8	10.7	34.2	100.0	2,669
South	12.6	8.2	24.3	12.1	42.8	100.0	1,824
West	13.7	16.0	26.6	13.1	30.5	100.0	1,160
<u>Age</u>							
15-24	13.1	12.1	26.7	13.4	34.7	100.0	2,282
25-34	19.0	14.3	25.8	11.6	29.3	100.0	2,422
35-44	20.5	14.6	26.2	10.1	28.5	100.0	2,424
<u>Education</u>							
< Complete secondary	9.5	8.5	27.4	13.7	40.2	100.0	778
Complete secondary	16.5	13.8	26.5	11.6	31.7	100.0	4,828
> Complete secondary	25.8	16.3	24.8	10.9	22.1	100.0	1,522

Table 9.8
 Percentage of women who read specific newspapers, by age, region, residence, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Newspapers read										Number of women
	Facti	Tele-nedelya	Silski Visty	Kiev vedomosti	Golos Ukraini	Uryadovi Kur'er	Business	Vesukrain. Vedomosty	Local	Never	
Total	27.3	14.7	4.8	3.9	2.4	2.4	1.9	1.4	17.1	30.8	7,128
<u>Residence</u>											
Urban	30.3	17.5	1.8	4.5	2.2	2.6	2.3	1.6	16.1	30.8	5,544
Rural	19.2	7.4	12.6	2.6	3.1	1.6	0.7	0.9	19.9	30.8	1,584
<u>Region</u>											
North	56.8	10.5	6.7	10.1	2.9	2.7	2.8	3.6	16.3	14.6	858
Central	21.6	4.2	18.1	4.9	4.5	4.5	0.8	2.5	22.4	24.5	617
East	22.9	23.3	1.3	1.3	1.5	1.4	1.7	0.3	16.2	34.2	2,669
South	18.7	21.1	0.8	3.4	1.7	1.6	2.9	1.2	15.4	42.8	1,824
West	22.7	3.6	5.5	3.8	3.2	3.2	1.3	1.3	17.8	30.5	1,160
<u>Age</u>											
15-24	23.9	17.7	3.7	4.1	2.2	1.4	1.8	0.9	14.2	34.7	2,282
25-34	27.6	14.1	4.7	4.1	2.1	2.3	1.8	1.6	17.4	29.3	2,422
35-44	30.2	12.4	5.9	3.7	1.9	3.3	2.1	1.8	19.7	28.5	2,424
<u>Education</u>											
< Comp. secondary	17.2	15.2	4.4	2.5	1.9	0.6	0.7	0.3	13.5	40.2	778
Comp. secondary	25.4	13.8	5.2	3.1	1.7	1.3	1.1	1.1	17.4	31.7	4,828
> Comp. secondary	38.9	17.6	3.5	7.6	5.1	7.0	5.3	3.1	18.4	22.1	1,522

Table 9.9
 Percent of women who had seen any family planning information on television or in newspapers or magazines
 within the previous six months,
 and percent who think such information should be broadcast, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	% who have seen FP information on TV	% who have seen FP info in newspapers or magazines	% who think FP information should be broadcast	<i>Number of women</i>
Total	35.5	39.4	84.7	7,128
<u>Residence</u>				
Urban	37.5	43.1	83.9	5,544
Rural	30.0	29.4	86.8	1,584
<u>Region</u>				
North	34.5	39.3	90.0	858
Central	34.6	33.3	88.9	617
East	38.4	42.3	83.5	2,669
South	38.8	45.6	85.8	1,824
West	28.9	32.5	79.7	1,160
<u>Age</u>				
15-24	39.1	46.2	87.3	2,282
25-34	37.4	40.4	85.3	2,422
35-44	29.9	31.7	81.7	2,424
<u>Education</u>				
<Comp. secondary	32.7	37.5	81.4	778
Comp. secondary	34.2	36.5	85.4	4,828
>Comp. secondary	41.1	50.2	84.1	1,522

Table 9.10
 Percentage distribution of current contraceptive use, by whether women
 had been exposed to family planning (FP) messages in the media, among fecund sexually active women
 1999 Ukraine Reproductive Health Survey

	Current contraceptive use			Total	<i>Number of women</i>
	Use modern methods	Use traditional methods	Do not use contraception		
Exposed to media FP messages	45.7	32.1	22.2	100.0	2,574
Not exposed to media FP messages	37.4	35.8	26.8	100.0	1,817
All women	42.2	33.7	24.1	100.0	4,391

CHAPTER X

SEXUAL EXPERIENCE

The URHS included a series of questions designed to study certain aspects of sexual behavior among survey respondents. All women were asked the month, year, and age at which they first had sexual intercourse, the length of time since they most recently had sexual intercourse, the frequency of sexual intercourse during recent months, and the number of recent and lifetime sexual partners. Since the issues of unintended pregnancy and sexually transmitted diseases among young adults have been of growing importance in eastern Europe and the former Soviet Union (as in much of the rest of the world), a set of questions regarding the initiation of sexual activity was asked of respondents between the ages of 15 and 24. These included questions on the respondent's relationship to her first sexual partner and use of contraception at first premarital intercourse, either to prevent pregnancy or prevent STDs.

It is traditional for Ukrainian women to marry and start childbearing at younger ages than in the most of western Europe and developed countries in other parts of the world. It has not been unusual for young couples to stay with their family, sharing income and responsibilities in raising children with parents and sometimes with grandparents. Sex education during the Soviet period tended to be neither comprehensive nor very informative, besides being almost inaccessible for most young women. Since premarital sex was customarily considered to be morally unacceptable (and abortion was available when pregnancy did occur), premarital childbearing was not very common. Lack of contraceptive knowledge and shame led to a very high percentage of non-marital pregnancies among teenagers resulting in abortions. Youth in rural areas, where the stigma associated with premarital pregnancy is even stronger, were more affected than others.

However, the issue of teen pregnancy was not a major concern of the Ukrainian public health establishment during the Soviet period. The economic, social, and political changes that took place in Ukraine after the fall of the Soviet Union, including the transformation of the country to a more open society with exposure to mass media and Western culture, changed many aspects of life among Ukrainians. Among the areas where changes have been occurring are sexual behavior, reproductive health, and marital and motherhood norms. Changes in the social and economic situation may be affecting the ability of women/families to support children born to unmarried teen mothers. Another cause for concern is the increase in sexually transmitted infections among the population, especially among young adults.

Sexual experience of all women

It has been hypothesized by some in the region that the dramatic declines in fertility and pregnancy rates in recent years in the former Soviet Union is attributable in large part to a decline in the sexual activity rate, rather than other more commonly mentioned factors, particularly economic and social factors leading couples to want fewer children. The URHS included several questions on the sexual experience of the respondents which helped to examine whether there might be any validity to this hypothesis. There are no comparison data from before the fertility decline, and the URHS did not collect information on trends in sexual activity, but the data available will provide some indication of whether sexual activity rates are at low levels or not.

Figure 10.1 displays the percent of 15-24 year-old respondents, by single year of age, who reported that they had ever had sexual intercourse. Relatively few females under the age of 17 reported being sexually experienced. Table 10.1 shows the proportion of respondents in each five year age cohort (except for 15-19 year-olds) who reported having sexual intercourse before selected ages. The results reveal that the proportion of women who have had sexual experience by the time they turn 16, 18 and 20 years old has increased in recent years, indicating that the age at first sex has been declining substantially. Twenty-three percent of respondents in the 20-24 year old cohort had their first sexual experience before age 16, about four times the percentage in the three oldest cohorts. The proportion of 20-24 year-olds who were sexually experienced by age 18 was 54%, also more than twice as high as for the older cohorts. The percentage of women who had intercourse by age 20 also increased, from 51% to 85%. Overall, urban women were slightly more likely than rural women to become sexually active by ages 16 or 18. The median age at first sexual intercourse (i.e., the age by which one-half of each cohort became sexually active) also reveals that sexual initiation has been becoming younger in recent years. The median age declined from 19.6 years for the 30-34 year-old cohort to 17.8 years for the 20-24 year cohort, representing a decline of almost two years in a ten year period. Figures were very similar for urban and rural Ukraine. These figures provide strong evidence that the age at which Ukrainian females have been becoming sexually active has been declining rapidly in the previous 10 years or so. The increase in sexual experience by ages 16, 18, and 20 can be seen graphically in Figure 10.2.

About three-fourths of all sexually experienced respondents reported having sexual intercourse in the previous thirty days (Table 10.2). Sixty-three percent had sexual relations in the previous week. When these tabulations are restricted to women currently in union, the proportion sexually active was 72% for the past week, 83% for the past month, and 95% for the past year. There was virtually no difference between urban and rural areas in reported sexual activity. Small differences were noted between the country's regions, with rates for the past week and month lowest in the South and highest in the Central region. Not surprisingly, there were differences according to the age of respondents. Weekly and monthly rates were highest among women between the ages of 20 and 34 years. It is assumed that the rates are slightly lower among 15-19 year-olds because a large proportion of the sexually experienced females at those ages are not currently in union. Among sexually experienced women who were never in union, who were primarily young women, 63% were sexually active in the previous 30 days.

Table 10.3 presents the percent distribution of the number of times sexually experienced respondents reported having sexual intercourse in the previous 30 days. The proportion of women in union who did not give any response or said that they did not remember, was almost one in four (24%). Of the 76% of women in union at the time of interview who responded to the question on coital frequency, four of every ten reported having intercourse at least 10 times in the previous 30 days. About one-third of that group reported having intercourse 20 or more times in the previous 30 days. About one of every three responding women in union reported having sexual relations from 3 to 9 times during the month. Ten percent of women in union said that they did not have sexual relations in the previous 30 days. The overall median for women in union was slightly over five times per month. As expected, coital frequency varied with women's ages; the proportion of women in union reporting intercourse at least 10 times in the previous 30 days declined from 38% for 15-24 year-olds to 22% for 35-44 year-olds. On the other hand, the proportion sexually inactive in the previous month varied relatively little with age, though it was slightly higher for 15-24 year-

olds and 35-44 year-olds than for 25-34 year-old women. The URHS does not provide information on trends in coital frequency.

The bottom panel of Table 10.3 examines coital frequency among sexually experienced women not in union at the time of interview. Over half of these women reported having no sexual intercourse in the previous 30 days. The sexual activity of these women was highly correlated with their age. Among the oldest cohort of women the percentage reporting no recent intercourse was almost twice as high as among the youngest women (65% vs. 39%). Based on the data showing an increase in sexual experience by age 16, 18 and 20, and the high proportion of women who said they were currently sexually active with high median coital frequency, even without the comparison to earlier years, it seems unlikely that the reported frequency of sexual intercourse is low enough to be responsible for significant declines in the pregnancy rate observed recently.

Information on the number of recent sexual partners women have had is useful in examining risks for sexually transmitted diseases as well as the analysis of sexual behavior with regard to women's health. The vast majority of sexually experienced women (92%) said that they had had no more than one sexual partner in the previous twelve months (Table 10.4). Only about 6% of women reported having multiple partners during that time. The likelihood of multiple sexual partners was highest among 15-19 year-old women (20%), never married women (27%), and residents of urban areas (7%). Women in the West region were much less likely than others (3%) to report having more than one partner in the previous 12 months. If these data are reliable, they indicate that few married Ukrainian women have more than one sexual partner over the course of a year. However, because this behavior is so stigmatized, it is not possible to say whether significant numbers of women underreport the number of partners they have had. The data also indicate that the proportion of women having large numbers of sexual partners is exceedingly small.

First sexual experience of young adults

The median age at first sexual intercourse, based on reports of whether young survey respondents had ever had sexual intercourse, was about 18.4 years. The proportion of 15-17 year-olds reporting that they had ever had intercourse was still relatively low, 18% (Table 10.5). However, many girls became sexually active at ages 18 or 19. More than half of 18-19 year-olds were sexually experienced, as were about four of every five 20-21 year-olds. The proportion sexually experienced reached its highest point at ages 22- 24, at 91 percent.

Clearly, premarital sexual intercourse has become common throughout Ukraine. About 30% of 15-19 year-old females reported having premarital intercourse, while this figure was 73% for females ages 20 to 24 years old. Altogether 60% of all 15-24 year-old respondents had had sexual intercourse before marriage. Eighty-five percent of all sexually experienced young women reported having premarital sexual intercourse. The proportion of sexually experienced young adults who had first intercourse before marriage, was inversely related to age at first intercourse, declining from 96% of 15-17 years olds to 81% of 22-24 years olds.

At ages younger than 18 years, urban females were slightly more likely than rural females to be sexually experienced, but the difference was not significant (Table 10.6). However, even this small urban-rural difference disappears after age 17. Sexually experienced urban women at all ages were

more likely than sexually experienced rural women to have had premarital intercourse.

Eighty-five percent of initial sexual experience for young women in Ukraine was premarital; only 15% of sexually experienced young women reported that their first sexual partner was their husband (Table 10.7). About equal numbers of women said that their first sexual partner was a fiancé or boyfriend (each 36%). An additional 9% of women said they first had sexual relations with a friend or acquaintance. First intercourse with a boyfriend or friend tended to be more common when it took place before age 18. First intercourse was described as forced intercourse or rape by 2% of young women. Even among those who first had sexual relations after their eighteenth birthday, for only 23% was first intercourse with their husband. Rural women were far more likely than urban women to report that their first sexual experience was after marriage or with a fiancé (64% vs. 46%).

Just under half of young women who first had intercourse before marriage (47%) reported that they or their partner used contraception during her first sexual experience (Table 10.8). Condoms accounted for over half of this contraception (27%), with withdrawal the only other commonly used method (13%). No other method was used by more than 2% of respondents. The predominance of condoms and withdrawal is not surprising, given that few other methods are readily accessible and do not require contact with a clinic or physician. Those females who first had premarital sex at age 18 or older were not significantly more likely than those starting at younger ages to report using contraception at first intercourse (48% vs. 45%). However, those who began sexual activity at age 18 or later were more likely than those beginning younger to report using withdrawal at their first premarital sexual experience. Females whose first sexual experience was at a younger age were slightly more likely to report using a condom at their first premarital sex. The relatively widespread use of condoms by the youngest women/couples at first premarital intercourse may indicate some success in the promotion of prevention strategies not only of unintended pregnancy, but sexual transmitted diseases as well.

Contraceptive use at first intercourse varied by residence. Overall, rural women/couples were only about two-thirds as likely as urban residents to use contraception during their first premarital intercourse (35% vs. 50%). There was an even greater difference in condom use, which was two to three times higher in urban areas than in rural areas, regardless of women's ages.

Table 10.9 displays data on premarital sexual experience and use of contraception for Eastern European countries in which similar reproductive health surveys have been conducted since 1993. Data from the USA is also shown for comparison purposes. The proportion of young women reporting premarital sexual experience in Ukraine was the third highest of the six countries, following the Czech Republic and Russia. Ukraine ranked fourth in overall contraceptive use, but had the second highest use of modern methods. All countries in the region registered contraceptive use well below that reported in the United States.

Figure 10.1
Percent of Respondents Who Have Ever Had
Sexual Intercourse, by Current Age
1999 Ukraine Reproductive Health Survey




Figure 10.2
Percent of Respondents Who First Had
Sexual Intercourse Before Selected Ages, by Current Age
1999 Ukraine Reproductive Health Survey




Table 10.1
 Percent of women 20-44 years of age who had sexual intercourse before ages 16, 18, 20,
 and median age at first sexual intercourse, by current age, by residence
 1999 Ukraine Reproductive Health Survey

Intercourse before:	Current age of respondent					Total
	20-24	25-29	30-34	35-39	40-44	
<u>Total</u>						
Age 16	22.5	9.1	4.3	6.1	6.0	9.6
Age 18	53.9	34.4	20.6	19.6	17.2	29.1
Age 20	84.7	70.4	57.4	53.1	50.8	63.2
Median age at first intercourse	17.8	18.8	19.6	19.8	20.0	19.2
<i>Number of women</i>	1,182	1,227	1,195	1,246	1,178	6,028
<u>Urban</u>						
Age 16	23.7	10.5	4.8	6.4	6.7	10.4
Age 18	54.9	37.3	21.2	19.8	17.6	30.1
Age 20	86.1	70.1	56.5	51.2	50.3	62.7
Median age at first intercourse	17.7	18.8	19.6	19.9	20.0	19.2
<i>Number of women</i>	919	941	908	990	925	4,683
<u>Rural</u>						
Age 16	19.1	5.7	3.2	5.3	4.2	7.3
Age 18	51.2	27.2	19.3	19.1	16.2	26.4
Age 20	80.9	70.9	59.6	58.4	52.3	64.4
Median age at first intercourse	17.9	19.0	19.6	19.6	20.1	19.2
<i>Number of women</i>	263	286	287	256	253	1,345

Table 10.2
 Cumulative percentage of respondents 15-44 years of age who reported having sexual intercourse within selected periods
 of time, by residence, region, age, and marital status, sexually experienced women
 1999 Ukraine Reproductive Health Survey

Characteristic	Percent with intercourse in:			<i>Number of women</i>
	Last week	Last month	Last year	
Total	63.4	74.9	90.9	5,646
<u>Residence</u>				
Urban	63.6	74.1	90.7	4,370
Rural	63.0	76.9	91.2	1,276
<u>Region</u>				
North	62.4	76.1	93.0	688
Central	67.5	79.7	92.6	533
East	65.7	75.8	90.4	2,068
South	58.8	69.9	91.1	1,486
West	62.0	73.8	88.7	871
<u>Age</u>				
15-19	54.9	67.6	95.9	315
20-24	62.9	73.5	93.9	968
25-29	67.9	78.1	93.0	1,108
30-34	70.0	80.2	92.9	1,096
35-39	63.3	71.4	89.4	1,101
40-44	55.4	70.5	83.7	1,058
<u>Marital status</u>				
Currently in union	71.9	83.1	95.2	4,380
Previously in union	29.9	40.4	67.1	736
Never in union	40.1	54.6	88.0	530

*342 women (5.7%) who did not respond to questions on most recent sexual intercourse are not included in this table

Table 10.3
 Percentage distribution of frequency of sexual intercourse in the previous 30 days,
 by age, by current marital status, among sexually experienced women
 1999 Ukraine Reproductive Health Survey

Characteristic	Number of times had sexual intercourse in the last 30 days							Total	Number of women
	0	1-2	3-4	5-9	10-19	20+	Don't remember/ No response		
<u>Currently in union</u>									
15-24	11.6	6.6	7.3	11.7	21.3	16.4	25.2	100.0	841
25-34	7.0	9.0	8.9	15.4	23.6	10.4	25.8	100.0	1946
35-44	11.1	14.4	13.4	18.0	16.3	5.7	20.9	100.0	1855
Total	9.5	10.7	10.4	15.8	20.2	9.6	23.8	100.0	4,632
<u>Not currently in union</u>									
15-24	38.9	15.6	6.8	8.4	8.9	6.0	15.3	100.0	521
25-34	49.7	12.9	6.9	7.8	8.5	4.8	9.1	100.0	376
35-44	64.9	10.8	7.3	6.2	4.2	0.9	5.8	100.0	459
Total	50.4	13.1	7.0	7.5	7.6	4.0	10.4	100.0	1,356

Table 10.4
 Percentage distribution of numbers of sexual partners in the previous 12 months,
 by age, region, residence, and marital status, sexually experienced women 15-44 years of age
 1999 Ukraine Reproductive Health Survey

Characteristic	Number of sexual partners in last 12 months						Total	Women
	0	1	2	3 - 4	5+	DR		
Total	0.7	91.5	4.4	1.5	0.4	1.5	100.0	5,564
<u>Age</u>								
15-19	0.0	78.2	13.3	4.9	1.6	1.1	100.0	327
20-24	0.4	87.9	7.2	2.4	0.4	1.7	100.0	971
25-29	0.3	92.8	4.0	1.8	0.2	0.9	100.0	1,111
30-34	0.6	94.1	3.5	0.3	0.4	1.2	100.0	1,097
35-39	0.7	94.2	2.3	1.2	0.2	1.4	100.0	1,079
40-44	1.8	92.5	2.5	0.6	0.4	2.2	100.0	997
<u>Region</u>								
North	0.3	91.0	4.5	2.6	0.4	1.2	100.0	673
Central	1.6	93.2	4.1	0.4	0.0	0.8	100.0	510
East	0.7	89.7	5.6	1.7	0.4	1.9	100.0	2,056
South	0.7	89.6	5.5	1.9	0.7	1.6	100.0	1,450
West	0.5	95.6	1.8	0.5	0.3	1.4	100.0	875
<u>Residence</u>								
Urban	0.8	90.4	5.0	1.9	0.3	1.7	100.0	4,297
Rural	0.5	94.4	3.0	0.5	0.7	1.0	100.0	1,267
<u>Marital status</u>								
Currently in union	0.1	97.2	1.5	0.5	0.2	0.5	100.0	4,495
Previously in union	5.0	68.1	15.0	4.5	0.4	6.3	100.0	558
Never in union	1.5	67.6	18.5	6.4	1.8	4.3	100.0	511

*Excludes 424 women (7.1%) who did not respond to the question.

*Table 10.5
Percent of respondents between the ages of 15 and 24
who have ever had sexual intercourse, by current age, by marital status at the time of intercourse
1999 Ukraine Reproductive Health Survey*

Current age	Ever had intercourse	First intercourse marital	First intercourse premarital	% Premarital	Number of women
15 - 17	17.8	0.7	17.0	95.9	657
18 - 19	54.5	5.4	49.1	90.0	422
20 - 21	82.4	11.6	70.8	85.9	437
22 - 24	90.9	17.0	73.8	81.2	714
15-19	32.1	2.6	29.6	92.0	1,079
20-24	87.7	14.9	72.7	82.9	1,151
15-24	59.6	8.7	50.9	85.4	2,230

**52 women (2.4%) who did not respond to question are not included in these tabulations.*

Table 10.6
Percent of respondents between the ages of 15 and 24
who have ever had sexual intercourse, by current age and residence, by marital status at the time of intercourse
1999 Ukraine Reproductive Health Survey

<i>Current age</i>	<i>Urban</i>					<i>Rural</i>				
	<i>% Ever had intercourse</i>	<i>First intercourse marital</i>	<i>First intercourse premarital</i>	<i>% Premarital</i>	<i>N</i>	<i>% Ever had intercourse</i>	<i>First intercourse marital</i>	<i>First intercourse premarital</i>	<i>% Premarital</i>	<i>N</i>
15 - 17	18.7	0.5	18.2	97.2	513	15.2	1.4	13.8	91.0	144
18 - 19	54.7	4.4	50.3	91.9	329	53.9	8.4	45.5	84.5	93
20 - 21	80.3	10.6	69.7	86.8	342	88.6	14.5	74.1	83.7	95
22 - 24	91.1	14.3	76.8	84.5	550	89.9	24.2	65.7	73.1	164
15-19	32.7	2.0	30.7	93.8	842	30.4	4.1	26.3	86.5	237
20-24	86.9	12.8	74.1	85.3	892	89.5	20.7	68.8	76.9	259
15-24	59.5	7.4	52.1	87.6	1,734	60.0	12.4	47.6	79.3	496

*52 women (2.4%) who did not respond to question are not included in these tabulations.

Table 10.7
 Percentage distribution of relationship to first sexual partner, by age at first sex, by residence
 15-24 year-old sexually experienced women
 1999 Ukraine Reproductive Health Survey

Relationship to first sexual partner	Residence/Age at first sex								
	Total			Urban			Rural		
	Total	<18	18+	Total	<18	18+	Total	<18	18+
Husband (after marriage)	14.5	8.3	22.7	12.3	6.6	20.4	20.5	13.7	28.4
Fiance	36.6	34.8	38.9	33.9	32.6	35.9	43.7	41.3	46.5
Boyfriend	35.9	40.5	29.9	39.6	44.1	33.3	25.7	29.5	21.3
Friend / acquaintance	8.9	11.4	5.7	9.5	11.4	6.7	7.6	11.5	3.1
Just met	1.0	0.9	1.1	1.3	1.1	1.6	0.4	0.7	0.0
Forced intercourse / rape	1.9	3.1	0.4	1.9	2.9	0.6	1.8	3.3	0.0
Family member / incest	0.1	0.2	0.0	0.1	0.2	0.0	0.0	0.0	0.0
Other	0.1	0.0	0.2	0.1	0.0	0.3	0.0	0.0	0.0
No response	0.9	0.8	1.0	1.1	1.1	1.1	0.3	0.0	0.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>1,336*</i>	<i>765</i>	<i>571</i>	<i>1,038</i>	<i>602</i>	<i>436</i>	<i>298</i>	<i>163</i>	<i>135</i>

*Total does not include 27 (2.0%) respondents who said they did not remember their age at first sexual intercourse.

Table 10.8
 Percentage distribution of contraceptive method used at first sexual intercourse,
 by age at first sex, by residence, 15-24 year-old women with premarital sexual experience
 1999 Ukraine Reproductive Health Survey

Contraceptive method	Residence/Age at first sex								
	Total			Urban			Rural		
	Total	<18	18+	Total	<18	18+	Total	<18	18+
Any method	46.5	45.1	48.4	50.0	45.5	50.9	35.4	30.9	41.9
Condoms	27.4	29.5	24.1	33.1	33.9	29.0	13.0	14.9	10.3
Withdrawal	12.9	10.6	16.6	11.8	9.9	14.9	16.3	12.6	21.6
Safe period	2.3	1.8	3.1	2.1	2.1	2.2	2.8	0.8	5.6
Pills	1.0	0.8	1.2	1.1	1.0	1.2	0.6	0.2	1.1
IUD	0.3	0.4	0.1	0.4	0.5	0.2	0.0	0.0	0.0
Vaginal methods	0.1	0.0	0.4	0.2	0.0	0.5	0.0	0.0	0.0
Other	0.2	0.3	0.0	0.3	0.5	0.0	0.0	0.0	0.0
Don't remember	2.3	1.7	3.0	2.1	1.6	2.9	2.8	2.4	3.3
No method	53.5	54.9	51.6	50.0	50.5	49.1	64.6	69.1	58.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of respondents</i>	<i>1,110</i>	<i>685</i>	<i>425</i>	<i>881</i>	<i>548</i>	<i>333</i>	<i>229</i>	<i>137</i>	<i>92</i>

*Total does not include 25 (2.4%) respondents who did not response

Table 10.9
 Percentage of young adult women 15-24 years of age reporting premarital sexual experience (PSE) and contraceptive use at first PSE
 Reproductive Health Surveys (RHS), Eastern Europe: 1993 -1999 and United States, 1995

Country	Year of survey	% reporting PSE		% 15-24 using contraception		
		15-19	20-24	Total	Modern methods	Traditional methods
Czech Republic	1993	36	93	57	28	29
Moldova	1997	14	40	33	14	19
Romania	1999	24	58	58	28	30
Russia*	1999	49	87	51	40	10
Ukraine	1999	30	73	47	32	15
Georgia	1999	**	2	3	3	0
USA	1995			76	70	4

*Three sites only: Ivanovo Oblast, Ekaterinburg city, and Perm city

**Less than 1%

CHAPTER XI

MATERNAL AND CHILD HEALTH / WOMEN'S HEALTH

The URHS questionnaire included a considerable amount of information on maternal and child health topics. With regard to health, there is great interest not only in the outcomes of pregnancies and complications associated with induced abortion, as discussed in previous chapters, but also with attitudes and practices of women and health care providers, and with facility practices that can have an impact on pregnancy outcomes and infant health and well-being. Each respondent who had given birth since the beginning of 1994 was asked a series of questions regarding her most recent pregnancies and deliveries, including such topics as prenatal care, utilization of various health services, breastfeeding, and cigarette smoking.

Prenatal care

Prenatal care is most beneficial when it begins early and continues regularly throughout a pregnancy. According to the Prenatal Care Program guidelines of the Ukraine Ministry of Health (MOH), the first prenatal care visit should occur before the thirteenth week of pregnancy. To evaluate the adequacy of prenatal care, it is necessary to monitor not only the timing of the initial visit, but also the total number of prenatal care visits during pregnancy. According to the prenatal care program implemented in 1995 by the MOH, healthy pregnant women should make one prenatal visit per month during first twenty weeks of pregnancy, two visits per months during the next ten weeks, and weekly prenatal visits during the final ten weeks.

About 10% of survey respondents reported that they received no prenatal care during their last pregnancy (since the beginning of 1994) leading to a live birth (Table 11.1 and Figure 11.1). Another 3% waited until the third trimester of pregnancy before they had any prenatal care. In about two-thirds of pregnancies (65%), prenatal care began during the first trimester, as recommended. Residents of the South region, those who were over 35 years old, those who had more than two children, and those who did not go beyond secondary education were the most likely to receive no prenatal care and were more likely to initiate it later. The likelihood of receiving early prenatal care was highest among urban women, residents of the Central region, the youngest women, those who had no children, and those with highest level of education.

Among women who received prenatal care, 81% made at least ten prenatal care visits (Table 11.2). About 10% of women made between five and nine visits, and only 7% of women made fewer than five visits during the last pregnancy resulting in a live birth since January 1994. Urban women made an adequate number of prenatal visits more often than rural women. However, in spite of the tendency to start prenatal care later, women over 35 years of age were at least as likely as younger women to make 10 or more visits.

Government facilities were the principal source for prenatal care among pregnant women with a live birth since January 1994, regardless of their characteristics (Table 11.3). Two-thirds of women received their prenatal care (or most of it) at women consultation clinics. However, rural women and residents of the Central region were somewhat less likely than others to use these facilities. These women received prenatal care at hospitals and maternity houses more often than other

Centers. Private clinics provided prenatal care for very few women (0.2%), mainly from the East region.

Fetal ultrasound is an important diagnostic procedure in obstetrics, which is valuable for: assessing gestational age, diagnosis of multiple pregnancy, malpresentations, hydatidiform moles, location of the placental site, and the detection of fetal abnormalities. Seventy-eight percent of women with recent live births had a diagnostic ultrasound during pregnancy (Table 11.4). Urban women (81%), Central region residents (82%), the youngest women (81%), and the most educated women (84%) reported the highest use of ultrasound.

Elevated blood pressure (BP) during pregnancy can be a sign of serious disease or pregnancy complications. Measurement of blood pressure should be done during every prenatal visit, in order to reduce the risk of adverse pregnancy and birth outcomes. Eighty-seven percent of women said that they had their BP measured during their pregnancy (Table 11.4, column 2). Differentials in BP measurement according to residence, age and education were very small. In addition to the 10% of women who received no prenatal care, 3% of women did not have their BP measured by their prenatal care provider. Urban women were more than three times more likely than rural women not to have their BP measured during their prenatal care visits. It is encouraging that BP was measured among all women over 35 years of age who visited prenatal care facilities.

Overall, 16% of women were told that they had high blood pressure during pregnancy (Table 10.5). Differences between population subgroups in reported high blood pressure rates tended to be very small. Thirty-four percent of women were told that they had anemia. This condition was reported most often by urban women, those who were younger than 35 years of age, and those with the highest level of education. Thirty-eight percent of women with recent births said they took iron supplements during pregnancy. Although women were not asked whether they took supplements as a preventive measure or because they had been diagnosed with anemia, it is likely that many of these women took supplements because of an anemia diagnosis. Iron supplements were taken most frequently by urban women, 15-24 year-olds, and the best educated.

Hospitalization during pregnancy

Through analysis of the frequency and length of the hospitalization of women during recent pregnancies (for reasons other than delivery) we tried to determine the likelihood and seriousness of health problems and pregnancy complications among Ukrainian women during pregnancy. Ukrainian physicians, like most of their eastern European counterparts, hospitalize women for pregnancy complications more readily and for longer periods of time than physicians in other industrialized countries. Furthermore, hospital stays for delivery tend to be longer than elsewhere. Because of the changing economic situation in Ukraine and increasing influence of medical practice from other areas it would not be surprising to observe decreases in the likelihood and length of hospitalization associated with pregnancy complications and other health problems.

URHS respondents were asked whether and, if so, for how long they were hospitalized because of pregnancy-associated problems prior to their most recent delivery in the previous five years. Thirty-two percent of women with deliveries leading to a live birth since January 1994 reported being hospitalized for prenatal problems (Table 11.6). There were considerable differences in the likelihood of hospitalization according to respondent characteristics. The probability of

hospitalization was inversely correlated with birth order, from 38% for first births to only 14% for third or higher order births. Urban women were more likely to be hospitalized than rural women. Hospitalization also decreased with increasing age, which is strongly correlated with birth order. The proportion hospitalized was highest in the East and lowest in the West. In spite of economic changes and on-going health sector reform, the proportion of women hospitalized was virtually unchanged between 1994-96 and 1997-99.

As anticipated, the duration of hospital stays tended to be very long (Table 11.7 and Figure 11.2). The median stay was between two and three weeks and two-thirds lasted for two weeks or more. Only 7% of stays were reported to be shorter than one week. Urban women and women from the North region tended to have longer hospital stays than other women. The duration of hospitalization appears to be little changed in recent years.

Labor and delivery

Typically labor and delivery in Ukraine takes place in government obstetric facilities such as maternity houses, obstetric departments of MCH centers and regional hospitals, where trained personnel can provide women with specialized obstetric care. The top panel of Table 11.8 shows that 94% of deliveries leading to live births since 1994 took place at maternity houses. Most of the remainder (5%) occurred in MCH centers and hospitals. Fewer than one percent of deliveries took place at home.

Fifty-eight percent of women with recent live births reported receiving a post partum check-up within six weeks of delivery of their last baby (Table 11.9). The proportion receiving check-ups appears to have decreased somewhat between 1994-96 and 1997-99, from 60% to 55%. Rural women (54%), as well as those from the South (53%) and West (54%), were less likely than others to have received check-ups. The likelihood of having an exam decreased substantially with increasing age at delivery and increasing birth order. Most noteworthy is the apparent decline in proportions of women undergoing post partum exams in recent years. These declines have been especially marked in rural areas and in western Ukraine

Cigarette smoking during pregnancy

There is strong evidence that tobacco use affects not only the health of women, but also increases the risk of adverse maternal and perinatal outcomes as well. Overall, 9% of women were cigarette smokers at the time they became pregnant with their most recent pregnancy leading to a live birth since January 1994 (Table 11.10). Urban women were three times more likely than rural women to smoke. About half of women who were cigarette smokers when they became pregnant reported that they stopped smoking during their pregnancy. The proportion of those who gave up smoking during pregnancy was about the same among urban and rural women. Better educated women were more likely than less educated women to quit smoking. It has been well documented that smoking during pregnancy increases the risk that an infant will be born with low birthweight. Respondents who delivered low birthweight infants (i.e., under 2500 grams) were almost twice as likely as women whose babies were normal birthweight to be smokers at the beginning of pregnancy and throughout the pregnancy versus 4%, respectively).

Breastfeeding

Breastfeeding practices have a significant influence on the health of the child, as well as the fertility of the mother. Breast milk not only provides complete nutrition for infants. It also contains antibodies that protect infants from infection before their immune system is fully mature, thus decreasing infant morbidity and mortality. An additional effect of breastfeeding is its suppression of ovulation following a birth, thus reducing the risk of pregnancy. In the URHS women were asked whether each of her recently born children had been breastfed, were currently being breastfed, and, if currently breastfed, whether they had started receiving other foods or liquids yet.

Breastfeeding remains the norm across Ukraine. According to results of the URHS, 92% of babies born to respondents since January 1994 reportedly had been breastfed (Table 11.11). The percentage of children who were ever breastfed did not differ greatly across residence, age, and education categories of the population, though the proportion breastfed was somewhat lower in the Central region (86%). The percentage breastfed has not changed significantly in recent years; the proportions were very similar for babies born in 1994-1995, 1996-1997, and 1998-1999. The mean duration of breastfeeding for those babies who were breastfed was 7.5 months. The percentage was slightly higher in the Central region (9.3 months) and lower in the North region (5.9 months) than elsewhere. Otherwise, the differentials observed were small.

In Table 11.12, which examines current breastfeeding status, infants have been split into three age categories: under 4 months, 4-5 months, and 6-11 months. Almost eight of every ten infants under four months of age and just over half of those four to five months of age were still being breastfed. These figures drop off sharply for 6-11 month-old infants, among whom one-third were still being breastfed. Since some of the health benefits of breastfeeding may be diminished by the early introduction of other foods and liquids, it is important to examine the extent of exclusive breastfeeding. An infant was considered to be exclusively breastfed if he/she receives only breast milk. UNICEF recommends that infants in the youngest age group (i.e., under four months) be exclusively breastfed, while the oldest group should be receiving breast milk as well as other foods. Although the prevalence of breastfeeding was high, for many young infants breastfeeding was only partial. Among infants under 4 months of age, only 31% were exclusively breastfed. Less than 1% of infants at least six months old were exclusively breastfed, while almost 70% were not being breastfed at all.

Figure 11.3 displays information on an assortment of indicators, both behaviors and circumstances that are negatively related to infant and/or maternal health: the proportions of women with live births since the beginning of 1994 who received no prenatal care during their first trimester, had fewer than 10 prenatal care visits, were hospitalized during pregnancy, smoked cigarettes during pregnancy, and did not breastfeed their infant.

Women's health behaviors

The URHS also included some questions that addressed issues of women's health. Two of these topics were cigarette smoking and general gynecologic check-ups.

Tobacco use is thought to have increased sharply throughout eastern Europe and the former Soviet Union for approximately the past decade. Nineteen percent of respondents said that they currently smoked cigarettes (Table 11.13). The most marked difference in smoking prevalence was

according to place of residence: urban women were three times more likely than rural women to be smokers (23% versus 7%). Smoking was most common among women at ages 20-24 (25%) and decreased with increasing age, such that it was lowest among the oldest respondents (14%). The fact that the proportions were lowest in the thirties and forties and that even the rates for teenagers exceeded those for the oldest respondents probably indicates that the prevalence of smoking has been increasing in recent years. There was only a weak correlation between smoking and level of education noted, with the least educated most likely to smoke. It is somewhat encouraging to note that few of the current smokers in the URHS could be considered heavy smokers, with only 3% of women reporting that they typically smoked more than ten cigarettes per day. A very small percentage of women (0.1%) reported that they smoked more than a whole pack of cigarettes per day.

It is recommended that women of childbearing age undergo a routine (i.e., not pregnancy related) gynecologic examination at least once per year. In fact, a majority of respondents (62%) said that they had had an exam during the previous 12 months (Table 11.14). However, a substantial number of women (11%) had never undergone such an exam. For very few women was the most recent exam more than four years previously. As expected, the distribution of years since the most recent gynecologic exam and whether women had ever had an exam were highly correlated with the age and marital status of respondents. The proportion without a recent exam was especially high among youngest women. The proportion without a recent exam was also higher among rural women compared to urban women, and was higher in the South and West regions than in the other regions.

Figure 11.1
Percentage Distribution of Time of First Prenatal Visit
for the Last Pregnancy Since January 1994
1999 Ukraine Reproductive Health Survey




Figure 11.2
Percentage Distribution of Length of Hospitalization
During Last Pregnancy Since January 1994
1999 Ukraine Reproductive Health Survey




Figure 11.3
Percent of Respondents Reporting Selected Behaviors/Circumstances
Related to the Health of Infants/Mothers for Last Pregnancies
Leading to Live Births Since January 1994
1999 Ukraine Reproductive Health Survey




Table 11.1
Percentage distribution of trimester of first prenatal care visit, by selected characteristics,
all pregnancies ending in live birth since January 1994
1999 Ukraine Reproductive Health Survey

Characteristic	Start of prenatal care					Total	Number of pregnancies
	1st trimester	2nd trimester	3rd trimester	No prenatal care	Do not remember		
<u>Total</u>	65.2	20.0	2.5	9.8	2.5	100.0	1,938
<u>Residence</u>							
Urban	66.6	18.2	2.9	9.8	2.5	100.0	1,389
Rural	62.5	23.4	1.7	9.7	2.7	100.0	549
<u>Region</u>							
North	67.9	16.9	1.7	11.3	2.2	100.0	228
Central	69.4	22.9	2.9	2.9	1.9	100.0	170
East	64.7	18.3	2.4	11.7	2.8	100.0	617
South	57.9	19.0	4.2	15.7	3.2	100.0	463
West	66.5	22.5	1.9	6.7	2.4	100.0	460
<u>Age at delivery</u>							
15-24	65.8	19.7	2.2	9.3	3.0	100.0	1,147
25-34	64.4	21.0	2.6	10.0	2.0	100.0	683
35-44	59.8	18.5	5.9	14.6	1.2	100.0	98
<u>Birth order</u>							
1	67.4	18.8	2.1	8.9	2.8	100.0	1,149
2	64.1	20.2	2.5	10.8	2.4	100.0	584
3 +	55.3	26.9	3.8	12.1	1.9	100.0	205
<u>Education</u>							
<Complete secondary	64.7	20.6	2.7	11.0	1.0	100.0	162
Complete secondary	63.6	20.7	2.5	10.4	2.8	100.0	1,395
>Complete secondary	71.3	16.9	2.4	6.9	2.5	100.0	381

Table 11.2
 Percentage distribution of number of prenatal care visits, by selected characteristics,
 all pregnancies with prenatal care ending in live births since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	Number of prenatal care visits				Total	Number of women
	1-4	5-9	10+	Don't remember		
<u>Total</u>	6.9	9.7	81.3	2.1	100.0	1,724
<u>Residence</u>						
Urban	6.0	8.8	83.4	1.7	100.0	1,269
Rural	8.8	11.4	77.2	2.7	100.0	455
<u>Regions</u>						
North	5.4	3.4	89.3	1.9	100.0	197
Central	5.5	6.7	86.7	1.2	100.0	153
East	8.1	10.6	78.3	3.0	100.0	572
South	5.2	9.3	83.5	1.9	100.0	427
West	8.1	13.2	77.0	1.6	100.0	375
<u>Age at delivery</u>						
15-24	6.6	10.4	80.9	2.1	100.0	938
25-34	7.5	9.5	80.9	2.1	100.0	684
35-44	9.1	2.2	87.2	1.5	100.0	97
<u>Birth Order</u>						
1	6.5	9.8	81.7	2.0	100.0	1,020
2	6.4	10.4	81.0	2.2	100.0	549
3 +	10.6	8.1	79.8	1.5	100.0	155
<u>Education</u>						
< Complete secondary	4.8	6.3	88.9	0.0	100.0	142
Complete secondary	7.2	10.1	80.2	2.5	100.0	1,223
> Complete secondary	7.2	9.6	82.0	1.2	100.0	359

Table 11.3
 Percentage distribution of place of (most) prenatal care visits, by residence and region,
 all pregnancies ending in live birth since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	Place of prenatal care visits							Total	Number of women
	Women's consult clinic	Hospital	Maternity house	MCH center	Private clinic	Other	No care		
<u>Total</u>	75.1	8.1	5.2	0.4	0.2	1.2	9.8	100.0	1,934*
<u>Residence</u>									
Urban	77.8	5.6	5.2	0.6	0.3	0.7	9.8	100.0	1,388
Rural	70.2	12.7	4.9	0.0	0.2	2.3	9.7	100.0	546
<u>Region</u>									
North	75.5	8.8	2.9	0.4	0.0	1.1	11.3	100.0	228
Central	62.7	21.9	10.7	0.0	0.0	1.8	2.9	100.0	169
East	77.6	5.2	4.8	0.1	0.6	0.0	11.7	100.0	617
South	73.4	2.3	7.0	0.8	0.0	0.8	15.7	100.0	461
West	78.2	8.5	3.9	0.4	0.2	2.1	6.7	100.0	459

* Does not include 4 women who did not give information on place of prenatal care.

Table 11.4
 Percent of respondents who had their blood pressure (BP) measured and percent who received an ultrasound during their most recent pregnancy ending in a live birth since January 1994, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	% who received ultrasound	% who had BP measured	Number of women
<u>Total</u>	78.2	86.9	1,938
<u>Residence</u>			
Urban	80.5	85.8	1,389
Rural	73.3	88.4	549
<u>Region</u>			
North	78.8	85.4	228
Central	81.5	94.1	170
East	79.2	84.2	617
South	77.2	81.7	463
West	76.0	89.7	460
<u>Age</u>			
15-24	81.3	87.9	1,147
25-34	77.0	85.9	683
35-44	72.6	85.3	98
<u>Education</u>			
< Complete secondary	71.0	84.3	162
Complete secondary	77.5	86.0	1,395
> Complete secondary	83.5	89.7	381

NOTE: Respondents who reported receiving no prenatal care are assumed not to have received the services listed.

Table 11.5
 Percent of respondents who were told they had high blood pressure (BP), percent who were told they had anemia,
 and percent who took iron supplements during their most recent pregnancy
 ending in a live birth since January 1994, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Percent told they had high BP	Percent told they had anemia	Percent who took iron supplements	<i>Number of women</i>
<u>Total</u>	16.3	33.8	38.3	1,938
<u>Residence</u>				
Urban	17.2	35.5	42.6	1,389
Rural	14.3	30.5	29.3	549
<u>Region</u>				
North	15.6	38.1	41.1	228
Central	19.7	36.8	38.1	170
East	17.1	34.2	38.6	617
South	14.6	33.6	39.1	463
West	15.0	30.2	35.9	460
<u>Age</u>				
15-24	14.9	34.1	41.4	1,147
25-34	16.8	34.2	37.4	683
35-44	18.5	30.9	31.2	98
<u>Education</u>				
< Comp. secondary	20.3	29.9	38.3	162
Complete secondary	16.7	33.1	36.7	1,395
> Comp. secondary	13.3	38.1	43.6	381

NOTES: Excludes 10 respondents who did not remember whether they received particular services. Respondents who reported receiving no prenatal care are included with the negative responses for all variables.

Table 11.6
 Percent of pregnancies ending in a live birth since January 1994
 during which women were hospitalized, by residence, region, and year of delivery
 1999 Ukraine Reproductive Health Survey

Characteristic	Percent hospitalized	Number of live births
<u>Total</u>	32.3	1,938
<u>Residence</u>		
Urban	34.8	1,389
Rural	27.5	549
<u>Region</u>		
North	34.8	228
Central	32.3	170
East	39.4	617
South	35.3	463
West	22.9	460
<u>Birth Order</u>		
1	37.9	1,149
2	27.8	584
3 +	13.5	205
<u>Age at delivery</u>		
15-24	35.8	1,147
25-34	30.1	683
35-44	23.0	98
<u>Education</u>		
<Complete Secondary	33.1	162
Comp Secondary	30.6	1,395
>Complete Secondary	32.8	381
<u>Year of delivery</u>		
1994-1996	32.8	1,046
1997-1999	31.8	881

Table 11.7
 Percentage distribution of length of hospitalization by residence, region, and year of delivery,
 pregnancies ending in a live birth since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	Length of hospitalization					Total	Number of hospitalizations
	1-6 nights	7-13 nights	14-29 nights	30+ nights	Don't remember		
<u>Total</u>	6.8	23.7	44.9	21.1	3.5	100.0	619
<u>Residence</u>							
Urban	6.4	24.1	43.5	23.6	2.5	100.0	463
Rural	7.8	22.8	48.4	15.2	5.9	100.0	156
<u>Region</u>							
North	6.4	14.1	50.0	25.6	3.9	100.0	78
Central	9.1	27.3	38.2	21.8	3.6	100.0	55
East	7.1	24.5	46.5	20.6	1.3	100.0	229
South	6.6	27.4	41.9	19.3	4.7	100.0	151
West	5.7	24.5	44.3	19.8	5.7	100.0	106
<u>Year of delivery</u>							
1994-1996	7.2	23.9	43.4	21.6	3.9	100.0	335
1997-1999	6.4	23.7	46.3	20.7	3.0	100.0	284

Table 11.8
 Percentage distribution of type of place of delivery for pregnancies ending in a live birth since January 1994, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Type of place of delivery						Total	Number of live births
	Maternity house	Hospital	MCH Center	Home	Other	Do not remember		
<u>Total</u>	93.8	4.5	0.8	0.6	0.2	0.2	100.0	1,938
<u>Residence</u>								
Urban	94.8	3.4	1.0	0.4	0.2	0.3	100.0	1,389
Rural	91.7	6.6	0.4	1.1	0.2	0.0	100.0	549
<u>Region</u>								
North	96.1	3.0	0.4	0.4	0.0	0.0	100.0	228
Central	91.8	6.5	0.0	1.2	0.6	0.0	100.0	170
East	95.7	2.4	1.4	0.5	0.0	0.0	100.0	617
South	95.6	0.9	1.3	1.3	0.0	0.1	100.0	463
West	90.6	8.3	0.4	0.2	0.4	0.0	100.0	460
<u>Age at delivery</u>								
15-24	94.8	4.2	0.6	0.2	0.0	0.2	100.0	1,147
25-34	93.8	4.8	0.8	0.7	0.2	0.2	100.0	683
35-44	92.6	3.5	1.6	1.6	0.7	0.0	100.0	98
<u>Birth order</u>								
1	94.6	4.1	0.8	0.0	0.3	0.2	100.0	1,149
2	94.0	4.1	1.0	0.7	0.0	0.2	100.0	584
3+	89.6	7.1	0.0	2.7	0.5	0.2	100.0	205
<u>Education</u>								
< Comp. secondary	94.4	3.9	0.0	1.8	0.0	0.0	100.0	162
Complete secondary	93.7	3.0	2.3	0.1	0.3	0.6	100.0	1,395
> Comp. secondary	93.7	4.9	0.5	0.6	0.2	0.1	100.0	381

Table 11.9
 Percentage of respondents who received a post partum check-up within six weeks of birth,
 by selected characteristics, by year of birth, for most recent birth since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	Year of birth		
	1994-1999	1994-1996	1997-1999
<u>Total</u>	57.6	60.1	55.0
<u>Residence</u>			
Urban	59.3	59.6	59.0
Rural	54.0	61.3	46.8
<u>Region</u>			
North	61.3	62.0	60.8
Central	62.1	68.8	55.3
East	60.1	58.3	62.2
South	52.5	56.6	48.8
West	54.2	59.8	48.4
<u>Age at delivery</u>			
15-24	60.5	64.1	56.7
25-34	55.0	55.1	54.8
35-44	43.0	48.3	38.7
<u>Birth order</u>			
1	61.5	65.5	57.6
2	54.0	53.7	54.4
3+	45.7	49.3	43.0
<u>Education</u>			
< Comp. secondary	60.2	63.5	58.0
Complete secondary	57.4	60.2	54.4
> Comp. secondary	57.3	58.7	55.6

NOTE: Excludes 14 women who did not provide information on whether they received a post partum check-up.

Table 11.10
 Percentage of women who smoked cigarettes at the time they became pregnant and percentage who continued smoking during pregnancy, by selected characteristics, most recent pregnancy ending in a live birth since January 1994
 1999 Ukraine Reproductive Health Survey

Characteristic	% who were smokers at time became pregnant	% who women who stopped for pregnancy	% of smokers who stopped for pregnancy	Number of women
<u>Total</u>	8.9	4.6	51.7	1,724
<u>Residence</u>				
Urban	11.5	5.9	51.3	1,269
Rural	3.7	2.1	56.8	455
<u>Age</u>				
15-24	9.3	5.4	58.1	938
25-34	7.9	3.5	44.3	689
35-44	12.3	5.5	44.7	97
<u>Education</u>				
< Complete secondary	10.7	3.5	32.7	142
Complete secondary	9.0	4.6	51.1	1,223
> Complete secondary	8.8	4.9	55.7	359
<u>Birth weight of baby</u>				
Less than 2500 grams	15.3	6.8	44.4	107
At least 2500 grams	8.6	4.4	51.2	1,617

Table 11.11
 Percent of babies born to respondents since January 1994 who were breastfed,
 and mean duration of breastfeeding for those ever breastfed*, by selected characteristics

Characteristic	Percent ever breastfed	Number of births since 1/94	Mean duration of breastfeeding (months)
<u>Total</u>	92.1	1,920	7.5
<u>Residence</u>			
Urban	92.4	1,188	7.6
Rural	91.2	732	7.2
<u>Region</u>			
North	93.9	225	5.9
Central	86.3	168	9.3
East	91.0	616	7.5
South	91.0	459	8.2
West	94.9	452	7.4
<u>Age of mother</u>			
15-24	93.1	683	7.2
25-34	91.8	1,049	7.6
35-44	89.8	188	**
<u>Year of birth</u>			
1994-95	91.7	672	NA
1996-97	91.8	741	NA
1998-99	93.0	507	NA
<u>Education of mother</u>			
<Complete secondary	89.1	159	6.3
Complete secondary	93.1	1,385	7.8
>Complete secondary	89.6	376	6.7

*Mean duration only for children who were ever breastfed, calculated using current status data.

**Too few recent births to calculate a stable estimate.

Table 11.12
 Percentage distribution of current breastfeeding status, by age
 for infants under one year of age at the time of interview
 1999 Ukraine Reproductive Health Survey

Residence/Age	Current breastfeeding status			Total	<i>Number of Babies</i>
	Exclusively breastfed	Partially breastfed	Not breastfed		
< 4 months	31.4	46.7	21.9	100.0	103
4-5 months	6.3	42.8	50.9	100.0	60
6-11 months	0.8	30.6	68.6	100.0	148

Table 11.13
 Percentage distribution of number of cigarettes typically smoked per day, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Cigarettes per day						Total	Number of women
	Do not smoke	< 1	1-10	11-20	>20	Don't know/ No answer		
<u>Total</u>	81.2	1.1	14.3	2.8	0.1	0.5	100.0	7,128
<u>Residence</u>								
Urban	76.8	1.4	17.6	3.6	0.2	0.5	100.0	5,544
Rural	92.9	0.4	5.5	0.7	0.0	0.5	100.0	1,584
<u>Age</u>								
15-19	81.1	1.2	15.9	1.2	0.0	0.5	100.0	1,100
20-24	75.1	1.4	20.0	2.9	0.1	0.4	100.0	1,182
25-29	78.3	1.1	16.4	3.7	0.1	0.5	100.0	1,227
30-34	82.3	1.0	12.7	3.6	0.2	0.3	100.0	1,195
35-39	83.7	0.8	12.2	2.7	0.0	0.6	100.0	1,246
40-44	86.4	1.2	8.9	2.6	0.3	0.9	100.0	1,178
<u>Education</u>								
< Complete secondary	77.5	1.0	16.8	3.6	0.2	0.9	100.0	778
Complete secondary	81.8	0.9	14.3	2.4	0.1	0.5	100.0	4,828
> Complete secondary	81.3	1.7	12.9	3.6	0.1	0.4	100.0	1,522

Table 11.14
 Percentage distribution of years since most recent (non pregnancy related) gynecologic exam, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Number of years since last gynecologic exam							DR	Total	Number of women
	Never	<1	1	2-3	4-9	10+				
<u>Total</u>	10.9	62.1	10.1	7.8	2.6	0.1	6.3	100.0	7,119*	
<u>Residence</u>										
Urban	9.9	62.9	9.7	7.9	2.6	0.1	6.9	100.0	5,536	
Rural	13.6	60.1	11.2	7.5	2.6	0.0	4.9	100.0	1,583	
<u>Region</u>										
North	7.3	67.8	11.0	7.5	1.6	0.1	4.8	100.0	858	
Central	9.1	70.2	8.0	7.0	1.8	0.0	3.8	100.0	616	
East	9.0	65.7	8.3	7.5	2.8	0.3	6.4	100.0	2666	
South	13.8	58.5	9.9	8.1	2.1	0.2	7.4	100.0	1822	
West	15.4	50.3	13.9	8.8	2.9	0.5	8.2	100.0	1157	
<u>Age</u>										
15-19	30.3	52.1	7.8	4.6	0.9	0.0	4.5	100.0	1,100	
20-24	12.4	65.8	10.0	4.7	1.6	0.0	5.4	100.0	1,182	
25-29	7.4	68.6	10.6	6.8	1.8	0.0	4.9	100.0	1,224	
30-34	5.4	65.9	11.5	9.5	2.2	0.1	5.3	100.0	1,195	
35-39	5.2	61.9	9.6	9.1	4.1	0.0	9.9	100.0	1,243	
40-44	5.1	58.5	11.2	12.2	4.9	0.3	7.9	100.0	1,175	
<u>Marital status</u>										
Currently in union	5.9	63.2	11.6	8.3	3.1	0.0	7.9	100.0	798	
Previously in union	6.9	65.1	10.1	8.6	2.8	0.1	6.4	100.0	4,789	
Never in union	24.9	53.1	9.5	5.3	1.7	0.1	5.4	100.0	1,532	

* Does not include 9 women, who did not respond whether they had had an exam.

CHAPTER XII

SEXUALLY TRANSMITTED INFECTIONS

Economic and social disruption often coincide with outbreaks or substantially increased incidence of adverse health conditions, especially infectious disease. It is now clear that the successor states of the former Soviet Union have been experiencing major epidemics of sexually transmitted infection (STI) (Renton et al., 1998; Waugh, 1999). For instance, the notification rate of new cases of syphilis in Ukraine increased from 6.0 per 100,000 population in 1990 to 144 per 100,000 in 1996 (Tichonova and Borisenko, 1997). Official statistics indicate that STI rates are even higher among younger people than in the overall population.

Even more alarming than the rapid rise in STI rates generally, has been the recent upsurge of HIV infection in the former Soviet Union, particularly among injecting drug users. There is already a major HIV epidemic in Ukraine and it has shown no sign of abating yet (Lifson and Preble, 1997; Barnett et al., 2000; Dehne et al., 1999). According to United Nations estimates, while there were 150,000 HIV infected individuals overall in western Europe by the end of 1997, in Ukraine alone there were as many as 180,000 people infected with the AIDS virus. Besides the obvious effects of this epidemic on the health of Ukraine, it also likely to have a dramatic effect social and economic impact on the nation.

Rates of sexually transmitted infection are largely determined by sexual behavior on the one hand and by the accessibility, acceptability, and effectiveness of services for early diagnosis and treatment on the other. Both are likely to have been radically altered by recent economic, political, and social changes that have generated huge income differentials and high rates of poverty, along with rising unemployment, especially among women. The opening of borders has increased migration both within the country and outside of the country. This migration, combined with dramatically increased communications and exposure to western media, has opened Ukraine and its neighbors to many new influences. These influences have contributed to changes in sexual behavior and sexual relationships, which in turn have impacted the prevalence of STIs, most notably in the young adult population.

The decentralization of medical services, along with other changes in society, has made it more difficult than previously to estimate the true size of the STI epidemic. During the Soviet period, the system of STI control in Ukraine was based on a centrally controlled nationwide clinical dermatovenerology service (DVS). The responsibilities of the DVS included free diagnosis and treatment, identification of sexual contacts for tracing, and partner notification. Diagnostic testing was also provided through active screening in clinically and occupationally defined groups. Diagnostic facilities included serological and bacteriological laboratories attached to clinics that were accredited and had high levels of quality control. The DVS structure has basically survived in Ukraine since independence. However, in addition, a large number of private dermatovenerology clinics and offices have been established, where patients pay for services and receive testing and treatment anonymously. For certain STIs, such as gonorrhea, patients are more likely to avoid government-provided services, for several reasons. The ability to remain anonymous attracts patients to these private clinics. Moreover, these clinics tend to be better equipped and provide higher quality service than public clinics. They can provide adequate testing and treatment for

those who have the ability to pay, while free-of-charge government clinics are sometimes unable to do so. As a result, higher quality services are available to the minority who can afford the fees, while lower quality services are provided to the majority. Individuals with STIs who wish to remain anonymous and have inadequate resources to pay for treatment in the private sector are often forced to try to treat themselves.

The combined effect of a number of factors, including the impact of political and market reforms, a decline in the standard of living for much of the population, a decline in the government's ability to fund health services, economic dislocation, a poorly controlled private medical sector, and the growing prevalence of self-medication, among others, has brought about an even greater need than in the past to emphasize primary prevention of STIs.

The STI module of the URHS was designed to help determine the population's knowledge about and experience with sexually transmitted infections/diseases and to identify the subgroups of the population with the greatest need for interventions to enhance primary prevention of STIs. These findings will also help to define factors which could influence the acceptability and effectiveness of primary prevention messages, in order to improve the likelihood of success of any educational efforts. The URHS included questions on awareness of STIs including HIV/AIDS, experience of specific STIs and conditions and symptoms often related to STIs, perceived risk of STIs, place of treatment for those reporting that they had been infected, and reasons for not seeking treatment. It should be kept in mind that, since these estimates are based on self-reports, there is a strong likelihood that the occurrence of STI's is underreported, owing to undiagnosed conditions and unwillingness to report their occurrence.

Awareness of STIs and lifetime history of STI diagnosis

The proportion of respondents who reported they had ever heard of selected sexually transmitted (or potentially sexually transmitted) conditions are displayed in Table 12.1. Knowledge of syphilis was nearly universal, with only 2% reporting they had never heard of it. The only other conditions known by a large majority of women were pelvic inflammatory disease (5% unaware), gonorrhea (9%), and genital ulcers (9%). The conditions with which the most respondents were unfamiliar were human papilloma virus (HPV) (74%), genital herpes (66%), and chlamydia (54%).

Awareness of STIs varied substantially according to respondent characteristics. Every condition asked about was much less widely known among rural women, 15-24 year-olds, those with incomplete secondary education, and the sexually inexperienced than among other groups of women. Also, women from West region were consistently less likely than residents of other regions to be aware of these conditions. The greatest differences tended to be for those conditions which were least widely known, such as chlamydia, genital herpes, and human papilloma virus.

A very high proportion of women reported having had pelvic inflammatory disease (PID) (38%) or genital ulcers (38%) at some time during their life (Table 12.2). These most common conditions are not specific diseases, nor are they necessarily sexually transmitted, but are often result from the presence of sexually transmitted infection. The diagnosis of these conditions does not always require special laboratory equipment or specialists, but can be done by obstetrician-gynecologists.

Other diseases that an at least 2% of respondents reported ever having been diagnosed with were: trichomoniasis (4%), syphilis (2%), gonorrhea (2%), and chlamydia (2%).

The lifetime incidence of almost every disease asked about was several times higher in urban areas than in rural areas. In the cases of syphilis and gonorrhea it was about seven times higher in urban areas. Only for PID and genital warts were the urban-rural differences relatively small. There is a possibility that those diseases that require special diagnostic laboratory equipment and methods for accurate diagnosis were underdiagnosed in rural areas.

The percentage of women ever diagnosed with almost all the STIs about which women were asked increased with respondents' age and education level. There is no obvious reason for a direct correlation between education and actual STI occurrence. It is more likely that the better educated women were more likely to have their conditions correctly diagnosed or to be aware of what their diagnosis was. This may support the hypothesis that there was underreporting by certain groups of women for many of the conditions asked about. The lifetime incidence of syphilis and gonorrhea among residents of the West region (6% and 5%, respectively) was at least three times higher than among respondents of the other regions. Trichomoniasis was much more likely to be reported by women in the South (6%) and East (4%) than elsewhere.

All sexually active women were asked whether they had experienced any selected symptoms potentially related to sexually transmitted infections during the past 12 months (Table 12.3). One-fifth of respondents reported experiencing vaginal discharge during that time. Vaginal discharge in combination with certain other symptoms is often indicative of STIs. Ten percent of women said that they experienced vaginal discharge accompanied by lower abdominal pain, 5% had vaginal discharge with itching, and 3% had vaginal discharge with painful urination. Fourteen percent of women complained of having sores or warts in the genital area during the previous 12 months. There were no substantial differences in the proportions reporting symptoms according to the residence, age, or educational attainment of women.

Treatment for possible symptoms of STIs

Almost one-third of respondents (31%) who experienced possible symptoms of STIs in the previous 12 months did not consult a health care provider for diagnosis or treatment (Table 12.4). The majority of respondents who experienced possible STI symptoms went to a women's consultation clinic or outpatient clinic for treatment (58%). The proportion of respondents who reported being treated at a private clinic or office was relatively small (3%). Fewer than 1% of women were treated at dermatovenereal disease (DVD) clinics. The proportion of women who did not seek treatment decreased with increasing education. It should come as no surprise that urban women were more likely than rural women to visit private clinics, probably because of better accessibility and higher incomes.

Respondents who did not seek treatment for symptoms of possible STIs gave a variety of reasons for not doing so (Table 12.5). Almost two-thirds of them (65%) reported that they thought their symptoms were not serious enough to warrant going for treatment. The only other reasons commonly given were: services were too far away or too expensive (11%); fear of poor treatment or confidentiality (6%); and embarrassment (3%). Feeling that the symptoms were not serious was the predominant reason for not seeking services among all categories of women. Distance and cost

(i.e., access to services) were most commonly mentioned by respondents with incomplete secondary education (22%). Fear of poor treatment and confidentiality was most common among urban respondents (7%), and previously married women (13%). Embarrassment was most likely to be mentioned by respondents who had never been in union and the youngest respondents. Not knowing where to go for services was rarely mentioned, except among 15-24 year-old respondents (4%).

Knowledge about prevention/Perceived risk of acquiring STIs

It is extremely difficult to significantly reduce the rate of STIs in a population that does not have knowledge about the risk factors, pathways, signs and symptoms, and preventive methods of these diseases. Each respondent was asked whether she was aware that individuals can be infected with HIV or STIs without showing any symptoms of disease. Survey responses showed considerable lack of knowledge among Ukrainian women about certain aspects of STIs and HIV/AIDS and their prevention (Table 12.6). About one-third of respondents were not aware that someone could be infected with HIV and exhibit no symptoms. About one-half of women lacked awareness that people with STIs could have no symptoms. Those respondents who lived in urban areas were somewhat more likely than rural women to be aware of this aspect of HIV and STIs. Lack of knowledge was especially common among the women residing in the West region of the country. There was little difference between age groups. Awareness of both HIV and STIs substantially increased with education. Sexually experienced women were more knowledgeable than inexperienced women, but only slightly so.

Multiple studies have demonstrated that condoms are highly effective in preventing many STIs, including HIV infection, when used consistently and correctly. Respondents were asked their opinions regarding the protection that condoms provide against HIV and other STIs. Only 6% of women thought that condoms provided excellent protection (Table 12.7). Roughly equal numbers of respondents said that condoms were good (30%) or fair (35%) at preventing transmission of infection. Twelve percent of respondents felt that condoms afforded poor protection. About 18% of women said they did not know about the effectiveness of condoms in preventing transmission of HIV/AIDS and other STIs. There was very little difference in opinions about condoms protective power between the subgroups of respondents, except for larger percentages answering "I do not know" among the less well educated and the sexually inexperienced. It is clear that Ukrainian women tend to underrate the protection afforded by condoms. There is a great need to inform the population better about the role of condoms in protection against the spread of infection, when used properly.

Perceptions of risk of acquiring STIs reflect the knowledge of a population about the basic risk factors and knowledge of means of preventing the transmission of these diseases. About 5% of the Ukrainian women perceived themselves at high risk for acquiring an STI, with another 6% saying they were at medium risk (Table 12.8). Another 29% of respondents felt that they were at low risk, while the majority, about six of every ten women considered themselves to be at no risk of contracting an STI. Women with multiple sex partners in the previous 12 months were substantially more likely to perceive themselves as being at high or medium risk of infection. Even so, it appears that many such individuals may have underestimated their risk, since about two-thirds considered themselves to be at low risk or no risk. Rural respondents, the youngest respondents, and those who reported having no sexual partners during the previous 12 months were the groups

of women who most often perceiving themselves not to be at risk for STIs.

Table 12.1
 Percentage of respondents who have never heard of selected STI's / conditions,
 by region, residence, age, education, and sexual experience
 1999 Ukraine Reproductive Health Survey

Characteristic	Syphilis	Gonorrhea	Trichomoniasis	Chlamydia	Genital herpes	Human papilloma virus (HPV)	Pelvic inflammatory disease (PID)	Genital ulcers	<i>Number of women</i>
<u>Total</u>	1.9	8.6	40.8	53.6	65.5	73.8	5.5	9.4	7,128
<u>Residence</u>									
Urban	1.8	6.5	35.4	48.1	60.7	69.8	4.9	8.9	5,544
Rural	2.5	14.2	55.0	68.3	78.1	84.5	7.2	10.6	1,584
<u>Regions</u>									
North	2.2	7.5	41.9	55.7	61.3	72.3	5.2	7.3	858
Central	1.4	9.9	40.5	64.3	66.7	73.1	4.2	8.6	617
East	2.5	6.2	35.8	45.8	62.4	69.4	3.6	8.5	2,669
South	0.7	5.4	31.4	45.3	60.7	73.1	5.3	9.6	1,824
West	2.3	15.3	56.4	66.5	77.2	83.5	9.9	12.6	1,160
<u>Age</u>									
15-24	2.9	15.3	52.6	63.6	73.9	82.5	9.4	24.0	2,282
25-34	1.5	5.4	34.4	48.2	60.6	69.7	3.5	1.6	2,422
35-44	1.5	5.1	35.4	49.1	62.1	69.4	3.7	2.6	2,424
<u>Education</u>									
< Complete secondary	4.5	23.6	64.1	75.6	82.7	90.4	10.9	35.5	778
Complete secondary	1.8	7.8	42.3	55.8	68.4	76.1	5.2	7.0	4,828
> Complete secondary	0.9	2.7	22.6	34.2	46.2	57.0	3.6	2.7	1,522
<u>Sexual experience</u>									
Experienced	1.6	6.2	36.4	50.2	63.2	71.7	3.9	3.9	6,218
Not experienced	4.5	23.8	68.5	75.3	79.9	87.2	15.9	43.5	910

Table 12.2
 Percentage of respondents who have ever been diagnosed with selected STIs /conditions,
 by region, residence, age, and education
 1999 Ukraine Reproductive Health Survey

Characteristic	Syphilis	Gonorrhea	Trichomoniasis	Chlamydia	Genital herpes	HPV	PID	Genital ulcers	<i>Number of women</i>
<u>Total</u>	2.1	1.7	3.7	1.5	0.8	0.9	38.2	37.8	7,128
<u>Residence</u>									
Urban	2.7	2.2	4.2	1.8	0.9	1.2	40.2	37.3	5,544
Rural	0.4	0.3	2.3	0.7	0.3	0.3	33.0	38.9	1,584
<u>Regions</u>									
North	0.3	0.3	2.9	1.0	1.4	0.9	41.8	38.1	858
Central	0.0	0.0	1.8	0.2	0.2	0.0	41.4	39.8	617
East	1.7	1.0	4.1	1.3	0.8	1.1	41.5	39.9	2,669
South	1.6	1.4	5.6	3.1	1.1	1.4	39.9	37.2	1,824
West	5.6	4.9	2.8	1.6	0.3	0.3	26.8	33.3	1,160
<u>Age</u>									
15-24	1.4	0.9	1.9	0.8	0.5	0.8	20.9	14.6	2,282
25-34	2.4	2.2	4.7	2.4	0.7	0.9	44.3	46.7	2,422
35-44	2.4	1.9	4.3	1.3	1.1	1.2	49.1	50.7	2,424
<u>Education</u>									
< Complete	1.3	0.9	1.7	0.7	0.4	0.2	23.8	16.4	778
Complete secondary	2.2	1.7	3.7	1.4	0.7	0.9	39.8	40.1	4,828
> Comp. secondary	2.4	2.1	4.7	2.4	1.3	1.6	41.1	41.9	1,522

Table 12.3
 Percent of sexually active women who have experienced selected symptoms
 consistent with sexually transmitted infections (STI) during the past 12 months, by age, education, and place of residence
 1999 Ukraine Reproductive Health Survey

Characteristic	Vaginal discharge	Vaginal discharge with itching	Vaginal discharge with painful urination	Vaginal discharge with lower abdominal pain	Sores or warts in the genital area	<i>Number of women</i>
<u>Total</u>	20.4	4.7	2.7	10.3	14.3	5,481
<u>Residence</u>						
Urban	20.3	5.2	2.8	10.0	13.7	4,252
Rural	20.7	3.2	2.7	10.9	15.8	1,229
<u>Age</u>						
15-19	20.3	2.7	1.6	8.9	7.0	398
20-24	24.1	6.1	3.0	10.4	14.8	955
25-29	23.5	4.9	2.2	11.9	15.7	1,084
30-34	22.4	6.1	4.4	11.8	15.8	1,066
35-39	16.9	3.6	2.1	8.9	14.1	1,041
40-44	15.0	3.4	2.3	8.6	14.1	937
<u>Education</u>						
< Complete secondary	18.2	2.7	2.7	10.4	15.2	405
Complete secondary	20.4	4.5	2.8	10.6	14.4	3,829
> Complete secondary	21.4	5.8	2.4	9.1	13.6	1,247

Table 12.4
 Percentage distribution of place of treatment for possible symptoms of STI, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Place of treatment						Total	Number of women
	No consultation or treatment	Women's consultancy / outpat. clinic	Dermato-venereal clinic or office	Private office/clinic	Other	Don't remember		
<u>Total</u>	30.6	58.3	0.7	3.4	3.6	3.4	100.0	1,445
<u>Residence</u>								
Urban	30.6	57.5	1.0	4.2	3.6	3.2	100.0	1,097
Rural	30.5	60.4	0.0	1.5	3.5	4.0	100.0	348
<u>Age</u>								
15-19	31.0	51.3	2.4	2.8	5.2	7.3	100.0	90
20-24	30.8	52.8	0.0	6.6	6.3	3.5	100.0	283
25-29	27.4	63.8	0.9	3.7	2.8	1.4	100.0	312
30-34	30.9	59.2	0.0	2.5	3.3	4.0	100.0	305
35-39	31.9	59.9	1.3	1.5	2.4	3.0	100.0	242
40-44	32.7	58.0	0.5	2.9	1.9	4.0	100.0	213
<u>Education</u>								
< Complete secondary	37.7	52.9	0.0	2.1	3.9	3.3	100.0	112
Complete secondary	31.1	58.6	0.8	2.9	3.2	3.6	100.0	994
> Complete secondary	26.4	59.5	0.9	5.8	4.6	2.8	100.0	339

Table 12.5
 Percentage distribution of reasons for not seeking treatment for possible STI symptoms,
 by place of residence, and union status
 1999 Ukraine Reproductive Health Survey

Characteristic	Primary reason for not seeking treatment							Total	Number of women
	Thought it was not serious	Services too far away / too expensive	Feared poor treatment / confidentiality	Embarrassment	Didn't know where to go	Other	DK/DR		
<u>Total</u>	65.2	11.4	6.4	2.9	1.3	8.5	4.3	100.0	471
<u>Residence</u>									
Urban	65.8	10.5	7.0	2.7	1.1	9.3	3.6	100.0	363
Rural	63.5	13.8	4.7	3.7	1.9	6.5	5.9	100.0	108
<u>Age</u>									
15-24	62.8	8.4	7.2	4.4	3.6	8.2	5.4	100.0	126
25-34	67.8	11.4	6.7	2.4	0.6	6.4	4.7	100.0	188
35-44	63.9	14.1	5.4	2.4	0.0	11.4	2.8	100.0	157
<u>Education</u>									
< Complete secondary	51.4	21.9	6.4	4.3	0.0	6.7	9.3	100.0	44
Complete secondary	66.7	10.8	5.9	3.5	1.1	8.3	3.7	100.0	323
> Complete secondary	66.1	8.7	8.1	0.6	2.5	10.2	3.8	100.0	104
<u>Marital status</u>									
Currently in union	65.8	11.7	5.7	2.1	1.2	9.8	3.7	100.0	389
Previously in union	66.4	16.4	12.6	3.6	0.0	1.0	0.0	100.0	34
Never in union	59.9	5.4	7.8	9.1	2.4	3.7	11.7	100.0	48

Table 12.6
 Percent of respondents who are aware that individuals can be infected with HIV/AIDS or have other sexually transmitted infections (STIs)
 without showing any signs or symptoms of disease, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	HIV / AIDS			STIs			Women
	Aware	Not aware	Don't know*	Aware	Not aware	Don't know	
<u>Total</u>	67.2	15.0	17.8	51.1	33.1	15.9	7,128
<u>Residence</u>							
Urban	69.6	14.1	16.3	52.5	33.0	14.5	5,544
Rural	60.8	17.3	21.9	47.2	33.2	19.6	1,584
<u>Regions</u>							
North	71.9	10.9	18.0	59.3	24.0	16.7	858
Central	72.2	12.6	15.2	58.1	27.2	14.7	617
East	70.6	14.5	15.0	47.2	39.4	13.4	2,669
South	69.4	13.0	17.5	55.9	31.2	12.9	1,824
West	54.1	21.8	24.1	43.7	33.8	22.5	1,160
<u>Age</u>							
15-24	66.4	17.6	16.0	49.4	33.6	17.0	2,282
25-34	68.8	14.1	17.2	55.2	30.9	13.9	2,422
35-44	66.5	13.4	20.1	48.7	34.6	16.7	2,424
<u>Education</u>							
< Comp. secondary	57.5	17.7	24.8	38.4	36.3	25.3	778
Comp. secondary	65.7	15.6	18.8	50.7	33.0	16.3	4,828
> Comp. secondary	77.3	11.5	10.7	59.3	31.4	9.3	1,522
<u>Sexual experience</u>							
Experienced	67.9	14.5	17.6	52.3	32.9	14.8	6,218
Not experienced	63.1	18.0	19.0	43.7	33.5	22.9	910

*Includes 7 women who said they had not heard of HIV/AIDS.

Table 12.7
 Percentage distribution of opinions regarding the protection that condoms gives against HIV and other STI's,
 by residence, region, age, education, and sexual experience
 1999 Ukraine Reproductive Health Survey

Characteristic	Protection against disease provided by condoms					Total	Women
	Excellent	Good	Fair	Poor	Don't know		
<u>Total</u>	5.8	29.7	35.1	11.8	17.5	100.0	7,128
<u>Residence</u>							
Urban	6.2	30.2	35.2	12.1	16.3	100.0	5,544
Rural	4.8	28.6	34.7	11.1	20.9	100.0	1,584
<u>Region</u>							
North	5.9	30.8	35.7	10.8	16.8	100.0	858
Central	7.4	32.2	31.5	12.0	16.9	100.0	617
East	6.8	27.1	36.6	11.8	17.6	100.0	2,669
South	4.9	31.1	36.3	14.3	13.4	100.0	1,824
West	4.0	30.8	32.9	10.5	21.8	100.0	1,160
<u>Age</u>							
15-24	5.2	31.4	33.3	11.7	18.4	100.0	2,282
25-34	6.9	30.3	36.5	11.9	14.5	100.0	2,422
35-44	5.4	27.5	35.5	11.9	19.7	100.0	2,424
<u>Education</u>							
< Comp. secondary	3.3	15.5	29.6	12.2	29.4	100.0	778
Comp. secondary	6.0	29.4	34.3	12.3	18.0	100.0	4,828
> Comp. secondary	6.6	33.1	40.8	10.1	9.5	100.0	1,522
<u>Sexual experience</u>							
Experienced	6.2	29.8	35.8	12.1	16.1	100.0	6,218
Not experienced	3.2	29.2	30.3	10.4	26.9	100.0	910

Table 12.8
 Percentage distribution of respondents' perceived risk of sexually transmitted infection,
 by residence, region, age, education, and number of recent sexual partners
 1999 Ukraine Reproductive Health Survey.

Characteristic	Perceived risk of sexually transmitted infection					Total	N
	High	Medium	Low	None	Don't know		
<u>Total</u>	5.2	6.4	22.6	59.0	6.3	100.0	7,128
<u>Residence</u>							
Urban	5.2	6.3	24.7	57.0	6.8	100.0	5,544
Rural	5.3	6.8	16.8	64.3	6.9	100.0	1,584
<u>Region</u>							
North	4.1	6.6	20.8	57.3	11.3	100.0	858
Central	6.2	5.9	19.4	64.0	4.5	100.0	617
East	4.9	5.5	24.9	57.9	6.8	100.0	2,669
South	6.4	9.4	27.9	52.2	4.1	100.0	1,824
West	4.7	5.7	16.9	65.4	7.1	100.0	1,160
<u>Age</u>							
15-24	5.9	6.6	19.1	61.1	7.3	100.0	2,282
25-34	5.4	6.5	26.1	55.5	6.6	100.0	2,422
35-44	4.3	6.1	22.5	50.5	6.7	100.0	2,424
<u>Education</u>							
< Comp. secondary	5.8	4.1	14.7	67.2	8.3	100.0	778
Comp. secondary	5.6	6.2	21.6	59.6	7.0	100.0	4,828
> Comp. secondary	3.5	8.5	30.1	52.5	5.5	100.0	1,522
<u>Recent sex partners</u>							
0	4.6	3.7	10.1	74.5	7.1	100.0	1,215
1	4.9	6.7	24.5	57.9	6.0	100.0	5,100
2	10.0	16.8	37.2	30.7	5.3	100.0	243
3 +	11.3	14.2	29.3	37.3	7.9	100.0	79
No response	4.6	1.9	26.8	49.4	17.3	100.0	471

CHAPTER XIII DOMESTIC VIOLENCE

Violence against women has been increasingly recognized as an important clinical and public health issue. Violence and the threat of violence can result not only in psychological dysfunction and physical trauma, but also has the ability to affect women's reproductive health. Intimate partner violence should be of particular concern to women of reproductive age and their health providers. Intimate partner violence can put women at higher risk for lack of contraceptive use or inconsistent use, causing women to remain unprotected against STIs and unintended pregnancy. Violence during pregnancy can also be associated with inadequate prenatal care (Martin et al., 1999), increased maternal morbidity (Parker 1994), and poor pregnancy outcomes (Petersen et al. 1997, Campbell et al. 1999). The children of women who are subjected to intimate partner violence can also be adversely affected. Children of such women have been shown to have elevated rates of emotional, behavioral, and cognitive abnormalities (Jaffe et al. 1990).

There is little, if any, nationally representative data available on the extent of the problem of domestic violence in Ukraine. "While information about domestic violence in Ukraine from official sources is limited, the interviews conducted by the Minnesota Advocates confirmed that it is a widespread problem in the country." (Minnesota Advocates for Human Rights, 2000). For instance, a survey by the World Bank and the KIIS showed that 12% of women under age 28 said they had experienced violence at the hands of their husbands. A large percentage of police calls are related to domestic abuse, as are a large percentage of emergency room visits.

The 1999 URHS included a module with questions about women's lifetime and previous year's experience of being subjected to physical violence or threats of violence by their partner. Although there is a high risk of underreporting of such incidents, the survey serves to demonstrate the minimum proportion of women who are victims of violent behavior or the threat of such behavior in their homes. The module also included questions on abuse between parents when the respondent was growing up and abuse of the respondent as a child. Childhood exposure to abuse can exert long-term negative effects on interpersonal relationships, emotional distress, somatic symptoms, and substance abuse (Felitti et al. 1999). Almost one of every five respondents (18%) recalled that their parent or stepparent had been abused by their partner while the respondent was growing up (Table 13.1). Almost 30% of women reported that they were physically abused as a child by someone in their household. There were some noticeable, but not extreme, differentials in the proportions reporting these adverse childhood experiences among residential and sociodemographic groups. There were no consistent differences according to residence or age. The least well educated were the most likely to have these experiences. With regard to marital status, violent experiences were most common among those who lived in unregistered marriage, followed by those who had been divorced or separated. The proportion with parents who abused each other was about twice as high among women in unregistered marriages as among the never married.

Each respondent who had ever been married or lived with a man was asked questions whether she had been subjected to particular types of physical violence or threats of violence committed by her partner, either at any time during her life or in the previous 12 months (Table 13.2). Overall, 19% of ever-married women had ever had a partner threaten to hit her, 18% had ever been pushed or slapped, 13% had been punched, kicked, or hit with an object, and 4% had been threatened with a

weapon. Twenty-one percent had had any of these acts committed against her by her partner. Eight percent had experienced any of these types of violence within the previous 12 months, with most of them having had her partner threaten to hit her or actually pushing or slapping her.

The proportions reporting that violent incidents or the threat of them had ever occurred was consistently, but only slightly, higher in rural areas than in urban areas. Differentials were somewhat greater for incidents in the previous 12 months. Women with incomplete secondary education were roughly twice as likely to report violent incidents as the most highly educated women, with the complete secondary group between them. As age increased, so did the likelihood of ever experiencing violent incidents. However, this appears to be solely a function of the number of years exposed to the risk of violence, since the proportions reporting such incidents in the past 12 months was fairly constant across ages for each of the specific types of violence. In fact, the proportion of 15-19 year-olds reporting any type of violence recently was slightly higher than for any other age group. Divorced women and women who lived in unregistered marriage reported being abused considerably more often than respondents in registered marriages. Particularly alarming is the high proportion of the youngest women who reported domestic violence incidents, especially threats with a weapon, during the past year. (Rates for the previous 12 months for divorced, separated, and widowed women are misleadingly low, since most of these women were not been exposed to the risk of partner abuse in the months leading up to interview.) It is noteworthy that women in unregistered marriages were about twice as likely as women in registered marriages to suffer abuse, both over their lifetime and recently.

Fifty-four percent of women who reported abuse by their partner in the previous 12 months reported sustaining injuries from these incidents. Seven percent of abused women sought medical treatment, with about half of them (3%) incurring injuries serious enough to be hospitalized (Table 13.3). Even though urban women were less likely than rural women to be victims of violence, abused urban women were more likely to say they sustained injuries and sought treatment than rural women. The percentage reporting injury tended to be highest among those who had highest education and those who were not in a registered marriage.

Most women who had been physically abused discussed these incidents with others. These discussions were most often with family members (68% of women subject to violent incidents) or friends (56%) (Table 13.4). Women were relatively unlikely to discuss incidents with anyone other than family and friends. Just 16% of women went to the police for the help. A relatively small proportion of them (9%), probably predominantly those who sought medical treatment, had discussions with medical or social workers. Rural women, younger women, and those in a registered marriage were less likely than others to discuss these incidents than other subgroups of women were. The groups most likely to report violent incidents to the police were divorced and separated women (49%) and women 35-44 years old (22%).

Alcohol use has been documented as a major health and social problem in much of the former Soviet Union. It has been shown to be a major contributor to declines in male life expectancy and to the deterioration of health status generally in Russia. It has also been linked to problems of violence generally, including the occurrence of intimate partner violence. Clearly, alcohol consumption was closely related to violent behavior among the URHS respondents. Among women subjected to intimate partner violence in the previous 12 months, 78% reported that their partner

had been drinking alcohol at the time of the most recent abuse (Table 13.5). Eighty-three percent said he had been drinking at any time in the past year when she was subjected to violent incidents. Violence was closely tied to violent acts throughout the population, but was even more prevalent among certain subgroups than among others. Violence was most often correlated with alcohol consumption among rural women, better educated women, and women ages 30-44.

Table 13.1
 Percent of respondents who recall parent/step-parent physically abused by partner
 and percent who were physically abused as children by someone in their household,
 by selected characteristics
 1999 Ukraine Reproductive Health Survey

<u>Characteristic</u>	Parent abused by his/her partner	Respondent abused as a child	Number of women
<u>Total</u>	18.6	29.3	7,128
<u>Residence</u>			
Urban	18.5	29.2	5,544
Rural	18.9	29.5	1,584
<u>Education</u>			
< Complete secondary	22.8	32.9	778
Complete secondary	19.4	30.0	4,828
> Complete secondary	13.8	24.9	1,522
<u>Age</u>			
15-24	16.9	29.3	2,282
25-34	19.9	31.2	2,422
35-44	19.0	27.4	242
<u>Marital status</u>			
Registered marriage	18.3	29.5	4,495
Unregistered marriage	31.2	42.2	301
Divorced / Separated	22.7	31.4	676
Widowed	22.3	28.2	124
Never married	15.4	25.9	1,532

Table 13.2
 Percentage of ever-married respondents subjected to particular types of violence ever and during the past 12 months
 1999 Ukraine Reproductive Health Survey

Characteristic	Type of violence										No. of women
	Threaten to hit or throw object at her		Push or slap her		Hit her with fist or an object, or kick her		Threaten her with a knife or other weapon		Any listed type of violence		
	Ever	Last 12 months	Ever	Last 12 months	Ever	Last 12 months	Ever	Last 12 months	Ever	Last 12 months	
Total	19.1	6.2	18.3	6.6	12.9	4.5	3.6	0.8	21.1	7.9	5,596
Residence											
Urban	18.5	6.2	17.7	5.9	12.2	3.9	3.6	0.8	20.4	6.9	4,312
Rural	20.6	8.6	19.6	8.3	14.7	6.1	3.6	0.9	22.3	9.7	1,284
Education											
< Complete secondary	26.1	9.8	25.5	9.9	19.6	6.8	6.0	1.2	28.1	11.5	335
Complete secondary	19.9	7.1	18.9	6.7	13.6	4.6	3.9	0.9	22.0	8.2	4,018
> Complete secondary	14.4	5.4	13.9	5.4	8.7	3.5	1.8	0.4	16.1	6.2	1,243
Age											
15-19	7.3	7.0	6.4	5.2	5.0	4.7	2.1	1.9	12.2	10.1	127
20-24	13.9	7.3	14.8	7.3	8.8	4.0	1.6	0.6	16.8	8.7	800
25-29	18.2	7.2	16.7	7.0	12.1	4.9	3.9	1.3	19.4	8.1	1,139
30-34	18.6	7.4	18.5	7.4	12.3	5.3	2.2	0.3	20.9	8.7	1,159
35-39	20.5	6.1	19.1	5.4	14.3	3.5	4.5	0.7	22.4	6.6	1,215
40-44	23.8	6.6	22.1	6.4	16.6	4.8	5.2	1.1	25.5	7.6	1,156
Marital status											
Registered marriage	14.6	6.5	14.0	6.4	9.1	4.1	2.1	0.8	16.5	7.6	4,495
Unregistered marriage	33.1	13.5	31.5	11.4	26.5	10.7	7.7	1.5	36.4	15.7	301
Divorced / Separated	43.2	7.4	40.3	7.1	32.1	5.4	11.5	1.1	45.4	8.4	676
Widowed	24.4	1.8	24.4	1.8	20.0	1.8	7.8	0.3	25.6	1.8	124

Table 13.3
 Percentage of respondents abused in past 12 months who
 sustained injuries, sought treatment, or were hospitalized as a result, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	% who sustained injuries	% who sought medical treatment	% who were hospitalized	<i>Number of women</i>
<u>Total</u>	53.9	7.3	3.1	392
<u>Residence</u>				
Urban	58.0	9.2	3.5	267
Rural	46.6	4.1	2.4	125
<u>Education</u>				
< Complete secondary	49.2	4.4	4.4	36
Complete secondary	52.6	7.7	2.5	285
> Complete secondary	62.6	7.3	7.3	71
<u>Age</u>				
15-19	*	*	*	13
20-24	51.5	5.1	1.9	62
25-29	49.6	8.5	2.8	76
30-34	62.3	6.2	1.3	91
35-39	48.0	6.7	5.1	69
40-44	57.0	9.4	5.1	81
<u>Marital status</u>				
Registered marriage	51.4	4.5	2.7	296
Unregistered marriage	59.7	9.2	0.8	44
Divorced / Separated	64.5	20.9	6.8	49
Widowed	*	*	*	3

*Fewer than 25 women in category

Table 13.4
 Percentage of respondents abused in past 12 months
 who discussed these incidents with various types of people, by selected characteristics
 1999 Ukraine Reproductive Health Survey

Characteristic	Person with whom discussed incident						Number of women
	Police	Family members	Friends	Medical/Social worker	Psychologist	Others	
<u>Total</u>	15.9	67.9	56.2	8.7	2.2	3.6	392
<u>Residence</u>							
Urban	18.4	69.6	58.1	8.9	2.9	3.8	267
Rural	11.4	65.0	52.6	8.1	0.8	3.3	125
<u>Education</u>							
< Complete secondary	12.7	78.7	52.9	4.9	0.0	3.4	36
Complete secondary	17.5	66.2	56.0	9.0	2.4	3.7	285
> Complete secondary	10.7	70.0	58.4	9.0	2.2	3.5	71
<u>Age</u>							
15-19	*	*	*	*	*	*	13
20-24	6.5	66.2	38.2	5.4	0.0	1.9	62
25-29	16.3	65.6	57.5	7.1	4.3	5.7	76
30-34	10.6	62.1	60.5	6.2	0.4	1.3	91
35-39	23.7	72.1	63.5	13.0	0.0	4.1	69
40-44	21.4	75.2	57.8	11.4	6.0	5.7	81
<u>Marital status</u>							
Registered marriage	10.4	66.4	53.2	4.9	1.9	3.7	296
Unregistered marriage	18.4	66.1	45.3	5.5	0.8	0.0	44
Divorced / Separated	48.9	79.3	83.1	3.5	4.7	4.7	49
Widowed	*	*	*	*	*	*	3

*Fewer than 25 women in category

Table 13.5

Percent of respondents abused in past 12 months whose partner had been drinking alcohol the last time he abused her, and percent whose partner had been drinking at any time in the past year when he abused her, by selected characteristics
1999 Ukraine Reproductive Health Survey

Characteristic	Partner had been drinking alcohol last time he abused her	Partner had been drinking alcohol any time in the past year when he abused her	<i>Number of women</i>
<u>Total</u>	78.2	83.2	392
<u>Type of residence</u>			
Urban	75.7	79.9	267
Rural	82.7	89.2	125
<u>Education</u>			
< Complete secondary	62.8	71.2	36
Complete secondary	79.9	83.6	285
> Complete secondary	78.4	87.4	71
<u>Age</u>			
15-19	*	*	13
20-24	61.6	70.8	62
25-29	72.5	76.2	76
30-34	86.0	92.2	91
35-39	88.2	92.9	69
40-44	86.2	87.6	81
<u>Marital status</u>			
Registered marriage	78.8	83.6	296
Unregistered marriage	75.6	78.6	44
Divorced / Separated	79.3	87.4	49
Widowed	*	*	3

*Fewer than 25 women in category

CHAPTER XIV

CONCLUSIONS

Analysis of data from the 1999 Ukraine Reproductive Health Survey has allowed us to draw a number of important conclusions. Some of the most significant are the following:

- The 1999 URHS appears to be highly representative of the population of women of childbearing age in Ukraine. Distributions of age and other characteristics of respondents closely match those from official sources.
- In key areas the data collected in the survey appear to be relatively reliable and complete. Survey fertility rates closely resemble official rates. Survey abortion rates exceeded those from vital statistics by a significant amount, indicating that the completeness of reporting was likely to be good. Response rates were high and refusal rates were low. Few respondents were unwilling to answer questions on sensitive topics, such as abortion, contraception, sexuality, and health problems.
- Although there are certainly some important differences across countries in the region, there are also great similarities between many countries that were either part of the Soviet Union or under the domination of the Soviet Union prior to its break-up. Because of the commonalities between these countries socially, culturally, economically, and demographically, many of the general conclusions derived from the 1999 URHS are likely to be applicable to neighboring countries, and can be valuable in suggesting reproductive health policies and priorities there.
- Not only did the surveyed population have extremely low rates of childbearing, there is no indication that Ukrainian women intend to have larger families than currently. Few women with more than one child desired to have additional children and large numbers of women with just one child said they planned to have no more. Such low levels of desired childbearing, especially with the limited use of effective long-term contraceptive methods and a typically early start and finish of desired fertility, enhances the probability of unintended pregnancies and abortions. Although a substantial number of women reported infertility problems, there is no reason to believe that infertility (or low rates of sexual activity) is an important factor in bringing about the low fertility rate.
- Rates of induced abortion, though not nearly as high as in some other countries in the region, such as Russia, Georgia, and Romania, remain very high by international standards. The induced abortion rate was more than 50 per 1,000 woman per year for the period preceding the survey and the number of abortions still exceeded the number of live births. However, it seems clear that the incidence of induced abortion has been declining at a relatively rapid rate. Obviously, there is considerably more progress that can be made in reducing the rate of abortion, both through increased use of modern contraception and

improved use of contraceptive methods.

- Rates of abortion complications remain relatively high. About one-sixth of abortions result in what women describe as complications, and one of every 20 abortions results in hospitalization for those complications. Even though we have no data on severity of complications, such a rate of complications is almost certainly producing much greater maternal morbidity and greater costs for the health care system than replacing most abortion with successful contraception would.
- Overall contraceptive prevalence rates among sexually active women were high, not far below rates for most other developed countries. The IUD and condoms were the only modern methods in widespread use nationally. In contrast to what some have contended, Ukrainian women were not relying predominantly on abortion for birth prevention. Women tend to have highly negative opinions about induced abortion, but have used it frequently seemingly because of inadequate supplies of modern contraceptives or lack of accurate information concerning contraception and high rates of pregnancy while using contraceptive methods. In spite of the relatively high rate of contraceptive prevalence, unmet need for contraception remains quite high, mainly because the desire for children is so low.
- Almost half of contraceptors were employing methods with high failure rates, contributing to Ukraine's high incidence of unintended pregnancy and induced abortion. Withdrawal was the most commonly used method, with periodic abstinence also commonly employed. Because of the high failure rates associated with these methods, substantial declines in abortion incidence depend to a certain extent on switching to more effective methods and not just promoting contraceptive use in general. Only about one of every five users of these methods were aware of their poor effectiveness relative to modern methods.
- Because of the typically early end of desired childbearing among Ukrainian women, a great need for expanded use of long-term contraceptive methods exists. Except for the IUD, long-term methods were rarely used in Ukraine. Clearly, there are major barriers (legal, social, resource-related) to rapid expansion of contraceptive sterilization, but steps can be taken to improve and increase acceptability and availability of these safe and effective procedures. Another relatively long-term method, contraceptive implants, is now becoming more widely available in Ukraine. Unless contraceptive sterilization and other long-term methods become a realistic option for Ukrainian women and couples, unintended pregnancies are likely to remain common.
- Contraceptive counseling following delivery and especially induced abortion should be given greater attention. Relatively few women receive contraception of prescriptions for methods following these events, leading to a higher incidence of unintended pregnancy and induced abortion than there would likely be with more and effective counseling.
- Sexual activity among younger teenagers, while still not the norm, has been increasingly

common. It is not unreasonable to expect that activity will increase even further. This increased activity, plus the fact that fewer than half of young respondents reported the use of contraception the first time they had sexual intercourse, as well as increases in rates of sexually transmitted infections, points toward a need for increased sex education and policies designed to protect the health of adolescents.

- About one in every ten respondents with a recent pregnancy resulting in a live birth reported that they received no prenatal care during that pregnancy, considerably higher than the reports from other countries in the region. This finding should be further investigated to determine the accuracy of the result. If it is accurate, it is important that actions be taken to increase prenatal care coverage to reduce both maternal and infant morbidity.
- Rates of pregnancy-related hospitalization, as well as the typical length of those hospitalizations, were high in Ukraine, as in much or all of the former Soviet Union. This situation should be further studied to determine how much of the hospitalization stems from conditions that truly warrant it and how much results from a greater tendency than in the West to hospitalize for relatively minor conditions. The findings could point toward a need for more and better prenatal services to reduce the likelihood or severity of conditions leading to hospitalization or toward a reduction in the incidence of hospitalization given the occurrence of certain conditions.
- Sexually transmitted infections are a serious problem in Ukraine and, by all indications, are a growing problem as well. Besides high lifetime incidence of several conditions and infections, substantial numbers of women reported recent combinations of symptoms that are most often associated with sexually transmitted infections. About one-third of women who experienced recent symptoms that possibly resulted from STIs did not seek treatment for those symptoms. Such untreated conditions can progress to more serious complications for women and can lead to greater spread of infection.
- There were very large percentages of women with inadequate knowledge of certain aspects of STIs and HIV, indicating a need for better communication of health messages to the population. For instance, about one-half of respondents did not know that people infected with STIs might exhibit no symptoms. Only about one-third said that condoms provided good or excellent protection from infection with HIV or other STIs.
- Domestic abuse is increasingly recognized as a major problem, not just in Ukraine, but in many countries around the world. It is commendable that those carrying out the survey were strongly encouraged to collect information on this sensitive issue. Even if the figures from the URHS are underestimated, it still demonstrates that violence by men against their partners and resulting injuries are a common occurrence in Ukraine. It is also quite clear that alcohol plays a significant role in domestic violence. These results and other recent work on the subject points out the need to take action to try to reduce the incidence of violence.

Overall, the URHS has provided a wealth of information on a wide range of reproductive health topics and yields a picture of much of the current reproductive health situation in the country. It, more importantly, yields information for the nation, donor organizations, international agencies, and NGOs that can be used to help determine the scope of various reproductive health needs and the types of interventions and policies that might help improve the situation. The information presented in this report is by no means an exhaustive analysis of the data collected. There remains a large amount of information on all topics that can be further analyzed both by in-country and external researchers and reproductive health experts in order to help improve reproductive health status in Ukraine.

REFERENCES

All-Russian Centre for Public Opinion and Market Research, U.S. Centers for Disease Control and Prevention, U.S. Agency for International Development. 1998. 1996 Russia Women's Reproductive Health Survey: A study of three sites, Final Report.

Barnett T, Whiteside A, et al. 2000. The HIV/AIDS epidemic in Ukraine: its potential social and economic impact. *Social Science and Medicine*. 51(9), 1387-1403.

Barr DA, Field MG. 1996. The current state of health care in the former Soviet Union: Implications for health care policy and reform. *American Journal of Public Health*, 86(3):304-320.

Bergthold, G, Rooks J, Stewart G. 1998. Evaluation of the Women's Reproductive Health Initiative in Ukraine. Population Technical Assistance Project, Arlington VA.

Blayo C. 1993. The role of abortion in the countries of central and eastern Europe. In: *International Population Conference*, Montreal Canada, 24 August-1 September, 1993. Compiled by the International Union for the Scientific Study of Population, Liege Belgium, IUSSP, pp. 235-252.

Campbell J et al. 1999. Physical and nonphysical partner abuse and other risk factors for low birth weight among full term and preterm babies: a multiethnic case-control study. *American Journal of Epidemiology* 150(7):714-726.

Dehne KL et al. 1999. The HIV/AIDS epidemic in eastern Europe: recent patterns and trends and their implications for policy-making. *AIDS* 13(7):741-749.

Felitti VJ, Anda RF, Nordenberg D et al. 1999. The relationship of selected health-risk behaviors, health status and disease in adulthood from childhood abuse and household dysfunction. *Am J Prev Med* 14:245-58.

Goldberg HI, Melnikova N. 2001. Effects of contraceptive behavior on abortion levels in Ukraine. Presented at the 2001 Annual Meeting of the Population Association of America, Washington DC, March 29-31.

Grischenko VI. Research and service needs in reproductive health in Ukraine. In: Assessment of research and service needs in reproductive health in Eastern Europe—concerns and commitments. Proceedings of a workshop organized by the ICPR and the WHO Collaborating Centre on Research in Human Reproduction in Szeged, Hungary, 25-27 October 1993, edited by E. Johannisson et al.: New York, NY, Parthenon Publishing Group.

Hatcher RA, Trussell J, Stewart F, et al. 1998. *Contraceptive Technology: Seventeenth Revised Edition*. Ardent Media Inc., New York.

Jaffe PG, Harley DJ, Wolfe D. 1990. Children's observations of violence: I. Critical issues in child development and intervention planning. *Can J Psychiatry* 35(6):466-70.

Lifson AR, Preble E, 1997. An assessment of the HIV/AIDS/STD Situation in Ukraine. Health Technical Services Project, Arlington VA.

Little RE, Monaghan SC, Gladen BC, Shkyryak-Nyzhnyk ZA, Wilcox AJ. 1999. Outcomes of 17137 pregnancies in 2 urban areas of Ukraine. *American Journal of Public Health*, 89(12): 1832-6.

Martin SL, Clark KA, Lynch SR, Kupper LL, et al. 1999. Violence in the lives of pregnant teenage women: associations with multiple substance abuse. *American Journal of Drug and Alcohol Abuse* 25(3): 425-40.

Minnesota Advocates for Human Rights. 1996. Domestic Violence in Ukraine. Minneapolis, MN

Parker B, McFarlane J, Soeken K. 1994. Abuse during pregnancy: effects on maternal complications and birth weight in adult teenage women. *Obstet Gynecol*: 323-8.

Petersen R et al. 1997. Violence and adverse pregnancy outcomes: a review of the literature and directions for future research. *American Journal of Preventive Medicine* 13(5): 366-373.

Popov AA. 1991. Family planning and induced abortion in the USSR: Basic health and demographic characteristics. *Studies in Family Planning*, 22(6):368-377.

Population Reference Bureau. 2000. World Population Data Sheet.

Renton AM, Borisenko KK, et al. 1998. Epidemics of syphilis in the newly independent states of the former Soviet Union. *Sex. Transm. Infect.*:165-6.

Savblom C. 1999. Highlight on women's health—Ukraine. *Entre Nous*; 45, pp. 13-14.

Steshenko T, Irkina T. 1999. Some Aspects of the Demographic Situation in Ukraine: Reproductive Health and Family Planning. United nations Population Fund, Kyiv.

Tichanova L, Borisenko K, Ward H, et al. 1997. Epidemics of syphilis in the Russian Federation: trends, origins and priorities for control. *Lancet*: 350:210-13.

Ukraine Cabinet of Ministers, Ukraine Ministry of Health. 1997. The Health of Women and Children in Ukraine. Kyiv.

Ukraine Central Statistics Committee. 1998. Children, Women, and Family in Ukraine, 1998. Kyiv.

UNICEF. 1996. Childhood and Motherhood in Ukraine '96. Kyiv.

United Nations Development Programme. 1997. Ukraine Human Development Report. Kyiv.

Vovk IV. 1997. The importance of family planning in Ukraine. In: Assessment of research and service needs in reproductive health in Eastern Europe—concerns and commitments. Proceedings of a workshop organized by the ICPR and the WHO Collaborating Centre on Research in Human Reproduction in Szeged, Hungary, 25-27 October 1993, edited by E. Johannisson et al.: New York, NY, Parthenon Publishing Group.

Waugh MA. 1999. Task force for the urgent response to the epidemics of sexually transmitted diseases in eastern Europe and central Asia.. *The Journal of Sexual Health & HIV*: 75(1): 72-73.

3. How many females between the ages of 15 and 44 live in this flat/house?

___ females

4. For each of these women could you give me the following information:

LIST FROM OLDEST TO YOUNGEST

Line	First name	Age	Marital status	Education	Selection	Result
1	_____	---	---	---	---	---
2	_____	---	---	---	---	---
3	_____	---	---	---	---	---
4	_____	---	---	---	---	---
5	_____	---	---	---	---	---
6	_____	---	---	---	---	---
			CODES:	CODES:	CODES:	CODES:
			1 Married	1 No secondary	1 Selected	1 Complete
			2 Unregistered marriage	2 Incomp Secondary	2 Not sel.	2 Not home
			3 Divorced	3 Comp. Secondary		3 Refusal
			4 Separated	4 Prof Technical Ed		4 Incompetent
			5 Widowed	5 CompSec + TechEd		5 Other: _____
			6 Single (Never married)	6 Technicum		
				7 Incomp. Postsec.		
				8 Complete Postsec.		
				9 Don't know		

HOUSEHOLD VISIT RECORD

Visit number	1	2	3	4
	Day Month	Day Month	Day Month	Day Month
Date of visit	--- ---	--- ---	--- ---	--- ---
Result*	---	---	---	---

***RESULT CODES**

- 1 Completed household interview--at least one women 15-44 living in household
- 2 Completed household interview--no women 15-44 living in household
- 3 Nobody at home/No adults at home
- 4 Refusal
- 5 Unoccupied dwelling
- 6 Other _____

1999 UKRAINE WOMEN'S REPRODUCTIVE HEALTH SURVEY
Individual questionnaire

I. BACKGROUND CHARACTERISTICS

100. In what month and year were you born?

Month ___ ___

Year 19 ___ ___

101. How old are you?

___ ___ years old

(MAKE SURE THAT AGE AND DATE OF BIRTH CORRESPOND)

102. What is the highest level of education you completed? (SHOW CARD)

- 1 Less than grade 7
- 2 Less than grade 10
- 3 Technical training after grade 7-8
- 4 Complete secondary (10-11 years)
- 5 Professional technical education after grade 10-11
- 6 Complete secondary + technical education
- 7 Incomplete postsecondary
- 8 Complete postsecondary

103. Are you still attending school?

- 1 Yes
- 2 No

104. What is your marital status (SHOW CARD)

- 1 Married
- 2 Living with a man (In unregistered marriage)
- 3 Separated
- 4 Divorced
- 5 Widowed
- 6 Never been in a registered or unregistered marriage----->**GO TO Q200**

105. How many times have you been in registered or unregistered marriages?

___ times

106. In what month and year did you begin living with your (first) husband/partner?

Month __ __

Year 19 __ __

107. What was the highest level of education that your (first) husband/partner completed? (SHOW CARD)

- 1 Less than grade 7
- 2 Less than grade 10
- 3 Technical training after grade 7-8
- 4 Complete secondary (10-11 years)
- 5 Professional technical education after grade 10-11
- 6 Complete secondary + technical education
- 7 Incomplete postsecondary
- 8 Complete postsecondary
- 77 Don't know/Don't remember

108. When you first got married/started living with a man how many children did you desire to have?

___ children

44 As many as we could afford

55 As many as possible

66 Up to God, fate, etc.

77 Were not sure

88 Other _____

II. FERTILITY/PREGNANCY

200. Are you currently pregnant?
- 1 Yes
 - 2 No----->GO TO Q204
 - 3 Not sure---->GO TO Q204
201. In what month of pregnancy are you?
- ___ month
- 7 7 Don't know
202. At the time you became pregnant, did you want to become pregnant then, want to wait longer to become pregnant or want no more children?
- 1 Wanted to become pregnant
 - 2 Wanted to wait longer---->**GO TO Q205**
 - 3 Wanted no more children---->**GO TO Q205**
 - 8 Don't know/Don't remember---->**GO TO Q205**
203. How long had you been trying to become pregnant?
(CONVERT YEARS TO MONTHS)
- ___ ___ ___ Months
- 6 6 6 Did not try
- 7 7 7 Don't remember
- GO TO Q205**
204. Have you ever been pregnant? (Including pregnancies that did not result in a live birth)
- 1 Yes
 - 2 No----->**GO TO Q266**
 - 7 Not sure----->**GO TO Q266**
205. Have you ever had any live-born children, regardless of how long they lived?
- 1 Yes
 - 2 No--->**GO TO Q210**
206. How many living children do you have, including those who do not live with you?
- ___ ___ children
207. In what month and year was your last baby born?
- Month ___ ___
- Year 19 ___ ___

IF CURRENTLY PREGNANT, CODE Q208 AS "YES" AND SKIP TO Q210.

208. Since that birth have you been pregnant again?

- 1 Yes
- 2 No---->**GO TO Q210**
- 3 Not sure---->**GO TO Q210**

209. How did your most recent pregnancy end? (SHOW CARD 209)

- 1 Stillbirth
- 2 Miscarriage
- 3 Induced abortion---->**GO TO PREGNANCY HISTORY, TABLE 1**
- 4 Miniabortion----->**GO TO PREGNANCY HISTORY, TABLE 1**
- 5 Ectopic pregnancy

210. Have you ever had any miniabortions, induced abortions, or done anything to terminate a pregnancy (even if it was a long time ago)?

- 1 Yes
- 2 No

TABLE 1: PREGNANCY HISTORY

Now I would like to talk to you about your past pregnancies. Please make sure you include all pregnancies, regardless of when they occurred and how they ended, whether in a live birth, an abortion, a miscarriage, or a stillbirth. Starting with your most recent pregnancy, please give me the following information: (IF CURRENTLY PREGNANT START WITH PREVIOUS PREGNANCY.)

211	212	213	214	215	216	217	218
LINE	When did this pregnancy end? (Month & yr)	When you got pregnant, did you want to become pregnant then, want to get pregnant later, or not want to have any more children?	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At what age did he/she die? (DAYS IF <1 MONTH, MONTH IF <2 YRS)
1	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 7 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
2	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 7 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
3	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 7 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
4	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 7 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
5	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 7 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years

211	212	213	214	215	216	217	218
LINE	When did this pregnancy end? (Month & year)	When you got pregnant, did you want to become pregnant then, want to get pregnant later, or not want to have any more children?	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At what age did he/she die?
6	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
7	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
8	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
9	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
10	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years

211	212	213	214	215	216	217	218
LINE	When did this pregnancy end? (Month & year)	When you got pregnant, did you want to become pregnant then, want to get pregnant later, or not want to have any more children?	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At what age did he/she die?
11	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
12	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
13	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
14	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
15	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years

211	21	213	214	215	216	217	218
LINE	When did this pregnancy end? (Month & yr)	When you got pregnant, did you want to become pregnant then, want to get pregnant later, or not want to have any more children?	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At what age did he/she die?
16	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
17	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
18	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
19	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->NEXT PREGNANCY 5 Miscarriage-->NEXT PREGNANCY 6 Induced Abortion->NEXT PREGNANCY 7 Miniabortion-->NEXT PREGNANCY 8 Self-induced abortion->NEXT PREG.	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years
20	Month ___ Year 19 ___ IF BEFORE 1994, GO TO Q214	1 Wanted to get pregnant then 2 Wanted to get pregnant later 3 Did not want any more children 8 Not sure/Don't remember	___ months 9=9+ 77=DK/DR	1 Live birth (single) --->Q216 2 Live births (multiple)--->Q216 3 Live birth + stillbirth (mult)---->Q216 4 Stillbirth--->q219 5 Miscarriage-->Q219 6 Induced Abortion->Q219 7 Miniabortion-->Q219 8 Self-induced abortion->Q219	1 Boy 2 Girl 3 Both	1 Alive->NEXT PREG 2 Dead 3 One living, One dead	___ Days ___ Months ___ Years

219. TOTAL NUMBER OF PREGNANCIES (INCLUDING CURRENT PREGNANCY) _____

220. TOTAL NUMBER OF ABORTIONS (CODES 6, 7, OR 8 FOR COLUMN 215) _____

IF ANY ABORTIONS SINCE THE BEGINNING OF 1994 GO TO Q221.**IF NO ABORTIONS SINCE THE BEGINNING OF 1994, BUT AT LEAST ONE LIVE BIRTH GO TO Q238.****IF NO LIVE BIRTHS OR INDUCED ABORTIONS SINCE THE BEGINNING OF 1994 GO TO INSTRUCTIONS BEFORE Q266.**

TABLE 2: QUESTIONS 221-230 ONLY FOR ABORTIONS THAT OCCURRED IN 1994 OR LATER

Now I would like to talk to you about the abortions you have had since the beginning of 1994, starting with the most recent abortion:

221	222	223	224	225	226	227	228	229	230
COPY LINE # FROM Q211	TYPE OF ABORTION (FROM Q215)	What was the main reason that you decided to have an abortion? (CODES BELOW)	In what type of facility was that abortion performed? (CODES BELOW) SHOW CARD	Was this a legal abortion	Soon after this abortion did you have any complications that required treatment?	What was the most serious complication that you experienced? (CODES BELOW) SHOW CARD	Did you stay in the hospital longer than expected or were you readmitted for this complication?	Did you have any related health problems more than 6 months later?	What was the most serious complication that you experienced? (CODES BELOW) (SHOW CARD)
___	1 MINIABORTION 2 REG. ABORTION 3 SELF INDUCED	1 2 3 4 5 6 7 77	1 2 3 4 5	1 Yes 2 No 7 DK	1 Yes--->Q227 2 No---->Q229 7 Dont know->Q229	1 2 3 4 5 6 7	1 Stayed longer 2 Readmitted 3 Both 4 No	1 Yes 2 No->NEXT LINE 7 Don't know--> NEXT LINE	1 2 3 4 5 6 7
___	1 MINIABORTION 2 REG. ABORTION 3 SELF INDUCED	1 2 3 4 5 6 7 77	1 2 3 4 5	1 Yes 2 No 7 DK	1 Yes--->Q227 2 No---->Q229 7 Dont know->Q229	1 2 3 4 5 6 7	1 Stayed longer 2 Readmitted 3 Both 4 No	1 Yes 2 No->NEXT LINE 7 Don't know--> NEXT LINE	1 2 3 4 5 6 7
___	1 MINIABORTION 2 REG. ABORTION 3 SELF INDUCED	1 2 3 4 5 6 7 77	1 2 3 4 5	1 Yes 2 No 7 DK	1 Yes--->Q227 2 No---->Q229 7 Dont know->Q229	1 2 3 4 5 6 7	1 Stayed longer 2 Readmitted 3 Both 4 No	1 Yes 2 No->NEXT LINE 7 Don't know--> NEXT LINE	1 2 3 4 5 6 7
___	1 MINIABORTION 2 REG. ABORTION 3 SELF INDUCED	1 2 3 4 5 6 7 77	1 2 3 4 5	1 Yes 2 No 7 DK	1 Yes--->Q227 2 No---->Q229 7 Dont know->Q229	1 2 3 4 5 6 7	1 Stayed longer 2 Readmitted 3 Both 4 No	1 Yes 2 No->NEXT LINE 7 Don't know--> NEXT LINE	1 2 3 4 5 6 7
___	1 MINIABORTION 2 REG. ABORTION 3 SELF INDUCED	1 2 3 4 5 6 7 77	1 2 3 4 5	1 Yes 2 No 7 DK	1 Yes--->Q227 2 No---->Q229 7 Dont know->Q229	1 2 3 4 5 6 7	1 Stayed longer 2 Readmitted 3 Both 4 No	1 Yes 2 No->NEXT LINE 7 Don't know-->NEXT LINE	1 2 3 4 5 6 7

CODES FOR Q223

- 1 Wanted no (more) children
- 2 Dangerous to her life/health
- 3 Risk of/Diagnosed fetal defect
- 4 Social/Econ/Preference reasons
- 5 Not married/No partner
- 6 Partner wanted abortion
- 7 Other (specify) _____
- 77 Don't know

CODES FOR Q224

- 1 Hospital/Polyclinic
- 2 Maternity house
- 3 Private clinic/physician
- 4 Not at a facility
- 5 Other _____
- 6 Other _____
- 7 Don't know

CODES FOR Q227

- 1 Perforation
- 2 Hemorrhage
- 3 Fever
- 4 Discharge
- 5 Pelvic pain
- 6 Other _____
- 7 Don't know

CODES FOR Q230

- 1 Pelvic pain
- 2 Sterility
- 3 Infection
- 4 Lack of menses
- 5 Irregular bleeding

IF MOST RECENT ABORTION WAS SELF-INDUCED (IF LINE 1, Q215=8) GO TO INSTRUCTIONS AFTER Q237.

IF MOST RECENT ABORTION WAS BEFORE 1997 (IF TABLE 1, LINE 1, Q215=8), GO TO Q232

231. How much did you pay (in Hryvnas) for all costs associated with your (most recent) abortion or miniabortion? This should include such costs as anesthesia, doctors' fees, blood tests and analysis, and any others. (CONVERT TO \$US, IF POSSIBLE)

___ ___ Hryvnas ___ ___ US Dollars ___ ___ German DM ___ ___ Russian Rubles

- 5 5 5 No charge
- 6 6 6 Nonmonetary payments (goods, services, etc.)
- 7 7 7 Don't remember

232. How many days did you spend in the place where you had your (most recent) abortion or minabortion?

- ___ ___ days
- 00 Less than a day
- 77 Don't remember

233. Did you receive local anesthesia or an injection so you would not feel pain during that abortion?

- 1 Yes
- 2 No
- 7 Don't remember

234. While you were in the hospital or clinic for your (most recent) abortion or miniabortion, did a doctor or nurse talk to you about ways to avoid another unplanned pregnancy?

- 1 Yes
- 2 No
- 7 Don't remember

235. Did a doctor or nurse refer you to another clinic or consultation for contraceptive counselling or services?

- 1 Yes
- 2 No---->**GO TO Q237**
- 7 Don't remember----->**GO TO Q237**

236. Did you go for those services?

- 1 Yes
- 2 No
- 7 Don't remember

237. After your (most recent) abortion or miniabortion, did you leave the clinic/hospital with a contraceptive method or a prescription for a contraceptive method?

- 1 Received contraceptive method
- 2 Received prescription for a contraceptive method
- 3 Received neither
- 7 Don't remember

IF RESPONDENT HAS HAD NO LIVE BIRTHS SINCE JANUARY 1994 GO TO Q266

IF RESPONDENT HAS HAD ANY LIVE BIRTHS SINCE JANUARY 1994 CONTINUE WITH Q238

TABLE 3:
QUESTIONS 238-249 ARE ONLY FOR LIVE BIRTHS SINCE JANUARY 1994.
IN COLUMN 238 FILL IN LINE NUMBERS FROM Q211 FOR ALL LIVE BIRTHS SINCE JANUARY 1994.

I would like to ask you some questions about your children born since 1994. Starting with the most recent child, please tell me:

238	239	240	241	242	243	244	245	46	2	47	2	8	24	249
COPY LINE # FROM Q211	Did you receive any prenatal care from a doctor, nurse or midwife for this pregnancy?	During what month of pregnancy did you make your first prenatal care visit?	How many prenatal care visits did you make altogether during that pregnancy?	Where did you receive most of your prenatal care?	During that pregnancy were you ever hospitalized for any problems?	How many days were you in the hospital for these problems?	Was this delivery by cesarean section?	Where did that delivery take place?	Did you breastfeed him/her?	Are you still breastfeeding him/her?			At what age did he/she start receiving foods or liquids other than breast milk?	
___	1 Yes----->Q240 2 No----->Q243 7 Not sure--->Q243	___ 77=Don't know	___ visits 55=As many as Dr. said 66=DR, but at least 10 77=Don't remember	1 Maternity house 2 Women's consult. 3 MCH Center 4 Hospital 5 Priv. clinic/office 6 Other _____	1 Yes 2 No--> GO TO Q245	___ days 77=don't rem. 99=99+ days	1 Yes 2 No	1 Mat House 2 MCH Ctr. 3 Hospital 4 Home 5 Other___	1 Yes 2 No--> NEXT LINE	1 Yes 2 No		___ months 00=<1 month 66=Not yet 77=Don't rem.		
___	1 Yes----->Q240 2 No----->Q243 7 Not sure--->Q243	___ 77=Don't know	___ visits 55=As many as Dr. said 66=DR, but at least 10 77=Don't remember	1 Maternity house 2 Women's consult. 3 MCH Center 4 Hospital 5 Priv. clinic/office 6 Other _____	1 Yes 2 No--> GO TO Q245	___ days 77=don't rem. 99=99+ days	1 Yes 2 No	1 Mat House 2 MCH Ctr. 3 Hospital 4 Home 5 Other___	1 Yes>NEXT LINE 2 No>Q249			___ months 00=<1 month 66=Not yet 77=Don't rem.		
___	1 Yes----->Q240 2 No----->Q243 7 Not sure--->Q243	___ 77=Don't know	___ visits 55=As many as Dr. said 66=DR, but at least 10 77=Don't remember	1 Maternity house 2 Women's consult. 3 MCH Center 4 Hospital 5 Priv. clinic/office 6 Other _____	1 Yes 2 No--> GO TO Q245	___ days 77=don't rem. 99=99+ days	1 Yes 2 No	1 Mat House 2 MCH Ctr. 3 Hospital 4 Home 5 Other___	1 Yes>NEXT LINE 2 No>Q249			___ months 00=<1 month 66=Not yet 77=Don't rem.		
___	1 Yes----->Q240 2 No----->Q243 7 Not sure--->Q243	___ 77=Don't know	___ visits 55=As many as Dr. said 66=DR, but at least 10 77=Don't remember	1 Maternity house 2 Women's consult. 3 MCH Center 4 Hospital 5 Priv. clinic/office 6 Other _____	1 Yes 2 No--> GO TO Q245	___ days 77=don't rem. 99=99+ days	1 Yes 2 No	1 Mat House 2 MCH Ctr. 3 Hospital 4 Home 5 Other___	1 Yes>NEXT LINE 2 No>Q249			___ months 00=<1 month 66=Not yet 77=Don't rem.		
___	1 Yes----->Q240 2 No----->Q243 7 Not sure--->Q243	___ 77=Don't know	___ visits 55=As many as Dr. said 66=DR, but at least 10 77=Don't remember	1 Maternity house 2 Women's consult. 3 MCH Center 4 Hospital 5 Priv. clinic/office 6 Other _____	1 Yes 2 No--> GO TO Q245	___ days 77=don't rem. 99=99+ days	1 Yes 2 No	1 Mat House 2 MCH Ctr. 3 Hospital 4 Home 5 Other___	1 Yes>NEXT LINE 2 No>Q249			___ months 00=<1 month 66=Not yet 77=Don't rem.		

249A. NUMBER OF PREGNANCIES ENDING IN LIVE BIRTHS SINCE JANUARY 1994 ___

**THE FOLLOWING QUESTIONS DEAL WITH THE MOST RECENT PREGNANCY
THAT LED TO A LIVE BIRTH SINCE 1/94.**

**IF RESPONDENT RECEIVED NO PRENATAL CARE FOR LAST BIRTH (SEE Q239, LINE 1)
GO TO Q255.**

250. During these visits did you have your blood pressure measured?

- 1 Yes
- 2 No----->**GO TO Q252**
- 7 Don't remember----->**GO TO Q252**

251. During these visits were you ever told that you had high blood pressure?

- 1 Yes
- 2 No
- 7 Don't remember

252. Were you ever told you had anemia during that pregnancy?

- 1 Yes
- 2 No
- 7 Don't know/Don't remember

253. Did you take iron supplements during that pregnancy?

- 1 Yes
- 2 No
- 7 Don't know/Don't remember

254. Did you have any ultrasound exams during your pregnancy?

- 1 Yes
- 2 No
- 7 Don't remember

255. Did you smoke cigarettes at the time you found out you were pregnant?

- 1 Yes
- 2 No--->**GO TO Q257**

256. Did you continue to smoke cigarettes during that pregnancy?

- 1 Yes
- 2 No

257. About how many times per week did you drink alcoholic beverages during that pregnancy?

- 1 At least 4 times per week
- 2 1-3 times per week
- 3 Less than once per week
- 4 Never
- 7 Don't remember
- 9 No response

258. How much did your last baby weigh when he/she was born? (IF SHE DOESN'T KNOW THE EXACT WEIGHT, FIND OUT IF THE BABY WAS AT LEAST 2,500 GRAMS.)
- ___ ___ ___ grams
 5 5 5 5= Don't know, but <2,500 grams
 6 6 6 6= Don't know, but at least 2,500 grams
 7 7 7 7= Don't know/Don't remember
259. During the first six weeks after birth, did you have any post-natal care visits to check on your health?
- 1 Yes
 2 No
 7 Don't remember
260. Before you left the health facility following your most recent birth did a doctor or nurse talk to you about or offer to talk to you about contraception?
- 1 Yes
 2 No
 7 Don't remember
261. During your hospitalization, did a doctor or nurse refer you to another clinic or consultation for contraceptive counselling or services?
- 1 Yes
 2 No---->**GO TO Q263**
 7 Don't remember----->**GO TO Q263**
262. Did you go for those services?
- 1 Yes
 2 No
 7 Don't remember
263. Before you left the hospital , did you receive a contraceptive method or a prescription for a contraceptive method?
- 1 Received a contraceptive method
 2 Received a prescription for a contraceptive method
 3 Received neither
 7 Don't remember
264. How many months after birth did your menstrual period return?
- ___ ___ Months
 00 = <1 month
 66 = Not yet
 77 = Don't remember
265. How many months after birth did you resume sexual relations?
- ___ ___ Months
 00=<1 month
 66 = Not yet
 77 = Don't remember

266. Have you ever tried to become pregnant?

1. Yes
2. No----->**GO TO Q300**

266A. Have you ever had a problem becoming pregnant when you wanted to?

1. Yes
2. No----->**GO TO Q300**
7. Don't remember----->**GO TO Q300**

267. How long ago did you start trying to become pregnant the last time this happened?

- ___ ___ Years ago
- 0 0 Less that 1 year ago
 - 7 7 Not sure/ Don't remember----->**GO TO Q300**

IF MORE THAN 10 YEARS GO TO Q300

268. How long did you try to become pregnant that time (IF ANSWER IS IN YEARS, CONVERT IT TO MONTHS.)

- ___ ___ ___ Months
- 1 1 1 Still trying
 - 6 6 6 Don't remember, but at least 2 years
 - 7 7 7 Don't remember

269. Did you or your partner seek medical treatment to help you become pregnant that time?

- 1 Respondent sought treatment
- 2 Partner sought treatment
- 3 Both sought treatment
- 4 No----->**GO TO Q275**

270. Where was the first place you/your partner sought help in getting pregnant? (SHOW CARD)

- 1 Women's consultation
- 2 MCH Center
- 3 Family planning center
- 4 Other public facility (specify)_____
- 5 Private clinic/physician
- 6 Private fertility clinic
- 7 Other in Ukraine (specify)_____
- 8 Outside Ukraine

271. Did you/your partner seek help anywhere else?

- 1 Yes
- 2 No----->**GO TO Q273**

272. Where was the most recent place? (SHOW CARD)
- 1 Women's consultation
 - 2 MCH Center
 - 3 Family planning center
 - 4 Other public facility (specify)_____
 - 5 Private clinic/physician
 - 6 Private fertility clinic
 - 7 Other in Ukraine (specify)_____
 - 8 Outside Ukraine
273. What was the reason you were given for your inability to become pregnant? (SHOW CARD)
- 1 Woman's hormone problems
 - 2 Man's hormone problems
 - 3 Problem with man's sperm
 - 4 Anatomical problem of the woman (e.g., blocked tubes)
 - 5 Complications from previous pregnancy
 - 6 Complications from previous abortion
 - 7 Pelvic inflammatory disease
 - 8 Complications from sexually transmitted disease
 - 9 Other (specify)_____
 - 66 No reason given
 - 77 Don't know/Don't remember
274. What types of treatment(s) did you/your partner receive? Mention all that you remember. (SHOW CARD)
1. Hormones
 2. Anti-Inflammatory drugs
 3. Treatment for Fallopian blockage
 4. Physiotherapy
 5. Laser therapy
 6. Laparoscopy/Microsurgery
 7. Spa/Relaxation therapy, etc.
 8. In Vitro Fertilization (IVF)
 9. Other_____
 10. No treatment
275. Were you finally able to become pregnant?
- 1 Yes
 - 2 No----->**GO TO Q300**
276. How did that pregnancy end? (SHOW CARD)
- 1 Currently pregnant
 - 2 Live birth
 - 3 Stillbirth
 - 4 Miscarriage
 - 5 Ectopic pregnancy
 - 6 Abortion
 - 7 Miniabortion

III. CONTRACEPTION

TABLE 4:

For each of the following methods of preventing pregnancy, please tell me:

METHOD	300. Have you ever heard of it?	301. Have you/ partner ever used it?	302. Do you know where to get it?
A. Female Sterilization (Tubal ligation)	1 Yes-->Q301 2 No--->B	1 Yes-->B 2 No--->Q302	1 Yes 2 No
B. Male Sterilization (Vasectomy)	1 Yes-->Q301 2 No--->C	1 Yes-->C 2 No--->Q302	1 Yes 2 No
C. Oral contraceptives (Pills)	1 Yes-->Q301 2 No--->D	1 Yes-->D 2 No--->Q302	1 Yes 2 No
D. IUD	1 Yes-->Q301 2 No--->E	1 Yes-->E 2 No--->Q302	1 Yes 2 No
E. Depo-Provera/ Injections	1 Yes-->Q301 2 No--->F	1 Yes-->F 2 No--->Q302	1 Yes 2 No
F. Implants/ Norplant	1 Yes-->Q301 2 No--->G	1 Yes-->G 2 No--->Q302	1 Yes 2 No
G. Condoms	1 Yes-->Q301 2 No--->H	1 Yes-->H 2 No--->Q302	1 Yes 2 No
H. Spermicide/Cream/Foam	1 Yes-->Q301 2 No--->I	1 Yes-->I 2 No--->Q302	1 Yes 2 No
I. Diaphragm/Cervical cap	1 Yes-->Q301 2 No--->J	1 Yes-->J 2 No--->Q302	1 Yes 2 No
			302. Do you know where to get information on it?
J. Safe period methods (Rhythm, etc.)	1 Yes-->Q301 2 No--->K	1 Yes-->K 2 No--->Q302	1 Yes 2 No
K. Withdrawal	1 Yes-->Q301 2 No--->L	1 Yes-->L 2 No--->L	
L. Any other method _____	1 Yes-->Q301 2 No-->INSQ303	1 Yes 2 No	

IF RESPONDENT HAS USED ANY METHOD (ANY "YES" FOR Q301) GO TO INSTRUCTION BEFORE Q304

303. So, you have never used any method or done anything to prevent pregnancy with any partner?

- 1 Never used----->**GO TO Q318**
2 Ever used--->**CORRECT Q301, THEN GO TO Q304**

IF CURRENTLY PREGNANT GO TO Q318

304. Are you (or your partner) currently using any method or doing anything to prevent pregnancy?

- 1 Yes
2 No--->**GO TO Q318**

305. What method or methods are you using?

- 1 Tubal ligation----->**GO TO Q306**
- 2 Vasectomy----->**GO TO Q307**
- 3 Pills----->**GO TO Q308**
- 4 IUD----->**GO TO Q308**
- 5 DepoProvera/Injections---->**GO TO Q308**
- 6 Implants/Norplant----->**GO TO Q308**
- 7 Condoms----->**GO TO Q308**
- 8 Spermicide/Cream/Jelly---->**GO TO Q308**
- 9 Diaphragm/Cervical cap---->**GO TO Q308**
- 10 Morning-after pill----->**GO TO Q308**
- 11 Lact. Amenorrhea Meth---->**GO TO Q311**
- 12 Safe period method----->**GO TO Q311**
- 13 Withdrawal----->**GO TO Q311**
- 14 Douching----->**GO TO Q311**
- 15 Condoms + Spermicide---->**GO TO Q308**
- 16 Other combination including condoms _____ ---->**GO TO Q308**
- 17 Other combination not including condoms _____ ---->**GO TO Q308**
- 18 Other method _____ ---->**GO TO Q311**

306. Was this operation done during hospitalization for a delivery or abortion?

- 1 Yes, after delivery
- 2 Yes, after abortion
- 3 No

307. In what month and year was this operation performed?

Month __ __ Year 19 __ __

308. Where do/did you get your family planning method? (MOST RECENT SOURCE OF SUPPLY) (SHOW CARD)

- 1 Women's consultation
- 2 Maternity house
- 3 MCH center
- 4 Hospital
- 5 Pharmacy
- 6 Drug kiosk
- 7 Private clinic/Physician
- 8 Commercial kiosk
- 9 Store
- 10 Family planning center
- 11 Outside Ukraine
- 12 Other _____
- 77 Don't know/Don't remember

309. Did you (or your partner or relative) pay for this method the last time?

- 1 Yes
- 2 No----->**GO TO Q313**

310. How much did you pay? (CONVERT SUM INTO \$US IF GIVEN IN OTHER CURRENCY)

- __ __ __ DOLLARS
- 6 6 6= Payment in goods or services
 - 7 7 7= Don't remember

GO TO Q313

311. Q311 TO Q312 FOR WOMEN NOW USING WITHDRAWAL, A SAFE PERIOD METHOD, OR DOUCHING TO AVOID PREGNANCY

You said that you are now using _____ (ANSWER IN Q305) to avoid becoming pregnant, rather than a method you might obtain from a doctor, health facility, or pharmacy. Please tell me whether each of the following was very important, somewhat important, or not important at all in your decision to use this method: (SHOW CARD)

	Very Important	Somewhat Important	Not Important	Not Sure
A. Difficulty of getting other methods	1	2	3	7
B. Cost of other methods	1	2	3	7
C. Lack of information about other methods	1	2	3	7
D. Health/Side effects of other methods	1	2	3	7
E. Husband/Partner preference	1	2	3	7
F. Religious beliefs	1	2	3	7
G. Naturalness of the method	1	2	3	7

312. Do you think that _____ (current method) is better than worse than or about the same as methods received from a doctor or pharmacy, like the IUD or pills, at preventing pregnancies?

(READ CHOICES)

- 1 Current method more effective
- 2 About equally effective
- 3 Current method less effective
- 7 Don't know/Not sure

313. Do you currently have any important problems or concerns with using _____ (current method)?

- 1 Yes
- 2 No--->GO TO Q315

314. What is the most important problem or concern with using this method? (SHOW CARD)

- 1 Side effects
- 2 Health concerns
- 3 Cost
- 4 Access/Availability
- 5 Effectiveness of method for preventing pregnancy
- 6 Sometimes forget to use
- 7 Sometimes difficult/inconvenient to use
- 8 Husband/partner disapproves
- 9 Irreversible/Doctor needed to discontinue
- 10 Other (specify)_____

315. Would you prefer to use a different method of preventing pregnancy from the one you are currently using?

- 1 Yes
- 2 No--->GO TO INSTRUCTIONS PRECEDING Q318
- 7 Don't know----> GO TO INSTRUCTIONS PRECEDING Q318

316. What method would you prefer to use? (SHOW CARD)

- 1 Tubal ligation
- 2 Vasectomy
- 3 Pills
- 4 IUD
- 5 DepoProvera/Injections
- 6 Implants/Norplant
- 7 Condoms
- 8 Spermicide/Cream/Jelly
- 9 Diaphragm/Cervical cap
- 10 Morning-after pill
- 11 Lact. Amenorrhea Meth
- 12 Safe period method
- 13 Withdrawal
- 14 Douching
- 15 Condoms + Spermicide
- 16 Other combination including condoms_____
- 17 Other combination not including condoms_____
- 18 Other method_____
- 77 Not sure----->**GO TO Q318**

317. What is the most important reason that you do not use that method now?

- 1 Doctor will not prescribe it
- 2 Cost
- 3 Difficult to get/Must go too far to get it
- 4 Don't know enough about method
- 5 Do not know how to obtain it
- 6 Husband objects to it
- 7 Religious reasons
- 8 Fear of health problems
- 9 Fear of side effects
- 10 Poor effectiveness
- 11 Current method is permanent/long-term
- 12 Method is not natural
- 13 Other_____
- 77 Don't know

IF RESPONDENT OR HER PARTNER HAS BEEN STERILIZED, GO TO Q321 (CALENDAR)

318. Do you plan to have any (more) children (after this pregnancy)?

- 1 Yes
- 2 No----->**GO TO Q321**
- 7 Not sure----->**GO TO Q321**

319. How many (more) do you plan to have?

- ___ children
- 55 As many as possible
 - 66 Up to God/Fate, etc.
 - 77 Not sure

320. When do you think you would like to become pregnant? (SHOW CARD)

- 1 As soon as possible/Now
- 2 Within 1 year
- 3 In 1-2 years
- 4 In 3-5 years
- 5 In more than 5 years
- 6 When/After I get married
- 7 Not sure/Don't know

321. **CONTRACEPTIVE USE/PREGNANCY CALENDAR:**

(INTERVIEWER: FILL IN ALL PREGNANCIES AND BIRTHS FROM COLUMN 1 OF THE PREGNANCY HISTORY BEFORE COLLECTING THE CONTRACEPTIVE HISTORY).

Starting at the beginning of 1994, please try to remember:

321.1: in which months you were pregnant and when and how the pregnancy ended. (CARD)

321.2: in which months you started and stopped use of contraceptive methods. (CARD)

321.3: why you stopped using the method. (CARD)

COLUMN 1

PREGNANCY

- 0 Not pregnant
- 1 Pregnant
- 2 Live birth
- 3 Stillbirth
- 4 Miscarriage
- 5 Miniabortion
- 6 Regular abortion

COLUMN 2

CONTRACEPTION

- 0 No method
- 1 Pills
- 2 IUD
- 3 Condoms
- 4 Injections
- 5 Safe period methods
- 6 Withdrawal
- 7 Sterilization
- 8 Other _____
- 9 Don't remember

COLUMN 3

REASON STOPPED USING

- 1 Pregnant while using method
- 2 Wanted to get pregnant
- 3 Husband objected
- 4 Side effects
- 5 Health concerns
- 6 Physician decision
- 7 Supply/Availability/Cost
- 8 Inconvenient method
- 9 Want better/more effective method
- 10 To give body a rest
- 11 Infrequent/No sex
- 12 Marriage/Relationship ended
- 13 Could no longer get pregnant
- 14 Other _____
- 77 Don't remember

TE	DA	1	2	3	TE	DA	1	2	3
1994					1997				
Jan	1				Jan	1			
Feb	2				Feb	2			
Mar	3				Mar	3			
Apr	4				Apr	4			
May	5				May	5			
Jun	6				Jun	6			
Jul	7				Jul	7			
Aug	8				Aug	8			
Sep	9				Sep	9			
Oct	10				Oct	10			
Nov	11				Nov	11			
Dec	12				Dec	12			
1995					1998				
Jan	1				Jan	1			
Feb	2				Feb	2			
Mar	3				Mar	3			
Apr	4				Apr	4			
May	5				May	5			
Jun	6				Jun	6			
Jul	7				Jul	7			
Aug	8				Aug	8			
Sep	9				Sep	9			
Oct	10				Oct	10			
Nov	11				Nov	11			
Dec	12				Dec	12			
1996					1999				
Jan	1				Jan	1			
Feb	2				Feb	2			
Mar	3				Mar	3			
Apr	4				Apr	4			
May	5				May	5			
Jun	6				Jun	6			
Jul	7				Jul	7			
Aug	8				Aug	8			
Sep	9				Sep	9			
Oct	10				Oct	10			
Nov	11				Nov	11			
Dec	12				Dec	12			

IF NOT USING A METHOD IN JANUARY 1994 GO TO INSTRUCTIONS BEFORE Q323

322. You said you were using _____ in January of 1994. When did you start using that method that time?
 Month ___ Year 19__

IF WOMAN HAS NOT HAD AN IUD INSERTED SINCE JANUARY 1994, GO TO INSTRUCTIONS BEFORE Q331.

323. You said you (last) had an IUD inserted in (CHECK CALENDAR):

Month __ __ Year 19 __ __

Is that correct? **(IF NOT CORRECT, CHANGE CALENDAR AND Q323.)**

IF NOT CURRENTLY WEARING AN IUD, GO TO Q326

324. Since it was inserted, has the IUD been checked by a physician or health worker to make sure it was in place?

- 1 Yes
- 2 No--->**GO TO Q326**
- 7 Don't know--->**GO TO Q326**

325. How long ago was the last such check?

- __ __ Months ago
- 66 More than 2 years ago
- 77 Don't remember

326. Was your IUD inserted immediately (within one week) after a delivery, abortion, or miniabortion?

- 1 Yes, after delivery
- 2 Yes, after abortion/miniabortion
- 3 No

327. When your IUD was inserted, how long did the physician tell you it could be left in?

- __ __ years
- 44 As long as I wanted
- 55 Other (specify) _____
- 66 Did not say how long
- 77 Don't remember

328. Did you have any health problems or side effects that you think are related to your IUD within one year of insertion?

- 1 Yes
- 2 No--->**GO TO INSTRUCTIONS BEFORE Q331**
- 7 Not sure--->**GO TO INSTRUCTIONS BEFORE Q331**

329. What kind of problem did you have? (IF MORE THAN ONE PROBLEM, CODE THE MOST SERIOUS ONE.) (SHOW CARD)

- 1 Cramping
- 2 Heavy bleeding during menstrual periods
- 3 Bleeding/Spotting between menstrual periods
- 4 Inflammation/Infection/Discharge
- 5 Husband/Partner complained about string
- 6 Other (specify) _____

330. Was the problem serious enough that you went to a doctor or clinic about it?

- 1 Yes
- 2 No
- 7 Don't remember

IF RESPONDENT DID NOT TAKE ORAL CONTRACEPTIVES SINCE JANUARY 1994, GO TO INSTRUCTION BEFORE Q337

331. You said you most recently started taking oral contraceptives in (CHECK CALENDAR):

Month ___ Year 19 ___

Is that correct? (IF NOT CORRECT, CHANGE CALENDAR AND Q331.)

IF NOT CURRENTLY TAKING ORAL CONTRACEPTIVES, GO TO Q333

332. What brand of pills do you currently use? (ASK TO SEE PACKAGE, IF AVAILABLE)

- | | | | |
|-------------|-------------------|--------------|-------------------------|
| 1 Marvelon | 9 Tri-Regol | 17 Bicecurin | 24 Obusmen |
| 2 Cilest | 10 Regividon | 18 LoFeminol | 25 Conceplan M |
| 3 Micronor | 11 Antiovin | 19 Demulen | 26 Kliogest N |
| 4 Trinovum | 12 Ovidur-Richter | 20 Ovulen-50 | 27 Continuin (minipill) |
| 5 Triqvilar | 13 Trisiston | 21 Non-Ovlon | 28 Other_____ |
| 6 Femoden | 14 Diane-35 | 22 Egestriol | 77 Don't know |
| 7 Milvanar | 15 Ovidon | 23 Exluton | |
| 8 Ovrette | 16 Postinor | | |

333. When you started taking pills, how long did your physician tell you that you could take them?

___ Years

- 44 As long as I wanted/Indefinitely
 55 Other (specify)_____
 66 Did not say how long/Never talked about it
 77 Don't remember/Don't know

334. During the first year you took pills did you have any health problems or side effects you think were related to using your pills?

- 1 Yes
 2 No--->GO TO INSTRUCTIONS BEFORE Q337
 7 Don't remember--->GO TO INSTRUCTIONS BEFORE Q337

335. What was the worst problem you had? (SHOW CARD)

- 1 Headaches
 2 Vision problems
 3 Bloating/Weight gain
 4 Nausea
 5 Bleeding between menstrual periods
 6 Other (specify)_____

336. Was this problem serious enough that you went to a doctor or clinic about it?

- 1 Yes
 2 No
 7 Don't remember

IF CURRENTLY USING ANY CONTRACEPTIVE METHOD (SEE Q304) GO TO INSTRUCTIONS BEFORE Q342

IF CURRENTLY PREGNANT (SEE Q200) GO TO INSTRUCTIONS BEFORE Q342

337. Do you think you are able to get pregnant at the present time?

- 1 Yes--->GO TO Q339
 2 No--->
 7 Not sure---->GO TO Q339

338. Why not?
- 1 Menopause/No menstrual periods \
 - 2 Has had an operation for medical reasons \ that makes pregnancy impossible \
 - 3 Husband/partner has had a medical operation \
 - 4 Has not gotten pregnant despite |----->**GO TO INSTRUCTIONS BEFORE Q347**
At least 2 years not contracepting /
 - 5 Doctor says she or partner is infertile /
 - 6 Not sexually active
 - 7 Postpartum/Breastfeeding
 - 8 Other (specify)_____

339. Are you trying to become pregnant now?

- 1 Yes
- 2 No--->**GO TO Q341**

340. How long have you been trying to become pregnant?
(IF ANSWER IS IN YEARS CONVERT TO MONTHS)

- ___ ___ months
- 5 5 5 = At least 10 years
 - 6 6 6 = Other_____
 - 7 7 7 = Don't remember

GO TO INSTRUCTIONS BEFORE Q342

341. What is the most important reason you are not using a method to avoid pregnancy now?

- 1 Want to become pregnant
- 2 Not sexually active/No partner
- 3 Only occasionally sexually active
- 4 Breastfeeding/Postpartum
- 5 Fear of side effects
- 6 Previous side effects
- 7 Fear of health effects
- 8 Husband/Partner objects
- 9 Religious reasons
- 10 Doctor's recommendation/Doctor won't prescribe method
- 11 Desired method not available/difficult to get
- 12 Too expensive
- 13 Don't know where to get method
- 14 Methods difficult to use
- 15 Prefer abortion
- 16 Haven't bothered, but would like to use method
- 17 Too old
- 18 Difficulty getting pregnant
- 19 Other (specify)_____
- 77 Not sure/Don't know

IF THE RESPONDENT PLANS TO HAVE ANY (MORE) CHILDREN (Q318=1) GO TO INSTRUCTIONS BEFORE Q344.

IF RESPONDENT OR PARTNER HAVE BEEN STERILIZED (Q305=1 OR 2) GO TO INSTRUCTIONS BEFORE Q347.

342. Would you be interested in an operation (sterilization) to prevent you from having any more children (after this pregnancy)?

- 1 Yes---->**GO TO INSTRUCTIONS BEFORE Q344**
- 2 No
- 7 Not sure---->**GO TO INSTRUCTIONS BEFORE Q344**

343. What is the most important reason you would not be/are not interested?

- 1 Health risks
- 2 Fear of operation
- 3 Husband/Partner would object
- 4 Religious reasons
- 5 Not culturally/socially acceptable
- 6 Cost of an operation
- 7 Might want another child
- 8 Don't know enough about sterilization
- 9 Haven't thought about it
- 10 No partner/Not sexually active
- 11 Other_____
- 77 Don't know

IF NOT MARRIED OR LIVING WITH A MAN (Q104=3-6) GO TO INSTRUCTIONS BEFORE Q347

344. Did you and your husband/partner ever talk about the number of children you wanted to have.

- 1 Yes
- 2 No

345. Have you and your husband/partner ever discussed using family planning?

- 1 Yes
- 2 No

346. Who should make decisions about whether to use contraception: the woman, the man, or both together?

- 1 The woman
- 2 The man
- 3 Both
- 7 Not sure

IF THE RESPONDENT HAS NEVER HAD AN ABORTION GO TO THE INSTRUCTIONS BEFORE Q349.

347. Does your husband/partner know about your (most recent) abortion?

- 1 Yes
- 2 No
- 3 Was not my partner at that time---->**GO TO INSTRUCTIONS BEFORE Q349**
- 7 Don't know

348. Did you discuss with him whether to get an abortion that time?

- 1 Yes
- 2 No
- 7 Don't remember

CONTRACEPTIVE COUNSELLING

IF RESPONDENT HAS NOT USED ORAL CONTRACEPTIVES, IUD OR INJECTIONS SINCE 1/94 (SEE CALENDAR, COLUMN 2) GO TO Q400.

349. **CURRENT OR MOST RECENT OF THESE METHOD**

1 ORAL CONTRACEPTIVES

2 IUD

3 INJECTIONS

350. The last time you started using oral contraceptives/an IUD/injections, did a health provider talk to you about various methods of family planning and the most appropriate method for you?

1 Yes

2 No---->**GO TO Q352**

7 Don't remember---->**GO TO Q352**

351. Who provided this counselling? (CIRCLE ONE ANSWER ONLY) (SHOW CARD)

1 Physician

2 Nurse

3 Midwife

4 Other health professional_____---->**GO TO Q400**

5 Other (not a health professional)_____---->**GO TO Q400**

7 Don't remember

352. Did you select the method you received, was it selected by the provider, or was it selected by both of you?

1 Respondent

2 Provider

3 Both

7 Don't remember

353. Did the provider explain the possible side effects of the method?

1 Yes

2 No

3 Received no counselling----->**GO TO Q400**

7 Don't remember

354. How easily could you understand the information concerning use of the method and its possible side effects? (SHOW CARD)

1 Did not understand at all

2 Mostly did not understand

3 Understood about half

4 Mostly understood

5 Completely understood

7 Don't remember

355. Did the provider explain to you when to return for removal, refill, or follow-up?

1 Yes

2 No

7 Don't remember

356. Did you receive a pelvic examination before being given the method?

1 Yes

2 No

7 Don't remember

IV. OPINIONS, ATTITUDES, IEC

400. How often do you watch television? (SHOW CARD)

- 1 Every day or almost every day
- 2 At least once per week
- 3 At least once per month
- 4 Less than once per month----->GO TO Q405
- 5 Hardly ever/Never----->GO TO Q405
- 6 Other (specify)_____

401. What times do you most often watch television? (SHOW CARD)

- 1 6:00-8:00
- 2 8:00-10:00
- 3 10:00-12:00
- 4 12:00-14:00
- 5 14:00-16:00
- 6 16:00-18:00
- 7 18:00-20:00
- 8 20:00-22:00
- 9 After 22:00
- 10 No regular times

402. What types of programs do you most often watch? (SHOW CARD)

- 1 News
- 2 Entertainment programs
- 3 Soap operas
- 4 Sports
- 5 Children's programs
- 6 Plays/Dramas
- 7 Church/Religious programs
- 8 Women's programs
- 9 Health programs
- 10 Political events
- 11 Business programs
- 12 Music programs, videos
- 13 Other (specify)_____

403. Within the past 6 months have you seen anything on television about family planning or methods of preventing pregnancy?

- 1 Yes
- 2 No
- 7 Not sure

404. Within the past 6 months have you seen anything on television about sexually transmitted diseases?

- 1 Yes
- 2 No
- 7 Not sure

405. How often do you listen to the radio? (SHOW CARD)

- 1 Every day or almost every day
- 2 At least once per week
- 3 At least once per month
- 4 Less than once per month----->**GO TO Q408**
- 5 Hardly ever/Never----->**GO TO Q408**
- 6 Other (specify)_____

406. What times do you most often listen to the radio? (SHOW CARD)

- 1 6:00-8:00
- 2 8:00-10:00
- 3 10:00-12:00
- 4 12:00-14:00
- 5 14:00-16:00
- 6 16:00-18:00
- 7 18:00-20:00
- 8 20:00-22:00
- 9 After 22:00
- 10 No regular times

407. What types of programs do you most often listen to? (SHOW CARD)

- 1 News
- 2 Personal announcements
- 3 Commercials
- 4 Sports
- 5 Children's programs
- 6 Plays/Dramas
- 7 Church/Religious programs
- 8 Women's programs
- 9 Health programs
- 10 Political events
- 11 Business programs
- 12 Music programs
- 13 Other (specify)_____

408. Do you think information about methods of preventing pregnancy should be broadcast on radio and television?

- 1 Yes
- 2 No
- 7 Not sure

409. How often do you read a daily newspaper? (SHOW CARD)

- 1 Daily/Nearly every day
- 2 About 3-4 times per week
- 3 Once or twice per week
- 4 Less than once per week
- 5 Never/Almost never--->**GO TO Q411**

410. Which newspaper(s) do you read most often?

- 1 Facti
- 2 Vceukrainsky Vegemosty
- 3 Golos Ukraini
- 4 Business
- 5 Uryadovi Kur'er
- 6 Silski Visty
- 7 Kiev Vedomosti
- 8 Telenedelya
- 9 Other (specify)_____

411. Within the past six months have you seen any newspapers or magazines that contained family planning information?

- 1 Yes
- 2 No
- 7 Don't remember

On a scale from 1 to 5, please rate each of the following contraceptive methods according to each of the characteristics I will mention:

412. How would you rate each of the following methods with regard to safety and health effects?
(5=completely safe, 1=extremely unsafe)

Oral contraceptives	1	2	3	4	5	7
IUD	1	2	3	4	5	7
Injections	1	2	3	4	5	7
Condoms	1	2	3	4	5	7
Female sterilization	1	2	3	4	5	7
Induced abortion	1	2	3	4	5	7
Miniabortion	1	2	3	4	5	7

413. How would you rate each of the following methods with regard to effectiveness at preventing pregnancy?
(5=completely effective, 1=completely ineffective)

Oral contraceptives	1	2	3	4	5	7
IUD	1	2	3	4	5	7
Injections	1	2	3	4	5	7
Condoms	1	2	3	4	5	7
Female sterilization	1	2	3	4	5	7

414. How would you rate each of the following methods with regard to cost?
(5=very inexpensive, 1=very expensive)

Oral contraceptives	1	2	3	4	5	7
IUD	1	2	3	4	5	7
Injections	1	2	3	4	5	7
Condoms	1	2	3	4	5	7
Female sterilization	1	2	3	4	5	7
Induced abortion	1	2	3	4	5	7
Miniabortion	1	2	3	4	5	7

415. Overall, how much do you like each of the methods of preventing births?
(5=like very much, 1=dislike very much)

Oral contraceptives	1	2	3	4	5	7
IUD	1	2	3	4	5	7
Injections	1	2	3	4	5	7
Condoms	1	2	3	4	5	7
Female sterilization	1	2	3	4	5	7
Induced abortion	1	2	3	4	5	7
Miniabortion	1	2	3	4	5	7

416. Please indicate whether you agree or disagree with the following statements about birth control pills: (SHOW CARD)

5=strongly agree
4=mildly agree
3=partially agree, partially disagree
2=mildly disagree
1=strongly disagree
7=don't know

1. They make women gain weight	1	2	3	4	5	7
2. They make women's periods more regular	1	2	3	4	5	7
3. Taking them too long can cause infertility	1	2	3	4	5	7
4. Women who take them have a higher risk of getting cancer	1	2	3	4	5	7
5. They are bad for blood circulation	1	2	3	4	5	7

417. When is it most likely for a woman to become pregnant? (SHOW CARD)

1 In the week before menstruation starts
2 During menstruation
3 In the week after menstruation ends
4 Halfway between her periods
5 It doesn't matter, all times are alike
6 Other (specify) _____
7 Don't know

418. Do you think it is harmful to the baby if a woman smokes while she is pregnant?

1 Yes
2 No
3 Depends on how much she smokes
7 Don't know

419. How do you think that breastfeeding affects a woman's chances of becoming pregnant?
(READ CHOICES)

1 Increases her chance of pregnancy
2 Decreases her chance of pregnancy
3 Does not affect her chance of pregnancy
7 Don't know

V. WOMEN'S HEALTH

500. In what month and year did you first have sexual intercourse, if ever?

Month: ___ ___ Year 19 ___ ___
 22=Never had sexual intercourse--->**GO TO Q517**
 33=No response--->**GO TO Q517**
 44=Don't remember

501. How old were you at that time?

___ ___ years
 77=Don't remember

IF RESPONDENT IS NOW OLDER THAN 24 YEARS, GO TO Q508

502. At that time what was your relationship to your first sexual partner? (SHOW CARD)

1 Husband--->**GO TO Q508**
 2 Fiance
 3 Boyfriend
 4 Friend/ Acquaintance
 5 Just met
 6 Forced intercourse/Rape----->**GO TO Q508**
 7 Family member/Incest---->**GO TO Q508**
 8 Other_____

9 No response--->**GO TO Q508**

503. Did you or your partner use a contraceptive method or do anything to prevent pregnancy at that time?

1 Yes
 2 No---->**GO TO Q506**
 8 Don't remember/Don't know---->**GO TO Q507**
 9 No response---->**GO TO Q507**

504. What method? (SHOW CARD)

1 Pills
 2 IUD
 3 Condoms
 4 Spermicide/Jelly/Cream
 5 Diaphragm
 6 Safe period method
 7 Withdrawal
 8 Douching
 77 Other_____

505. Who took the initiative to use this method?

1 Respondent
 2 Partner
 3 Both
 7 Don't remember

IF USED CONDOMS (CHECK Q504) GO TO Q508, OTHERWISE GO TO Q507

506. Why didn't you or your partner use a contraceptive method?

- 1 Did not expect to have sex
- 2 Did not know any methods
- 3 Hard for young people to get contraception
- 4 Did not know how/where to get contraception
- 5 Wanted to get pregnant
- 6 Health concerns about contraception
- 7 Wanted to use, but didn't have any
- 8 Did not think she could get pregnant
- 9 Partner refused to use contraception
- 10 Other _____
- 77 Don't know/Don't remember

507. Did your partner use a condom to prevent a sexually transmitted disease (STD) at that time?

- 1 Yes
- 2 No
- 7 Don't remember

508. Have you had sexual intercourse in the last 30 days?

- 1 Yes
- 2 No---->GO TO Q510
- 9 No response---->GO TO Q517

509. How many times?

- ___ ___ times
- 77 Don't remember
 - 99 No response

510. About how long has it been since you last had sexual intercourse? (GET ANSWER IN DAYS, WEEKS, MONTHS, OR YEARS)

- 1 ___ ___ days
- 2 ___ ___ weeks
- 3 ___ ___ months
- 4 ___ ___ years-->**GO TO Q517**
- 777 Don't remember
- 999 No response--->**GO TO Q517**

511. During the past 12 months, with how many men have you had sexual intercourse?

- ___ ___ men
- 77 Don't remember
 - 99 No response---->**GO TO Q517**

IF RESPONDENT HAS NOT HAD SEXUAL INTERCOURSE IN PAST 3 MONTHS? (SEE Q510), GO TO Q517

512. Did any partner ever use a condom with you in the past three months?

- 1 Yes
- 2 No---->**GO TO Q517**

QUESTIONS 513-516 REFER TO MEN WITH WHOM THE RESPONDENT HAS HAD SEXUAL INTERCOURSE IN THE PAST THREE MONTHS. IF MORE THAN THREE MEN, LIST ONLY THE THREE MOST RECENT.

	513	514	515	516
	What was your relationship with him?	Has he used a condom with you in the past three months?	Did he use a condom the last time you had sex with him?	Would you say that he uses a condom every time you have sex with him?
1 Most recent partner	1 Husband/Live-in partner 2 Regular boyfriend/Finace 3 Occasional boyfriend 4 Acquaintance 5 Other_____	1 Yes 2 No> NEXT LINE	1 Yes 2 No->NEXT LINE	1 Yes 2 No
2 2nd most recent partner	1 Husband/Live-in partner 2 Regular boyfriend/Finace 3 Occasional boyfriend 4 Acquaintance 5 Other_____	1 Yes 2 No> NEXT LINE	1 Yes 2 No->NEXT LINE	1 Yes 2 No
3 3rd most recent partner	1 Husband/Live-in partner 2 Regular boyfriend/Finace 3 Occasional boyfriend 4 Acquaintance 5 Other_____	1 Yes 2 No>GO TO Q517	1 Yes 2 No->GO TO Q517	1 Yes 2 No

517. Have you ever had a regular (not pregnancy related) gynecologic exam?

- 1 Yes
- 2 No---->GO TO Q519
- 7 Don't know---->GO TO Q519

518. When was the last time you had a gynecologic exam?

- ___ __ years ago
- 66 Less than 1 year ago
- 77 Don't remember
- 88 Don't remember, but more than one year ago

519. Has a health care provider ever discussed with you how to prevent getting sexually transmitted diseases?

- 1 Yes
- 2 No
- 7 Don't remember

520. What are all the signs or symptoms can you think of that a woman might have a sexually transmitted disease (STD)?
(DO NOT READ LIST)

- 1 Vaginal discharge
- 2 Genital itching
- 3 Painful urination
- 4 Lower abdominal pain
- 5 Sore or wart in the genital area
- 6 Other_____
- 7 Don't know any
- 9 Refusal to answer

IF RESPONDENT HAS NOT HAD SEX IN THE PAST 12 MONTHS (SEE Q510) OR REFUSED TO ANSWER Q511, GO TO Q527

521. In the past 12 months have you had any vaginal discharge that was not menstruation?

- 1 Yes
- 2 No----->GO TO Q523
- 7 Don't remember/Don't know----->GO TO Q523

522. Along with the discharge, did you have any:

	<u>Yes</u>	<u>No</u>	<u>Don't remember</u>
1. Itching?	1	2	7
2. Painful urination?	1	2	7
3. Lower abdominal pain?	1	2	7

523. In the past 12 months have you had any sores or warts in the genital area?

- 1 Yes
- 2 No
- 7 Don't remember/Don't know

IF Q521 AND Q523 ARE NOT ANSWERED "YES", GO TO Q527

524. The last time you had any of these symptoms did you see or consult anyone for advice or treatment?

- 1 Yes
- 2 No----->GO TO Q526
- 7 Don't remember----->GO TO Q527

525. Where did you go or whom did you see? (SHOW CARD)

- 1 Women's consultation----->GO TO Q527
- 2 Dermato/Venereal Clinic----->GO TO Q527
- 3 Gynecology office at a polyclinic----->GO TO Q527
- 4 Dermato/Venereal Office at a polyclinic----->GO TO Q527
- 5 Private office or clinic----->GO TO Q527
- 6 Pharmacy
- 7 Friend, relative, etc.
- 8 Local healer
- 9 Other (specify)_____

526 Why didn't you visit a health professional? (CODE THE MOST IMPORTANT REASON ONLY.)

- 1 Did not think it was serious enough/Was not necessary/Cured without a visit
- 2 Feared poor treatment by health staff
- 3 Services too far away
- 4 Services too expensive
- 5 Confidentiality/Didn't want people to know about sexual activity
- 6 Embarrassment
- 7 Did not know where to go
- 8 Other (specify) _____
- 77 Don't know/Don't remember

527. Do you think a person can be infected with the AIDS virus and not have any symptoms or signs of the disease?

- 1 Yes
- 2 No
- 3 Has not heard of AIDS
- 7 Don't know

528. Do you think condoms provide excellent, good, fair, or poor protection against most sexually transmitted diseases and infection with the AIDS virus? (**READ CHOICES**)

- 1 Excellent
- 2 Good
- 3 Fair
- 4 Poor
- 7 Don't know/Not sure

529. Do you think a person can be infected with an STD and not have any symptoms or signs of the disease?

- 1 Yes
- 2 No
- 7 Don't know

530. What do you think is your risk of getting an STD? (SHOW CARD)

- 1 High risk
- 2 Medium risk
- 3 Low risk
- 4 No risk
- 7 Don't know

For each of the following conditions/diseases, please tell me:

CONDITION	531. Have you ever heard of it?	532. Have you ever had it or been diagnosed with it?
A. Genital ulcer	1 Yes-->Q532 2 No--->B	1 Yes 2 No
B. Syphilis	1 Yes-->Q532 2 No--->C	1 Yes 2 No
C. Gonorrhea	1 Yes-->Q532 2 No--->D	1 Yes 2 No
D. Chlamydia	1 Yes-->Q532 2 No--->E	1 Yes 2 No
E. Pelvic inflammatory Disease	1 Yes-->Q532 2 No--->F	1 Yes 2 No
F. Genital Herpes	1 Yes-->Q532 2 No--->G	1 Yes 2 No
G. Human papilloma virus	1 Yes-->Q532 2 No--->H	1 Yes 2 No
H. Trichomoniasis	1 Yes-->Q532 2 No--->Q533	1 Yes 2 No

533 Do you currently smoke cigarettes?

- 1 Yes
2 No---->**GO TO Q535**

534. How many cigarettes do you smoke per day, on average?

- ___ cigarettes
66 = Rarely smoke/Less than 1 per day

535. What types of alcoholic drinks do you drink most often?

- 1 Vodka, cognac, other strong liquor (GE 30%)
2 Wine, champagne, liqueurs (9-29%)
3 Beer, other low alcohol drinks (LE 8%)
4 Rarely drink

536. In a typical week how much grams of _____(the strongest drink mentioned in Q535) do you drink?

- ___ ___ grams
6 6 = Usually none
7 7 = Don't know

9 9 = No response

VI. SOCIOECONOMIC CHARACTERISTICS

600. What is your nationality?
- 1 Ukrainian
 - 2 Russian
 - 3 Jewish
 - 4 Other (specify) _____
 - 9 No response
601. What language do you most often speak at home?
- 1 Ukrainian
 - 2 Russian
 - 3 Mixed Ukrainian-Russian
 - 4 Sometimes Russian, sometimes Ukrainian
 - 5 Other (specify) _____
602. What is your religion?
- 0 No religion---->**GO TO Q604**
 - 1 Orthodox
 - 2 Catholic
 - 3 Muslim
 - 4 Jewish
 - 5 Other (specify) _____
 - 9 Refused/Not stated--->**GO TO Q604**
603. About how often do you usually attend religious services? (SHOW CARD)
- 1 At least once a week
 - 2 At least once a month, but less than once a week
 - 3 Less than once a month
 - 4 Only on holidays
 - 5 Never/Almost never
 - 6 Other (specify) _____
604. Are you currently employed?
- 1 Yes--->**GO TO Q606**
 - 2 Yes, but on maternity/pregnancy leave--->**GO TO Q607**
 - 3 Self-Employed--->**GO TO Q606**
 - 4 No
605. Which of the following best describes your situation?
- 1 Can't find a job/Unemployed
 - 2 Factory/Former place of employment closed
 - 3 Don't want to work/No need to work
 - 4 Unable to work
 - 5 Student
 - 6 Other (specify) _____

GO TO Q607

606. Do you currently work one job or more than one job?

- 1 One job
- 2 More than one job

607. Please tell me whether this household or any member of it has the following items:

	<u>Yes</u>	<u>No</u> _____
A. Bathroom/Shower	1	2
B. Color television	1	2
C. VCR	1	2
D. Automobile	1	2
E. Auto Washing machine	1	2
F. Telephone	1	2
G. Personal computer	1	2

VII. VIOLENCE

700. Thinking back to your childhood and adolescence, did you ever see or hear your parents or step-parents physically abuse each other?

- 1 Yes
- 2 No
- 3 Did not live with 2 parents
- 7 Don't remember
- 9 Refuse to answer

701 Do you recall ever being physically abused as a child by anyone in your household or family?

- 1 Yes
- 2 No
- 7 Don't remember
- 9 Refuse to answer

702. This next set of questions is about violence and physical abuse that may have happened between you and a partner or ex-partner. When we say a partner, we mean a husband, ex-husband, as well as any other person you have been living with, without being married. Did a partner or ex-partner ever threaten to hit you, shove, or slap you, threaten you with a knife or other weapon, or actually hit you?

- 1. YES
- 2. NO----->END OF INTERVIEW
- 3. NEVER HAD A PARTNER ----->END OF INTERVIEW
- 7. DO NOT REMEMBER----->END OF INTERVIEW

703. Please tell me if you have ever been physically abused, by a partner or ex-partner in the following ways (READ A-D):		704. When was the last time this (A-D) happened to you?
A. Threaten to hit you or throw something at you?	1 YES---> Q704 2 NO----> Q703B 7 DON'T REMEMBER----> Q703B 9 REFUSE---> Q703B	1. WITHIN THE LAST YEAR 2. 1-3 YEARS AGO 3. 4-5 YEARS AGO 4. AT LEAST 5 YEARS AGO 7. DON'T REMEMBER GO TO 703B
B. Push or shove you?	1 YES---> Q704 2 NO----> Q703C 7 DON'T REMEMBER----> Q703C 9 REFUSE---> Q703C	1. WITHIN THE LAST YEAR 2. 1-3 YEARS AGO 3. 4-5 YEARS AGO 4. AT LEAST 5 YEARS AGO 7. DON'T REMEMBER GO TO 703C
C. Kick or hit you with a fist or with something?	1 YES---> Q704 2 NO----> Q703D 7 DON'T REMEMBER----> Q703D 9 REFUSE---> Q703D	1. WITHIN THE LAST YEAR 2. 1-3 YEARS AGO 3. 4-5 YEARS AGO 4. AT LEAST 5 YEARS AGO 7. DON'T REMEMBER GO TO 703D
D. Threaten you with a knife or other weapon?	1 YES---> Q704 2 NO----> Q705 7 DON'T REMEMBER----> INST. BEFORE Q705 9 REFUSE---> INST BEF Q705	1. WITHIN THE LAST YEAR 2. 1-3 YEARS AGO 3. 4-5 YEARS AGO 4. AT LEAST 5 YEARS AGO 7. DON'T REMEMBER

IF NONE OF THE INCIDENTS IN A-D HAPPENED IN THE LAST YEAR (Q704A, B, C, OR D=1), END INTERVIEW.

705. In the past 12 months, did you have any swelling, bruises, cuts, or other physical injuries as a result of your partner's actions?

- 1. Yes
- 2. No----->GO TO Q708
- 7. Don't remember----->GO TO Q708

706. Did you see a doctor, or other medical care provider for medical treatment of these injuries?

- 1. Yes
- 2. No----->GO TO Q708
- 7. Don't remember ----->GO TO Q708

707. Were you hospitalized as a result of these injuries?

- 1. Yes
- 2. No
- 7. Don't remember

708. Did you talk about this(these) incident(s) with (READ 1-5)?

	<u>YES</u>	<u>NO</u>
1. Police	1	2
2. Family member	1	2
3. Friend	1	2
4. Health provider/Social Worker	1	2
5. Psychologist	1	2
6. Other (Specify)_____	1	2

709. The last time your husband hit or threatened you had he been drinking alcohol?

- 1 Yes---->END OF INTERVIEW
- 2 No
- 7 Don't know/Don't remember

710. At any time in the past year when he hit or threatened you had he been drinking alcohol?

- 1 Yes
- 2 No
- 7 Don't know/Don't remember

END OF INTERVIEW

TIME INTERVIEW ENDED ___ : ___

INTERVIEWED BY: _____

QUESTIONNAIRE REVIEWED BY: _____

DATA ENTRY OPERATOR # ____