From Cough to Cure: A Path of Ideal Behaviors in Tuberculosis Control

LEVELS * Poor knowledge Misperception of Poor knowledge Poor knowledge of Poor knowledge of Poor knowledge of of TB symptoms costs of diagnostic steps length of treatment length of treatment length of treatment Poor knowledge Preference for non- Expectations of TB care and DOTS health services about medical * Stigma * Stigma Stigma cure services (get meds * Attitude about not tests) ndividual Stigma related to health services TB diagnosis Stigma Low risk preception * Social Norm Continue Seek Go to Complete Begin Complete IDEAL BEHAVIOR DOTS Diagnosis Treatment Treatment Care Treatment & follow-up smears & final smear * Time, cost, distance Providers' poor * Time, cost, distance * Time, cost, distance * Time, cost, distance to DOTS facility knowledge of to DOTS facility to DOTS facility to DOTS facility correct procedures Lack of linkages * Poor quality of * Poor quality of * Poor quality of Providers' poor between DOTS and services services services inter-personal other providers (noncommunication DOTs & HIV care) * Health providers fall Health providers fail Health providers fail to give adequate to give adequate to give adequate Lack of resources, Missed diagnosis information information information including human and / or lack of resources * Lack of Lack of Lack of referral by non-DOIS Poor quality of providers medications medications medications services (hours, waittime) Time, cost and distance

