



Mapping hotspots of climate change and food insecurity in the global tropics

A report by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS)

Project team: Polly Ericksen, Philip Thornton, An Notenbaert, Laura Cramer, Peter Jones, Mario Herrero.

Advance Copy.
EMBARGOED FOR RELEASE UNTIL 00:01 GMT ON FRIDAY, 3 JUNE 2011

This research was coordinated by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), in collaboration with the International Livestock Research Institute (ILRI).

CCAFS Coordinating Unit - Department of Agriculture and Ecology, Faculty of Life Sciences, University of Copenhagen, Rolighedsvej 21, DK-1958 Frederiksberg C, Denmark.
Tel: +45 35331046; Email: ccaafs@life.ku.dk Website: ccaafs.cgiar.org

CCAFS is a strategic partnership of the Consortium of International Agricultural Research Centers (CGIAR) and the Earth System Science Partnership (ESSP). The program is supported by the European Union (EU), the United States Agency for International Development (USAID), the Canadian International Development Agency (CIDA), New Zealand's Ministry of Foreign Affairs and Trade, the Danish International Development Agency (Danida) and the UK Department for International Development (DFID), with technical support from the International Fund for Agricultural Development (IFAD). The views expressed in this document cannot be taken to reflect the official opinions of these agencies, nor the official position of the CGIAR or ESSP.

©2011. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS).

Correct citation

Ericksen P, Thornton P, Notenbaert A, Cramer L, Jones P, Herrero M. 2011. Mapping hotspots of climate change and food insecurity in the global tropics. CCAFS Report no. 5 (Advance Copy). CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). Copenhagen, Denmark. Available online at: www.ccaafs.cgiar.org.

Executive summary

This study was coordinated by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) to identify areas that are food insecure and vulnerable to the impacts of future climate change, across the priority regions for the CGIAR centres. The research was undertaken by a team of scientists from the International Livestock Research Institute (ILRI). The study relied on maps: first, of variables that indicate the different aspects of food security (availability, access and utilization), and second, of thresholds of climate change exposure important for agricultural systems. Vulnerability was assessed using a domain approach based upon the Intergovernmental Panel on Climate Change (IPCC) framework of vulnerability as a function of exposure, sensitivity and coping capacity. Nine domains were identified; for each domain areas of the tropics were classified by high or low exposure, high or low sensitivity, and high or low coping capacity.

Length of growing period declines by 5% or more across a broad area of the global tropics, including heavily cropped areas of Mexico, Brazil, Southern and West Africa, the Indo-Gangetic Plains, and Southeast Asia. This suggests that at a minimum, most of the tropics will experience a change in growing conditions that will require adaptation to current agricultural systems. High temperature stress (above 30°C) will be widespread in East and Southern Africa, north and south India, Southeast Asia, northern Latin America and Central America. Length of growing period flips to less than 120 days in a number of locations across the tropics, notably in Mexico, northeast Brazil, Southern and West Africa and India. This is a critical threshold for certain crops and rangeland vegetation; hence these are important target areas for high exposure to climate change. Reliable crop growing days decrease to critical levels, below which cropping might become too risky to pursue as a major livelihood strategy in a larger number of places across the global tropics, including West Africa, East Africa, and the Indo-Gangetic Plains. Much of the tropics already experiences highly variable rainfall, above the median of 21% for cropped areas. Thus any increases in this variability will make agriculture more precarious.

In terms of food security, the net food production index is stagnant in all areas of interest, with differences between countries rather than regions. GDP per capita is low in many countries in Africa, as well as in Afghanistan, Nepal, Bangladesh, Laos and Cambodia. Poverty hotspots are West, Central and East Africa, India and Bangladesh and Southeast Asia. Africa and south Asia are clearly much more chronically food insecure regions than Latin America or China.

The most vulnerable domain for most exposures is high exposure, high sensitivity and low coping capacity (HHL). Such areas are highly vulnerable to climate change and have significant agriculture and high levels of food insecurity. Under exposure 1 (LGP decreases more than 5%), HHL is the category with the most people, followed by HHH. For exposure 2 (LGP flips) HHL is very small in terms of people; most people are in the categories LHL or LHH. Exposure 3 (reliable crop growing days - RGCP flips) has about 10 million more people in the HHL category, but again most people are in LHL or LHH. Under exposure 4 (maximum temperature -Tmax flips), the vulnerable population more than triples relative to exposure 3. Under exposure 5 (temperature flips) again more people are in the vulnerable categories. Under exposure 6 (rain per rainy day decrease), the most vulnerable population drops to 27.5 million, while under exposure 7 (rain per rainy day increase) 45.7 million are in the HHL category. This suggests that the choice of domain variables makes a big difference in terms of areas included.

Contents

Section 1: Introduction	9
Food security concepts and indicators.....	9
How might climate change increase risk of food insecurity?	14
Section 2: Climate change hotspot indicators across the global tropics.....	17
Threshold maps.....	18
Section 3: Food security maps across the global tropics.....	32
Availability indicators.....	32
Access indicators.....	39
Utilization indicators.....	42
Resource pressure	46
Section 4: Vulnerability domains	48
The criteria and their thresholds.....	48
The vulnerability domains, with different exposure variables.....	52
Section 5: Regional maps	62
East Africa	62
West Africa.....	70
South Asia	77
Conclusion: Analysis of hotspots and implications for CCAFS research.....	84
References	87

List of Maps

Map 2.1 The agricultural land area for regions of interest to CCAFS. Pa = pasture, Cr = irrigated cropping, Lg = length of growing period ≥ 60 days.....	18
Map 2.2.Areas that will experience more than a 5% reduction in LGP.....	19
Map 2.3.Areas that will flip from LGP >120 days in the 2000s to LGP < 120 days by 2050	20
Map 2.4. Areas that flip from > 90 reliable crop growing days (RCGD) per year in the 2000s to <90 RCGD by 2050.....	21
Map 2.5. Areas where the average annual temperature flips from $<8^{\circ}\text{C}$ in the 2000s to $> 8^{\circ}\text{C}$ by 2050.	22
Map 2.6. Areas where average annual maximum temperature will flip from $< 30^{\circ}\text{C}$ to $> 30^{\circ}\text{C}$	23
Map 2.7. Areas where maximum temperature during the primary growing season is currently $< 30^{\circ}\text{C}$ but will flip to $> 30^{\circ}\text{C}$ by 2050 (during the primary growing season).	24
Map 2.8. Areas where CV of rainfall is currently high.....	25
Map 2.9. Areas where CV of rainfall is more than 21%.	26
Map 2.10. Areas where rainfall per day decreases by 10% or more between 2000 and 2050.	27
Map 2.11. Areas where rainfall per rain day increases by 10% between 2000 and 2050.	28
Map 2.12. Number of identified climate change thresholds	29
Map 2.13. The frequency SPI defined as the average number of drought events per year per pixel (for the period 1974-2004).....	30
Map 2.14. Average flood frequency based upon data from the Dartmouth Flood Observatory.....	31
Map 3.1. Maize yields mapped by pixel across the global tropics.	32
Map 3.2. Rice yields mapped by pixel across the global tropics.....	33
Map 3.3. Millet yields mapped by pixel across the global tropics.	33
Map 3.4 Bean yields mapped by pixel across the global tropics.....	34
Map 3.5. Wheat yields mapped by pixel across the global tropics.	35
Map 3.6. Sorghum yields are mapped by pixel across the global tropics.	36
Map 3.7. Cassava yields mapped by pixel across the global tropics.	37
Map 3.8. Average food production index 2003-2007.	38
Map 3.9. GDP per capita 2005.	39
Map 3.10. Population living on less than USD 2 per day.	40
Map 3.11. Transport time to markets.	41
Map 3.12. Stunting prevalence.....	42

Map 3.13. Wasting prevalence.....	43
Map 3.14. Malnourished children per sq km.....	44
Map 3.15. Population using unimproved water source.....	45
Map 3.16. Annual population growth.....	46
Map 3.17. Arable land per capita.....	47
Map 4.1. Reliable crop growing days flip to less than 90 days.....	49
Map 4.2. Areas with greater than 16% cropping (assumed sensitive to change in climate).	50
Map 4.3. Areas with high and low coping capacity for a change in climate.	51
Map 4.4. Exposure 1: Areas where there is greater than 5% change in LGP.	52
Map 4.5. Exposure 2: LGP flips from more than 120 days to less than 120 days.....	54
Map 4.6. Exposure 3: RCGDs flip from more than 90 days to less than 90 days (threshold 2).	55
Map 4.7. Exposure 4: Maximum daily temperature flips from <30 deg C to >30 deg C (threshold 4).....	56
Map 4.8. Exposure 5: Maximum daily temperature during the growing season flips from <30 deg C to > 30 deg C (threshold 5).	57
Map.4.9. Exposure 6: Rain per rainy day decreases by more than 10% (threshold 6).....	58
Map 4.10. Exposure 7: Rain per rainy day increases by more than 10% (threshold 7)	59
Map 4.11. Exposure 8: CV rainfall currently greater than 21%	60
Map 4.12. Exposure 9: Mean annual temperature flips from less than 8 deg C to more than 8 deg C.	61
Map 5.1. Threshold 1 (Length of growing period flips from more than 120 days to less than 120 days);.....	62
Map 5.2. Threshold 2 (Number of reliable crop growing days flips from more than 90 to less than 90).	62
Map 5.3. Threshold 4 (maximum temperature flips to more than 30 deg C).	63
Map 5.4.Threshold 5 (maximum temperature in the growing season flips to more than 30 deg C).	63
Map 5.5. Threshold 6 (Average rainfall per rainy day increases by more than 10%).....	64
Map 5.6. Threshold 7 (Average rainfall per rainy day decreases by more than 10%).	64
Map 5.7. Coefficient of rainfall variability (mode is 21% for global tropics).....	65
Map 5.8. Percent area cropped	65
Map 5.9. Economic access: % population on less than USD 2 a day	66
Map 5.10. Physical access: travel time to nearest town >250K	66
Map 5.11. Kilocalories available per person per day	69

Map 5.12. Stunting prevalence (% under five).....	69
Map 5.13. LGP flip.....	70
Map 5.14. RCGD flip.....	70
Map 5.15. Tmax flip.	71
Map 5.16. Tmax growing season flip.	71
Map 5.17. Rainfall per rainy day increase.....	72
Map 5.18. Rainfall per rainy day decrease.....	72
Ma 5.19. Percent area cropped	73
Map 5.20. Coefficient of rainfall variability (mode is 21% for global tropics)	73
Map 5.21. Economic access: % population on less than USD 2 per day	74
Map 5.22. Physical access: travel time to nearest town >250K	74
Map 5.23. Kilocalories available per person per day	76
Map 5.24. Stunting prevalence (% under five).....	76
Map 5.25. LGP flip.....	77
Map 5.26. RCGD flip.....	77
Map 5.27. Tmax flip.	78
Map 5.28. Tmax growing season flip.	78
Map 5.29. Rainfall per rainy day increase.....	79
Map 5.30. Rainfall per rainy day decrease.....	79
Map 5.31. Coefficient of rainfall variability.....	80
Map 5.32. Percent area cropped.	80
Map 5.33. Economic access: % population on less than USD 2 per day.	81
Map 5.34. Physical access: travel time to nearest town >250K.	81
Map 5.35. Kilocalories available per person per day.	82
Map 5.36. Stunting prevalence (% under five).....	82

List of Figures

Figure 1. The components of food security.	10
Figure 2. Conceptual diagram of food system vulnerability.	14
Figure 3. Price volatility Nairobi.	67
Figure 4. Price volatility Kampala.	67
Figure 5. Price volatility Addis Ababa.	68
Figure 6. Price volatility Accra.	75
Figure 7. Price volatility Bamako.	75
Figure 8. Price volatility Niamey.	75
Figure 9. Price volatility Bangladesh.	83
Figure 10. Price volatility Delhi.	83

List of Tables

Table 1. Vulnerability of determinants of food security to water availability.	15
Table 2. Vulnerability domains based on exposure, sensitivity and coping capacity.	48
Table 3. Area and population included in the vulnerability domain exposure 1.	53
Table 4. Area and population included in the vulnerability domain exposure 2.	54
Table 5. Area and population included in the vulnerability domain exposure 3.	55
Table 6. Area and population included in the vulnerability domain exposure 4.	56
Table 7. Area and population included in the vulnerability domain exposure 5.	57
Table 8. Area and population included in the vulnerability domain exposure 6.	58
Table 9. Area and population included in the vulnerability domain exposure 7.	59
Table 10. Area and population included in the vulnerability domain exposure 8.	60
Table 11. Area and population included in the vulnerability domain exposure 9.	61

Section 1: Introduction

The CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) commissioned a small team of ILRI/ CCAFS Theme 4 staff to conduct a rapid assessment across the global tropics of the vulnerability of food security to climate change. The goal was to identify 'hotspot' locations where climate change impacts are projected to become increasingly severe by 2050 and food insecurity is currently a concern, using a range of indicators. The project is intended to help CCAFS by giving input to the selection of new target regions (i.e. multi-country) ex ante, and also as an ex post check of the three target regions chosen at the start of the project: East Africa, West Africa and the Indo-Gangetic Plains (IGP). In addition, the project is the start of a process that will link to regional scenarios and regional and local quantification work. Using maps as aids in the visualization of the possible impacts of climate change across the tropics and within regions will be important. This project also demonstrates the multiple indicators of food security that can be mapped and will interact with climate change. Finally, the project contributes to CCAFS work by including methods for mapping both food insecurity and the impacts of climate change on agriculture and food security, and giving guidance on interpreting results, particularly overlap, or lack of it, between the two categories of hotspots.

The work unfolded in several steps. First, a data scoping study of food security indicators was undertaken to identify both indicators of the components of food security and indicators that could be mapped from local to global level. This study also reviewed the recent debates about measuring food insecurity. Second, a workshop was held to determine which food security indicators could best be mapped. The workshop participants¹ also discussed which climate change impact indicators are available and can be mapped, and how these are relevant to food insecurity. A framework for assessing the vulnerability of food security to climate change was discussed and future work to develop this for CCAFS identified. Then both the food (in)security and the climate change impact indicators were mapped and hotspots identified. Finally we mapped nine vulnerability domains, combining exposure to climate change impacts, sensitivity to this change and coping capacity for adverse impacts on food security. All results are mapped for the entire region of interest.

Food security concepts and indicators

Definition

Food security is defined as a situation that exists when all people, at all times, have physical and economic access to sufficient, safe and nutritious food to meet their dietary needs and food preferences for an active and healthy life (FAO in Stamoulis and Zezza 2003). In order for a population to be characterized as food secure, they have to have enough nutritious, yet affordable food, and be sufficiently protected against future disruptions to the access of adequate food. For analytical purposes, the complex definition of food security can be broken up into three components: availability, access and utilization. A fourth dimension, stability, refers to the requirement that food secure people have access to appropriate food at all times. Ericksen (2008a) defined access, availability and utilization as shown below. Note that there is overlap among these dimensions; for example both availability and access discuss the need for equitable distribution of food and both access and utilization reflect the requirement for food secure people to have food that meets social and cultural preferences.

Food **availability** refers to the amount, type and quality of food a unit (such as community, household, individual) has at its disposal to consume. It can be further broken down into production, distribution and exchange (see Figure 1).

¹ Alison Misselhorn, Keith Weibe, Tanya Beaudreau, Erin Lentz, Jeronim Capaldo, Jean Balie, and Mark Smulders.

Access to food refers to the ability of a unit to obtain access to the type, quality, and quantity of food it requires. The subcomponents of access are affordability, allocation and preference.

Food **utilization** refers to the individual or household capacity to consume and benefit from food. The three subcomponents are nutritional value, food safety and social value.




Figure 1. The components of food security.

Note that there is a hierarchical relationship among these three components: “food availability is necessary but not sufficient for access, and access is necessary but not sufficient for utilization” (Webb et al 2006). Thus there may be plenty of food in the markets, but if it is too expensive for people to buy then they are food insecure. Or the food that is available and affordable may be of inferior quality and hence people may be micronutrient deficient. Diseases such as HIV can also interfere with people’s ability to utilize foods properly while a hazard or shock could adversely affect any one of these aspects (make food more expensive, cause shortfalls in production, or affect food safety); hence stability over time is also included. Food security status is also linked to a number of socio-economic characteristics, including wealth, age and status within a household. In addition, the gender dimension of food security has been explored in terms of the differences between food security outcomes for men and women in the same household and with regard to the key role that women play in attending to the food security of their children (FAO 2011).

An important distinction exists between chronic and transitory food insecurity, both in terms of indicators and underlying causes. Transitory food insecurity implies that there has been a temporary shock to food security but that the situation will return to normal after a period of time. Chronic food insecurity

means people cannot meet their basic requirements for a significant period of time with a more long term outcome. The drivers of transitory and chronic food insecurity are often different: transitory results from variability in production, food prices, or incomes, while chronic is the result of systemic or structural failure such as poverty or political marginalization. The two do interact, as chronic food insecurity results from one or a series of transitory shocks causing very vulnerable households to lose the ability to cope with any future shocks. Chronic food insecurity, such as high malnutrition rates or low household incomes, can make populations much more vulnerable to severe transitory shocks like a price increase (Misselhorn et al 2010). This distinction is relevant for CCAFS because climate change will have longer term impacts on production trends but in the short term it will increase variability and extreme events. Food insecurity is always due to multiple stressors, so the impact of climatic shocks has to be in the context of this multiple exposure.

Measuring Food (In)Security

The continued refinement of the definition of food security over the past three decades in response to improved empirical evidence has brought a greater understanding of both the components and the underlying causes of food insecurity. Despite this, the measurement and evaluation of food insecurity remains difficult. For much of the 1970s there was a bias equating food security with availability; this began to change after Sen's 1981 book articulating the idea that famines are not necessarily caused by a lack of food, but a lack of access to it (Webb et al 2006). Given the acknowledged multiple dimensions of food security, one indicator does not give the entire picture. Because it is a complex concept, involving the interaction of many different factors, there is no single measurement for food security (Riely et al 1999). Multiple indicators, most of which are proxy indicators, are used to determine food security status, and these vary depending on the global, national, district, household, or individual level.

Webb et al (2006) outline three recent advances in understanding and measuring food security. The first is a shift from measuring availability and utilization to measuring "inadequate access." Researchers and policy makers are increasingly acknowledging that purchasing power is the key to access. This shift happened after research showed a weak relationship between food availability and nutritional status, at national, household and individual levels. While the shift to measuring access is a crucial development to better understanding food insecurity, there are no exact indicators of access failure. Widely accepted proxies of both food supply failure and impaired utilization are available, but the relatively new attempt to measure inadequate access is not as yet developed. Households have many ways to mitigate or cope with negative shock and the search for measures of access failure has recently focused on measuring the potential use of these coping behaviours as indicators (Coates et al 2006).

The second shift in food security measurement is an expanding effort to find fundamental measures instead of relying on proxy measures. Much of the food security measurement that takes place in developing countries is 'derived measurement', which relies on proxy measures such as food consumption, income, or assets - thought to be indicators of food security. This is problematic because:

- There is no empirical link between the measures and actual food insecurity.
- The strength of correlation may differ between contexts and causes and consequences may also differ.
- The actual causes and consequences of food insecurity may be overlooked when using derived measures.

However proxy measures are often all there are to use, given the absence of fundamental measures (Webb et al 2006).

The third shift identified by Webb et al (2006) is a shift from focusing on objective to subjective measures: moving away from absolute measures, such as poverty lines and expenditure on goods and services, to experiential and perception measures that can be analyzed using econometric methods. For example, Coates et al (2006) analyse 22 separate scales and ethnographies exploring food insecurity and find four universal domains: uncertainty or worry, insufficient quantity, inadequate quality and social unacceptability. The first and last of these are subjective assessments but are fundamental to the experience of food insecurity.

In addition to being complex, measuring food security is time consuming and it is expensive to gather data at individual and household levels. Availability estimates can be generated at relatively low cost, especially at national and global scales, but they hide the difficult aspect of understanding access and utilization patterns at household and individual levels; for example differences between males and females, young and old. "The global figures mask considerable heterogeneity among and within regions. ... Food security measures based on household and individual data routinely generate higher estimates of food insecurity than those derived from more aggregate data" (Barrett 2010, p.826). To fully understand food security, we must measure more than just nutritional status and availability of supply; the element of vulnerability must be captured as well (Barrett 2010). This is a difficult concept to assess. Most measures of food insecurity (such as food intake, food production or income, and nutritional status) are static in nature and do not predict the possibility of having inadequate food in the future (Christiaensen and Boisvert 2000). It is not possible from such measures to assess if those who are currently food secure will become insecure in the future or vice versa. "Observational data necessarily report on the past... An ideal food security indicator would reflect the forward-looking time series of probabilities of satisfying the access criteria" (Barrett 2010 p.826).

The problem of defining thresholds complicates the conceptual complexity of food security and its difficulties of measurement. There is little agreement on which thresholds should be used to describe the difference between food security and food insecurity, with the exception of FAO's standard of undernourishment, and perhaps the WHO growth standards for children under five (WHO 2009). Two widely applied food security classification systems, the Famine Early Warning System Network (FEWSNET) and the Integrated Food Security Phase Classification System (IPC), differ in several key aspects of classification. FEWSNET classifies food security outcomes (such as food deficits and malnutrition), but not underlying conditions (for example, poor crop production, chronic poverty, or high food prices). In contrast, the IPC takes into account underlying conditions affecting food security, such as civil strife and hazards. The IPC also sets more quantitative thresholds than FEWSNET, including crude mortality rates, acute malnutrition, stunting, caloric intake, and water availability. Also, FEWSNET colours map areas with the food security level of the majority of the poorest wealth group in a given area, while the IPC colours a region based on the highest level of food insecurity found there. A further complication is the use of stunting (low height for age) for chronic food insecurity and the use of wasting (low weight for height) for transitory or acute food insecurity. Food security responses are often based on the indicators collected, so measuring either stunting or wasting may produce different response options, either in the form of direct interventions or policies. Also for consideration in the complexity of measuring and setting thresholds for food insecurity is the issue of lagging versus early indicators. As indicated by Barrett (2010), most measures of food security are reporting on the past: nutritional indicators, whether of stunting or wasting, reflect that food insecurity has already occurred. While monitoring food prices can provide an indicator of future food insecurity to some extent, researchers are still working to find predictive indicators that can be used to design preventative measures against food insecurity, rather than learning about food insecurity once it has occurred and only then intervening to mitigate it.

A final point is the issue of level of measurement. As discussed above, referring to Barrett (2010), global food security measures are broad and hide significant disparities between regions. National indicators can offer comparability between countries, but also obscure great differences at provincial or district levels

within a single country. Given the complex nature of food security, household level surveys and local studies are more suited to explain processes than global aggregate measures, but lack comparability and broad coverage. Global, national, and regional levels of production and availability figures can paint a broad view of the first pillar of food security while sub-national measures of livelihood and coping strategies offer a more detailed picture of access. Lastly, stability is not only difficult to define but also to measure, and even more difficult to predict.

To find suitable indicators to map, the project team compiled a large database of food security measures for each of the components, first at the global level, then for each of the three regions of interest for CCAFS, initially East Africa, West Africa and the Indo-Gangetic Plains, and for each country in the region. These indicators and descriptions are attached in Appendix 1. We identified indicators for access, availability and utilization, at each of the geographic levels of interest. We considered a number of indicators for stability and these are also shown, but as this requires predictive capacity, which at the moment this exercise does not include, we left this for future work. However we will include a temporal and dynamic dimension in the scenarios and modelling work.

How might climate change increase risk of food insecurity?

A unit of analysis or a system is vulnerable to an adverse shock or change if it will suffer harm from which it is difficult to recover. Vulnerability to climate change can be conceived as a function of exposure to a hazard (such as changes in temperature or precipitation from climate change), sensitivity to that hazard (for example, maize yields are highly sensitive to drought) and finally, adaptive capacity in the face of the hazard. Adaptive capacity reflects the ability of a system or community to manage the impacts of a shock. Figure 2 below, illustrates these concepts for food systems. If people have sufficient assets or strategies to manage a shock without suffering harm, then they will not be vulnerable (McCarthy et al 2001).




Figure 2. Conceptual diagram of food system vulnerability. GECAFS 2005.

Combining the hazard indicators with indicators of sensitivity and adaptive capacity helps to evaluate vulnerability, as exposure to changed climate patterns alone will not necessarily lead to increased vulnerability. As explained above, food insecurity is a function of multiple stressors, and climate change will add more stress, or potentially increase food insecurity if people, households or geographic areas are highly sensitive to the climate hazard and have insufficient coping capacity. Most of the measures of food (in)security collected are outcome indicators: e.g. the malnutrition rate in a community rather than the drivers or underlying processes that contribute to that malnutrition, such as chronic poverty, disease, or lack of access to diverse and nutritionally adequate diets. To have predictive capacity the exercise should model the drivers of food security, of which climate is only one, as they evolve over time to deliver food security outcomes. The table below is a simplified example of such an approach, based upon work by the Global Environmental Change and Food Systems (GECAFS) programme in the Indo-Gangetic Plains (Ericksen2008b).

Table 1. Vulnerability of determinants of food security to water availability

Key determinant	Determinant characteristics	Sensitivity to water availability	Adaptive capacity	Vulnerability to water availability
NUTRITIONAL VALUE Food diversity	A diet of rice and lentils supplemented by milk	Cows need four months of rain to produce milk	No functioning milk market to buy from when own production fails	High , due to no ability to purchase milk
Primary protein	Lentils are eaten every day	Lentils need two months of rain	Lentil market functions so can always buy them	Low , because can purchase lentils
AFFORDABILITY Household incomes	Agriculture is the main source of income	Agricultural earnings depend upon good yields and functioning markets	When crops fail, some work can be found in towns or further away	Moderate , due to social and economic constraints on migration

However, this is really hard to do across the global tropics at the necessary geographic level – that is, sub national – without household survey data to explain why people are food insecure; for example, how they meet their food basket requirements, and which entitlements fail as the result of a (climate) shock. Devereux (2007) uses data for Malawi to explore how droughts and floods affect each of four food security entitlements: production-based, labour-based, trade-based and transfer-based. With good household data for Malawi, Devereux can evaluate the impact of each type of entitlement failure. He illustrates that production-based entitlements are affected by harvest failure in the short term if households are highly dependent upon agriculture for food, and in the longer term through increased risk, which has been shown to dampen investment in increasing agricultural productivity. In the good year of 2000/2001, only one in four Malawian farmers was self sufficient in maize, and in the crisis year of 2001/2002 only 2.6% of surveyed households were self-sufficient, while 92% had run out of their own farmed maize after nine months. The impact of these transitory climate shocks on production-based entitlements is compounded by other processes such as decreasing land holdings and a decline in access to inputs. Labour-based entitlements are affected if off-farm employment

opportunities to earn cash to purchase food are influenced by a drought or flood. As so few Malawians are self-sufficient in maize, many turn to rural off-farm labour for both cash and food, but these opportunities are declining. In 2001/2002 the number of people seeking off-farm work during the crisis was much higher than the number of available jobs, so relative to the crisis of 1991/1992 labour-based entitlements were much harder to satisfy. Failures of trade-based entitlements occur when weather shock causes food prices to rise, while at the same time asset prices (such as for livestock) fall in market value. In Malawi, there are predictable increases in food prices every year during the hungry season, and in the 2001/2002 crisis retail prices of maize and cassava increased 300% in January 2002. As agricultural markets in Malawi often fail, these high food prices do not attract traders to supply more food. Finally, transfer-based entitlement can fail if informal social mechanisms are overwhelmed in a prolonged food crisis, as well as due to long-term trends. Several factors combined in rural Malawi, among them social change, the high prevalence of HIV/AIDS, and a low level of urbanization. These combined factors mean there is relatively less remittance income coming from urban areas in times of crisis.

In this project we mapped:


- current food insecurity outcome indicators;
- climate change hotspots in 2050;
- the overlap between these.

Section 2: Climate change hotspot indicators across the global tropics

In this project, we used modelled predictions of changes in temperature and precipitation up to 2050 to derive indicators that are relevant for food systems and food security. The data are available on www.futureclim.info and described in Jones et al (2009). For this report, we used the mean climatology of the four general climate models (GCMs) available from futureclim.info to generate daily weather data and define thresholds important for agriculture and food security. Of the 22 or so climate models used for the IPCC's Fourth Assessment Report (2007), output data are not always readily available for the core variables that are needed to drive many crop and pasture models: precipitation, maximum daily air temperature, and minimum daily air temperature. From the World Climate Research Program's (WCRP) Coupled Model Intercomparison Project phase 3 (CMIP3) multi-model dataset, we obtained data for three GCMs: the CNRM-CM3 model from France, CSIRO-Mk3.0 from Australia, and MIROC 3.2 (medium resolution) from Japan. We also obtained data for the ECHam5 model (from Germany) from the Climate and Environmental Data Retrieval and Archive (CERA) database at the German Climate Computing Centre (DKRZ). These and other climate models are extensively described in Randall et al. (2007).

The maps below are for the entire global tropics and illustrate differences among regions and within continents. In Section 3 we show maps by regions, to better illustrate variability within regions and countries.

The agricultural land area for regions of interest to CCAFS (between 35 °S and 45 °N, masking out Europe, the US, Argentina, Chile, Australia and New Zealand) are shown in Map 2.1. For our purposes, agricultural land area was defined as places in which the length of growing period (LGP) is greater than or equal to 60 days (i.e. agriculture is possible), plus areas identified as pasturelands and irrigated croplands from satellite imagery (see Ramankutty et al, 2008). The different categories, including overlap, are mapped below.




Map 2.1 The agricultural land area for regions of interest to CCAFS. Pa = pasture, Cr = irrigated cropping, Lg = length of growing period ≥ 60 days.

Threshold maps


We then defined nine types of climate change hotspots using thresholds for 2050. We used thresholds rather than continuous variables to define discrete areas. The climate change hotspot indicators across the global tropics described and mapped here were derived from the mean outputs of four climate models. There are many uncertainties associated with these indicators, not least the fact that different climate models give different results. These differences may be quite large, particularly for projected changes in rainfall patterns and amounts. The essential reason for this is that there are still many unknowns about the details of how climate may change in the future due to anthropogenic forcings. The climate models are still rather imperfect representations of reality, and as different teams of scientists build these models, these imperfect representations can differ substantially. In Appendix 2, we present probability maps of the eight thresholds derived from GCMs (thus coefficient of variability [CV] rainfall is not included).

1. Areas that will experience more than a 5% reduction in LGP (Map 2.2). LGP is defined by the average number of growing days per year, in which a growing day is one in which the average air temperature is greater than 6 °C and the ratio of actual to potential evapo-transpiration exceeds 0.35 (Jones and Thornton 2008). The growing season begins once five consecutive growing days have occurred and ends once 12 consecutive non-growing days occur.



Map 2.2. Areas that will experience more than a 5% reduction in LGP.

2. Areas that will flip from LGP greater than 120 days in the 2000s to LGP less than 120 days by 2050 (Map 2.3). Cropping is very difficult in places with an LGP less than 120 days. For example maize is considered marginal in areas with LGP between 121 and 150 (Nachtergaele et al 2002 in Jones and Thornton 2008). Grazing area can also be lost as LGP decreases. Mexico, northeast Brazil, a strip across the African Sahel, Morocco, areas of Southern Africa, and parts of India are highlighted as hotspots with this threshold.



Map 2.3. Areas that will flip from LGP >120 days in the 2000s to LGP < 120 days by 2050

3. Areas that flip from more than 90 reliable crop growing days (RCGD) per year in the 2000s to less than 90 RCGD by 2050 (Map 2.4). RCGD estimates the total number of reliable growing days over multiple seasons, for those regions with multiple cropping seasons. It also incorporates the changing probability of crop failure. Ninety RCGD is the equivalent of 120 day LGP, so when RCGD drops below 90 days cropping becomes very difficult. RCGD is a more discriminating indicator than LGP for rain fed agricultural crops. The area highlighted expands with this threshold, to include a range of areas across the global tropics in south Asia, Southern Africa, northeast Brazil, west Mexico, East and West Africa and east Asia.




Map 2.4. Areas that flip from > 90 reliable crop growing days (RCGD) per year in the 2000s to <90 RCGD by 2050

4. Areas where the average annual temperature flips from less than 8°C in the 2000s to more than 8°C by 2050 (Map 2.5). This could expand crop suitability of these areas. The Andes, parts of central and highland south Asia and south China are highlighted.




Map 2.5. Areas where the average annual temperature flips from <8°C in the 2000s to > 8°C by 2050.

5. Areas where average annual maximum temperature will flip from under 30°C to over 30°C (Map 2.6). **While** this is the maximum temperature that beans can tolerate, rice and maize yields suffer at higher temperatures than this, as do other staple crop yields. Grazing vegetation will also suffer at such high temperatures and we could see switches in species with implications for palatability for livestock. Higher temperatures also affect food safety, for example milk storage, and disease transmission patterns, such as malaria. Significant areas of Latin America, Africa and Southeast Asia are highlighted.



Map 2.6. Areas where average annual maximum temperature will flip from < 30°C to > 30°C .


6. Areas where the maximum temperature during the primary growing season is currently less than 30°C but will flip to more than 30°C by 2050, during the primary growing season (defined as the longest for a given area) (Map 2.7). This shrinks the highlighted area somewhat, but these areas are more vulnerable to increased riskiness of cropping because it is specific to the primary growing season for a given area.



Map 2.7. Areas where maximum temperature during the primary growing season is currently < 30°C but will flip to > 30° C by 2050 (during the primary growing season).


The next three thresholds attempt to characterize how climate change may affect variability. Many believe that climate change will increase variability (Easterling et al 2007 p 283), although at the moment climate modellers cannot predict this with any accuracy.

7. We first show where coefficient of variability of rainfall is currently high (Map 2.8), as increases would make cropping even riskier in such areas. In most heavily cropped areas (see Map 2.1) CV of rainfall is less than 25%, with the exception of India. There is very little cropping in areas with CV of rainfall greater than 45%. But large areas of Africa, south Asia, Mexico, and the Middle East have CV greater than 25%.




Map 2.8. Areas where CV of rainfall is currently high.

An analysis of percent area cropped and CV of rainfall shows that the majority of cropped areas fall within a CV range of 12% to 37%, with the mode occurring at 21%. We can use this to divide areas of the tropics as shown in Map 2.9, below. Those croplands with rainfall CV less than 21% are in white and rainfall CV more than 21% are in red. Large areas of the tropics are already in the over 21% zone, including heavily cropped areas such as south Asia, Mexico, Southern Africa, and northern Nigeria.




Map 2.9. Areas where CV of rainfall is more than 21%.

8. Changes in variability could mean less rain per rainfall event. Map 2.10 shows areas where rainfall per day decreases by 10% or more between 2000 and 2050. In areas that are already arid or semi-arid this poses a significant problem for cropping or livestock grazing.




Map 2.10. Areas where rainfall per day decreases by 10% or more between 2000 and 2050.

9. Conversely, in Map 2.11 below, we identify where the amount of rainfall per rain day increases by 10% between 2000 and 2050. This is a proxy for increased rainfall intensity, which can cause soil erosion and greater runoff and therefore limit the effectiveness of rainfall or increase flooding.



Map 2.11. Areas where rainfall per rain day increases by 10% between 2000 and 2050.

In a last step of the climate threat analysis, we classified areas by the number of climate change thresholds identified (excluding LGP decrease by more than 5%). For each pixel, the number of potential climate threats was calculated. In case the pixel is exposed to a positive temperature flip (from less than 8° C to more than 8° C), we lowered the number of threats by one.



Map 2.12. Number of identified climate change thresholds

Southern Africa has the largest area (across Namibia, Angola, Zambia, Botswana, Mozambique and South Africa) with multiple threats, followed by northeastern Brazil, Mexico, Guyana, Nicaragua, and small areas in Tanzania, Ethiopia, the DRC, Uganda, India, and Pakistan, as well as the Middle East.


We calculated domains for each individual threshold, rather than this combined one, as each threshold represents a different climate change impact, and hence the vulnerability domain size and implications differed. For a targeting exercise, exploring these differences is useful.

Risk Maps

The UNEP GRID project has compiled maps of global disaster risk (UNEP 2011). We show the maps for drought and flood here.

Drought

The risk of drought is an important factor in considering the potential for food insecurity because droughts reduce food availability through their impact on local production. Map 2,13 below shows the frequency SPI (Standardized Precipitation Risk), which is defined as the average number of drought events per year per pixel (for the period 1974-2004), where drought events are identified as three consecutive months with less than 50% of precipitation as compared with average.



Map 2.13. The frequency SPI defined as the average number of drought events per year per pixel (for the period 1974-2004).

Flood

The risk of flood is also important when considering the potential for food insecurity because floods destroy crops and reduce food availability. Map 2.14 below shows average flood frequency based upon data from the Dartmouth Flood Observatory. The Global Flood Hazard Frequency and Distribution is a 2.5 by 2.5 minute grid derived from a global listing of extreme flood events between 1985 and 2003 (poor or missing data in the early to mid 1990s), compiled by Dartmouth Flood Observatory and georeferenced to the nearest degree. The resultant flood frequency grid was then classified into 10 classes of approximately equal number of grid cells. The greater the grid cell value in the final data set, the higher the relative frequency of flood occurrence. The dataset is a result of the collaboration between the Center for Hazards and Risk Research, and the Columbia University Center for International Earth Science Information Network. South and Southeast Asia are highlighted in particular.



Map 2.14. Average flood frequency based upon data from the Dartmouth Flood Observatory.

In section four of the report, these maps are compared with future drought and intense rainfall likelihoods.

We did not map areas at risk of sea level rise, but Nicholls et al (2007) suggest that certain coastal environments are at risk: deltas, coral reefs, low-lying coastal wetlands, small islands, and soft areas of coastline. They identify south and Southeast Asian deltas and the Nile delta as particularly vulnerable because of high population densities.


Section 3: Food security maps across the global tropics

An expanded data set was collected and is shown in Appendix 1. For the mapping exercise, we were limited by the indicators available for the countries of interest - most are national level data. The indicators are described and mapped by the food security component each one represents. We endeavoured to map at least two indicators per food security component. Note that these are indicators of food security outcomes rather than drivers of food (in)security. In this section maps for the entire global tropics are shown. Regional maps are in Section 5.

Availability indicators


Current crop yields

Current crop yields are mapped using data from the International Food Policy Research Institute (You et al 2000). These figures represent food availability, the first pillar of food security. Yields are mapped by pixel across the global tropics for maize, rice, millet, sorghum, beans, cassava and wheat for the years 1999-2001.




Map 3.1. Maize yields mapped by pixel across the global tropics.

Map 3.1 illustrates the low maize yields across much of Africa (except parts of Southern Africa and Cameroon), south Asia, and much of Southeast Asia. Yields are higher in China, south India and much of Latin America. Note that this does not tell us about preferences for maize, which are, for example, high in Kenya, Tanzania and across Southern Africa.



Map 3.2. Rice yields mapped by pixel across the global tropics

Again, yields are low for rice across most of Africa and central India. Yields are variable across Southeast Asia and Latin America but higher generally than in Africa.



Map 3.3. Millet yields mapped by pixel across the global tropics.

Map 3.3 shows the low millet yields across all of the tropics, except parts of China. Note that millet is not grown in Latin America.




Map 3.4 Bean yields mapped by pixel across the global tropics.

Map 3.4 shows that beans are extensively grown across south and Southeast Asia, as well as much of Latin America. They are hardly grown in West and Central Africa. Yields only exceed 2.5 t ha in east China and parts of North Africa.

Wheat (see Map 3.5) is hardly grown in Africa or Brazil, except in the highland areas. It is extensively grown across central and south Asia, as well as China. Yields range from less than 1 t ha to 5-10, with notable spots of high productivity in India, China, Egypt, Zimbabwe and Mexico.




Map 3.5. Wheat yields mapped by pixel across the global tropics.



Map 3.6. Sorghum yields are mapped by pixel across the global tropics.

The next crop mapped (Map 3.6) is sorghum. In addition to the regional differences where it is grown, the yield differences are notable. Yields are higher in China and Latin America than in Africa or India.




Map 3.7. Cassava yields mapped by pixel across the global tropics.

The last crop mapped is cassava (Map 3.7), which is extensively grown in Latin America and Southeast Asia, as well as some parts of Africa. It is an important food security crop as it grows in marginal conditions.

Net food production index number per capita

The per capita net food production index number (PIN) is a national level indicator with data obtained from the FAO statistics division, FAOSTAT. A country's per capita PIN illustrates the relative level of the aggregate volume of food production per capita for each year in comparison with the base period 1999–2001. The category of food production includes commodities that are considered edible and that contain nutrients. As such, coffee and tea, although edible, are excluded, along with inedible commodities, because they have little to no nutritive value. This indicator is included to demonstrate the pillar of availability of food from national production, as it monitors trends in production per capita. If the index is above 100, production is increasing relative to the base year. The average for 2003–2007 is shown below.




Map 3.8. Average food production index 2003-2007.

Map 3.8 shows interesting differences by country, as well as by region. Africa has the most countries with PIN less than or equal to 105. In Latin America, Panama, Columbia, Venezuela, Suriname and Guyana have stagnant PIN, while Mexico, Brazil, Bolivia and the rest of Central America (except El Salvador and Belize) have PIN above 100. Much of Central and Southern Africa have PIN below 105, while several countries in East and West Africa are above 105. China and Southeast Asia are also above 105 for the most part, while the countries of the Indo-Gangetic plains are around 100. Very few countries have high growth (over 125).

Access indicators

GDP per capita (current USD)

GDP per capita is the gross domestic product of a country divided by its midyear population. Data are shown for 2005 and come from World Bank national accounts data and OECD national accounts data files. The information was downloaded from the World Development Indicators database of the World Bank databank, World Development Indicators, (<http://databank.worldbank.org>). This is a national level indicator that reflects the ability of consumers to purchase food, as it is a proxy for available income per capita.




Map 3.9. GDP per capita 2005.

GDP per capita is also variable across countries and regions (Map 3.9). In Latin America, Mexico and Venezuela have GDP per capita over USD 5000, while Nicaragua is between USD 500 and 1000. In Africa, there are a number of countries with GDP below USD 500; South Africa, Guinea Bissau, Namibia, Gabon, Angola, Tunisia, Morocco, Libya, Botswana and Algeria have GDP higher than USD 1000. In south Asia, India and Pakistan have GDP of USD 500–1000, contrasting with Nepal and Bangladesh. In Southeast Asia, Vietnam, Cambodia, Laos and Papua New Guinea are all below 1000.

Current poverty levels: % population living below USD 2 a day

Percent of population living with less than USD 2 per day is a poverty indicator mapped by the CGIAR CSI (Wood et al 2010). Current poverty levels are an indicator of food insecurity because they point to the income people have to spend on food and hence their vulnerability to an increase in the price of food. It is well known that the poorer a household is the greater the percent of income spent on food. This is another indicator of economic access to food (affordability).




Map 3.10. Population living on less than USD 2 per day.

As these data are available at the sub-national level (for those countries reporting), more heterogeneity can be analyzed (see the regional maps in Section 5). In Map 3.10 note that a country such as South Africa, which has a high GDP, also has many people living below the poverty line. Regionally, sub-Saharan Africa, south and Southeast Asia and much of west China stand out, for having more than 60% of the population living on less than USD 2 per day. Within Latin America, Nicaragua, Brazil, Bolivia and Venezuela have significant portions of the country with greater than 40% below the USD 2 per day threshold. In Africa, Sudan and Tunisia stand out for their wealth; note that these data mask within country differences. In Southeast Asia, Thailand and Malaysia are wealthier.

Transport time to markets

The time that food takes to reach markets affects households' ability to buy it (Nelson 2008). In Map 3.11 below, within-region differences are most noticeable; for example East Africa (especially Ethiopia and Southern Sudan) has fewer urban centres than West Africa, while India stands out for its high level of urbanization. This is one indicator of physical access to food, but it assumes that closer proximity necessarily increases food security, which is not the case. If other issues such as poverty or high HIV incidence affect food security then physical access will not make a difference. The increasing data about urban food insecurity illustrate this point. This is why no single indicator of food security can explain food security status overall.



Map 3.11. Transport time to markets.


Monthly staple food prices

Monthly prices of staple foods from the capital city of each focus country are graphed to illustrate price variability over time at the sub-national level. Prices have been obtained from FAO's Global Information and Early Warning System (GIEWS) price tool (www.fao.org/giews/pricetool/). The nominal monthly prices were converted to real prices using consumer price indices (CPI) of each country from the World Development Indicators database (<http://databank.worldbank.org>). Consumer price indices reflect changes in the cost to the average consumer of acquiring a basket of goods and services. It is generally assumed that greater price volatility seasonally or annually affects low income consumers' ability to access sufficient food. As we were only able to get data for the national capitals these data are discussed in the regional reports.

Utilization indicators

Malnutrition prevalence - stunting

Prevalence of child malnutrition is the percentage of children under 5 whose height for age is more than two standard deviations below the median for the international reference population - known as stunting (ages 0-59 months). For children up to two years old height is measured by recumbent length, and for older children height is measured by stature while standing. The data are based on the World Health Organization's new child growth standards released in 2006 and downloaded from the World Development Indicators database. Stunting is a lagging indicator and reveals chronic food insecurity within a population. Food insecurity does not always lead to undernourishment.




Map 3.12. Stunting prevalence.

In Map 3.12 sub-Saharan Africa, south and Southeast Asia stand out among the regions for having most countries reporting stunting prevalence greater than 40%.

Malnutrition prevalence - wasting

Prevalence of child malnutrition is the percentage of children under age 5 whose weight for age is more than two standard deviations below the median for the international reference population (ages 0-59 months) – known as wasting (Map 3.13). The data are based on the WHO's new child growth standards


released in 2006 and also downloaded from the World Development Indicators database. As with stunting, low weight for age (underweight) indicates chronic food insecurity. The Millennium Development Goal on reducing hunger tracks countries' progress in reducing this prevalence by half (UN 2010).



Map 3.13. Wasting prevalence.

Wasting is a stricter indicator than stunting because it indicates a more severe level of malnutrition.

The total number of malnourished children per square kilometre is shown in the Map 3.14 below. Malnourished is defined as above for wasting (Herrero et al 2009).




Map 3.14. Malnourished children per sq km.

Map 3.14 shows the density of hungry people: most of south Asia, Malawi, Ethiopia, Uganda, parts of Kenya and Tanzania, much of West Africa, and large areas of China, Vietnam, Laos, Indonesia, Guatemala and Nicaragua.

Population using unimproved water source

The percentage of a country's population using improved water sources can act as an indicator of the utilization aspect of food security, as contaminated water often spreads diseases that affect the body's ability to make use of food consumed. In Map 3.15 much of sub-Saharan Africa stands out for having more than 30% of the population reliant on unimproved water, as does Afghanistan and Papua New Guinea. Data for this indicator were downloaded from the WHO/UNICEF Joint Monitoring Program (JMP) for Water Supply and Sanitation, (www.wssinfo.org/data-estimates/table/).



Map 3.15. Population using unimproved water source.

Unimproved sources of drinking water are defined as:


- Unprotected spring. This is a spring that is subject to runoff, bird droppings, or the entry of animals. Unprotected springs typically do not have a "spring box".
- Unprotected dug well. This is a dug well for which one of the following conditions is true: 1) the well is not protected from runoff water; or 2) the well is not protected from bird droppings and animals. If at least one of these conditions is true, the well is unprotected.
- Cart with small tank/drum. This refers to water sold by a provider who transports water into a community. The types of transportation used include donkey carts, motorized vehicles and other means.
- Tanker-truck. The water is trucked into a community and sold from the water truck.
- Surface water is water located above ground and includes rivers, dams, lakes, ponds, streams, canals, and irrigation channels.
- Bottled water is considered to be improved only when the household uses drinking water from an improved source for cooking and personal hygiene; where this information is not available, bottled water is classified on a case-by-case basis. (WHO / UNICEF 2010)

Resource pressure

We also mapped two indicators of resource pressure, as a possible indicator of future vulnerability.

Population growth rate

The annual population growth rate is based on the de facto definition of population, which counts all residents regardless of legal status or citizenship, except refugees not permanently settled in the country of asylum, as they are generally considered part of the population of the country of origin. Information was downloaded from the World Development Indicators database. This is an indicator of resource pressure, which can affect future food security if agricultural productivity does not meet population growth and trade in food does not rise to meet food needs. Much of sub-Saharan Africa reports annual growth rates of over 2%, as do Pakistan and Afghanistan, as well as Guatemala, Papua New Guinea and the Gulf states (Map 3.16).



Map 3.16. Annual population growth.

Agricultural area per capita

This indicator has been calculated using data from FAOSTAT on agricultural area and national population (FAOSTAT 2011). Agricultural area (expressed in 1000 hectares) is the sum of areas under arable land², permanent crops³ and permanent meadows and pastures⁴. When divided by population, this indicates a country's available resources for producing its own food and long-term trends reflect the amount of resource pressure being exerted by growing populations. South and Southeast Asia have very low arable land per capita; Ethiopia, Uganda, China, Cameroon, the DRC, Benin, Guatemala, Honduras El Salvador and Panama are also less than 0.5 (Map 3.17).



Map 3.17. Arable land per capita.

² Land under temporary agricultural crops, temporary meadows for mowing or pasture, land under market and kitchen gardens and land temporarily fallow (less than five years). The abandoned land resulting from shifting cultivation is not included in this category. Data for arable land are not meant to indicate the amount of land that is potentially cultivable.

³ Land cultivated with long-term crops which do not have to be replanted for several years (such as cocoa and coffee); land under trees and shrubs producing flowers, such as roses and jasmine; and nurseries (except those for forest trees, which should be classified under 'forest').

⁴ Land used permanently (five years or more) to grow herbaceous forage crops, either cultivated or growing wild (wild prairie or grazing land).

Section 4: Vulnerability domains

In this part of the report, we attempted to overlay the climate change hotspots with the food insecure hotspots. We took a 'domain' approach to vulnerability, overlaying indicators for exposure, sensitivity and coping capacity (using the definition of vulnerability given earlier) and then classifying areas of the tropics (or domains) accordingly. Although we collected data for a number of food security outcomes, for this exercise we used only one. We strongly recommend using other indicators for regional, national and sub-national analyses of vulnerability.

Based on 3 criteria, we constructed 8 vulnerability domains:

Table 2. Vulnerability domains based on exposure, sensitivity and coping capacity.

Domain	Exposure	Sensitivity	Coping capacity
HHL	High	High	Low
HHH			High
HLL		Low	Low
HLH			High
LHL	Low	High	Low
LHH			High
LLL		Low	Low
LLH			High

The criteria and their thresholds

Exposure

We used the nine different climatic thresholds (see Section 1), with one addition to indicate climate change exposure. For each threshold, if there was no change or flip the area was classified as having low exposure; if there was a change or flip then exposure was called high.

For example: the reliable crop growing days flip. Areas in red are 'high' exposure; everywhere else is 'low'.



Map 4.1. Reliable crop growing days flip to less than 90 days


The different exposure metrics used are:

1. Lgpdelt: more than 5% decrease in length of growing period (LGP).
2. Lpgflip: LGP flips from more than 120 days to less than 120 days (threshold 1).
3. Rcgd: reliable crop growing days (RGCD) flip from more than 90 days to less than 90 days (threshold 2)
4. Tmax: maximum temperature flips from <30 deg C to > 30 deg C (threshold 4)
5. Tgrow: maximum temperature during the growing season flips from <30 deg C to > 30 deg C (threshold 5).
6. Rdaydec: rainfall per rainy day decreases > 10% (threshold 6)
7. Rdayinc: rainfall per rainy day increases > 10% (threshold 7)
8. CV: coefficient of variability of rainfall currently greater than 21%
9. Tmean: mean annual temperature flips from < 8 deg C to >8 deg C (threshold 3)

These each yield different exposure domains (see the maps in Section 2).

Sensitivity


Areas with more dependence on agriculture (both cropping and livestock based) are assumed to be more sensitive to a change in climate. Therefore, the greater an area is cropped (whether planted or grazed), the higher the sensitivity of that area. Based on the Ramankutty et al (2008) dataset we used percentage cropping as a proxy for sensitivity: areas having greater than 16% of the pixels were classified as under cropping (the mode for the global tropics) and considered highly sensitive.



Map 4.2. Areas with greater than 16% cropping (assumed sensitive to change in climate).

Coping capacity


We considered that chronic food insecurity could be a proxy for coping capacity, as an inability to tackle chronic food insecurity indicates a number of institutional, economic and political problems. We used stunting with a threshold of 40% prevalence as the cutoff between high and low (maps for 30% and 50% also available). (Note that data are unavailable for a few countries, appearing as white in the domain maps).



Map 4.3. Areas with high and low coping capacity for a change in climate.

The vulnerability domains, with different exposure variables

For each domain, we also calculated the vulnerable area (in km²) and the number of vulnerable people. The highest vulnerability is the domain HHL: high exposure, high sensitivity and low coping capacity. The least vulnerable is the domain LLH: low exposure, low sensitivity and high coping capacity. The red and orange colours are for the high exposure areas; green and blue are for low exposure. The paler colours are high capacity.




Map 4.4. Exposure 1: Areas where there is greater than 5% change in LGP.

This threshold includes large portions of the global tropics. India, parts of China, Southeast Asia, and West Africa, along with areas of East and Southern Africa and South America fall into the HHL and HHH categories. The number of people in the most vulnerable category (HHL) is 369.1 million, and the area is over 5 million km².

Table 3. Area and population included in the vulnerability domain exposure 1

Domain	Area (000 Km ²)	Population (million)
LLL	3,652	38.6
LLH	10,577	93.8
LHL	1,412	89.9
LHH	3,322	219.7
HLL	10,506	118.3
HLH	15,725	141.4
HHL	5,173	369.1
HHH	5,076	238.3




Map 4.5. Exposure 2: LGP flips from more than 120 days to less than 120 days.

This is a more restrictive exposure threshold, with only a few areas in south Asia in the HHL category, parts of Mexico in the HHH category, and areas of Africa in the HLL category. The population in the HHL (most vulnerable) category is 7.9 million, covering an area of only 81,000 km²

Table 4. Area and population included in the vulnerability domain exposure 2

Domain	Area (000 Km ²)	Population (million)
LLL	13,951	155.5
LLH	26,018	232.3
LHL	6,504	451.2
LHH	8,248	452.8
HLL	206	1.4
HLH	284	3.0
HHL	81	7.9
HHH	150	5.2




Map 4.6. Exposure 3: RCGDs flip from more than 90 days to less than 90 days (threshold 2).

This exposure threshold includes a slightly larger area, with more of each continent appearing in the high exposure category. However most of these areas are in the low sensitivity category, meaning less than 16% of the area is cropped. The population in the HHL category is 11.7 million.

Table 5. Area and population included in the vulnerability domain exposure 3

Domain	Area (000 Km ²)	Population (million)
LLL	13,745	152.9
LLH	25,625	228.0
LHL	6,424	447.4
LHH	8,129	446.7
HLL	412	3.9
HLH	677	7.3
HHL	161	11.7
HHH	269	11.3




Map 4.7. Exposure 4: Maximum daily temperature flips from <30 deg C to >30 deg C (threshold 4).

This is a more inclusive exposure threshold, with more of the tropics appearing in the high exposure category, especially in Africa. Interestingly, most of India is not highly exposed under this threshold, although part of the Indo-Gangetic Plains is in the HHL category. The population in the HHL category is 44.6 million.

Table 6. Area and population included in the vulnerability domain exposure 4

Domain	Area (000 Km ²)	Population (million)
LLL	10,782	130.6
LLH	22,165	214.8
LHL	5,952	414.4
LHH	7,440	430.5
HLL	3,375	26.3
HLH	4,137	20.5
HHL	633	44.6
HHH	957	27.5




Map 4.8. Exposure 5: Maximum daily temperature during the growing season flips from <30 deg C to > 30 deg C (threshold 5).

This category is obviously more restrictive than the previous thresholds. However new areas in India and China and West Africa, appear in the HHL or HHH categories, as well as Central America. The population in the HHL category is 55.8 million.

Table 7. Area and population included in the vulnerability domain exposure 5

Domain	Area (000 Km ²)	Population (millions)
LLL	12,531	140.3
LLH	24,704	221.7
LHL	5,697	403.2
LHH	7,873	426.1
HLL	1,626	16.6
HLH	1,598	13.6
HHL	888	55.8
HHH	525	31.9




Map.4.9. Exposure 6: Rain per rainy day decreases by more than 10% (threshold 6).

This threshold is also quite restrictive in terms of land area, but it is interesting to note the areas in Nigeria and India that fall into the HHL category. The other highly exposed areas are largely in the low sensitivity domain, as not much is cropped. The population in the HHL category is 27.5 million.

Table 8. Area and population included in the vulnerability domain exposure 6

Domain	Area (000 Km ²)	Population (millions)
LLL	13,749	151.6
LLH	25,073	226.4
LHL	6,296	431.6
LHH	8,097	440.7
HLL	408	5.3
HLH	1,229	8.9
HHL	289	27.5
HHH	301	17.3




Map 4.10. Exposure 7: Rain per rainy day increases by more than 10% (threshold 7)

This exposure includes a number of more densely populated areas with low coping capacity, again in West and Southern Africa, and south and Southeast Asia. The vulnerable population is 45.7 million.

Table 9. Area and population included in the vulnerability domain exposure 7

Domain	Area (000 Km ²)	Population (millions)
LLL	11,709	135.5
LLH	24,473	213.1
LHL	5,679	413.4
LHH	7,911	424.7
HLL	2,449	21.4
HLH	1,828	22.2
HHL	906	45.7
HHH	486	33.3




Map 4.11. Exposure 8: CV rainfall currently greater than 21%.

This is a very inclusive exposure threshold, with most of Africa and south and east Asia included in the high exposure category. The population in the HHL category increases to 270.4 million.

Table 10. Area and population included in the vulnerability domain exposure 8

Domain	Area (Km ²)	Population (millions)
LLL	7,298	86.6
LLH	13,337	141.0
LHL	2,614	188.7
LHH	4,519	268.8
HLL	6,859	70.2
HLH	12,965	94.3
HHL	3,971	270.4
HHH	3,878	189.2



Map 4.12. Exposure 9: Mean annual temperature flips from less than 8 deg C to more than 8 deg C.

This is a different type of exposure from the other eight, as these areas represent a lifting of current temperature constraints (average temperature too low), and so cropping potential may increase. Thus the domains to focus on are HLH or HHH, on the map indicated by “positive change”. The combined population in these two is 2.9 million and concentrated in the Andes and central China.


Table 11. Area and population included in the vulnerability domain exposure 9

Domain	Area (000 Km ²)	Population (millions)
LLL	14,124	156.3
LLH	26,119	233.3
LHL	6,573	458.9
LHH	8,380	457.1
HLL	33	0.6
HLH	183	2.0
HHL	13	0.15
HHH	17	0.89

Section 5: Regional maps

For targeting within regions, we also made regional maps, to highlight the variation within them (as opposed to among) more easily. Here we show the current CCAFS regions: East Africa, West Africa, and south Asia. The other regional maps are available as needed. The thresholds and food security indicators are as defined in the global maps.


East Africa



Map 5.1. Threshold 1 (Length of growing period flips from more than 120 days to less than 120 days);


Map 5.2. Threshold 2 (Number of reliable crop growing days flips from more than 90 to less than 90).

These maps (5.1 and 5.2) both indicate increased cropping risk. Threshold 2 (Map 5.2) includes a range of areas throughout East Africa.




Map 5.3. Threshold 4 (maximum temperature flips to more than 30 deg C). Map 5.4. Threshold 5 (maximum temperature in the growing season flips to more than 30 deg C).

Threshold 5 (Map 5.4) is more restrictive, but northern Uganda, parts of Ethiopia and central Kenya are highlighted.




Map 5.5. Threshold 6 (Average rainfall per rainy day increases by more than 10%). Map 5.6. Threshold 7 (Average rainfall per rainy day decreases by more than 10%).

Threshold 6 (Map 5.5) is much more common than threshold 7 (Map 5.6).




Map 5.7. Coefficient of rainfall variability (mode is 21% for global tropics)




Map 5.8. Percent area cropped

Note in Map 5.8 the high cropping density in southern Sudan, the highlands of Ethiopia and much of Uganda relative to Tanzania and Kenya. The eastern areas have higher CV of rainfall (Map 5.7).



Map 5.9. Economic access: % population on less than USD 2 a day



Map 5.10. Physical access: travel time to nearest town >250K

Kenya has lower poverty rates than Ethiopia, Tanzania, Uganda, Burundi and Rwanda (Map 5.9).

Physical access is restricted in much of Ethiopia, north Kenya, north Somalia and parts of Tanzania (Map 5.10).




Figure 3. Price volatility Nairobi.




Figure 4. Price volatility Kampala.






Figure 5. Price volatility Addis Ababa.

Figures from 2005 to 2009; prices are adjusted for inflation.

Maize prices are more stable than bean or rice prices. All three graphs in Figure 3 show the 2008 price spike.




Map 5.11. Kilocalories available per person per day



Map 5.12. Stunting prevalence (% under five)

Kilocalorie availability is lower in Uganda, Sudan and Somalia (Map 5.11). However stunting prevalence is highest in Ethiopia, followed by Burundi. Kenya has the lowest stunting prevalence (Map 5.12).


West Africa



Map 5.13. LGP flip.

Map 5.14. RCGD flip.


Thresholds 1 and 2. The length of growing period flip is restricted to a band across Senegal, southern Mali and Burkina Faso, with a bit in north Nigeria (Map 5.13). RCGD flip includes many more scattered areas throughout the region south of the Sahara (Map 5.14).



Map 5.15. Tmax flip.

Map 5.16. Tmax growing season flip.


Thresholds 4 (Map 5.15) and 5 (Map 5.16). Threshold 5 is common in the southern parts of the region, along the coast, much of Cameroon and the Central African Republic. Note that this overlaps with intensive cropping.




Map 5.17. Rainfall per rainy day increase.

Map 5.18. Rainfall per rainy day decrease.

Thresholds 6 and 7. Rainfall intensity may increase in the northern areas of Nigeria and much of Niger (Map 5.17). Rainfall per rainy day may decrease in the lower parts of Nigeria and across central Senegal (Map 5.18).




Map 5.19. Percent area cropped




Map 5.20. Coefficient of rainfall variability (mode is 21% for global tropics)

The CV of rainfall increases as one moves north (Map 5.20); some of this area is currently cropped quite intensively along the Niger/Nigeria border (Map 5.19). Map 5.19 also shows the intensive cropping throughout Nigeria, Ghana, Togo, Benin and Cote D'Ivoire, as well as northern Cameroon and eastern Senegal and the Gambia. Many of the climate change exposure thresholds coincide with these areas. It is notable the LGP flip is concentrated in the high CV rainfall band.



Map 5.21. Economic access: % population on less than USD 2 per day



Map 5.22. Physical access: travel time to nearest town >250K

As seen in Map 5.21 there are high concentrations of poverty across the region. The difference between urban and rural areas in Nigeria is stark while Cote d'Ivoire and Cameroon are wealthier. There are also sub-national differences in Nigeria, Mali, Burkina Faso and Ghana. Physical access is good in many of the cropped areas of the region (Map 5.22).





Figure 6. Price volatility Accra.

Figure 7. Price volatility Bamako.

Rice is more expensive than millet, sorghum, beans or maize, but there are no major differences in the volatility of prices. The impact of the food price crisis of 2008 can also be seen (Figs 6 & 7).






Figure 8. Price volatility Niamey.

The spike of 2005 was huge in Niamey, even relative to 2008 (Fig 8).




Map 5.23. Kilocalories available per person per day



Map 5.24. Stunting prevalence (% under five)

As seen in Map 5.23 kilocalorie availability is lowest in Chad, Niger, and the Central African Republic. Stunting (Map 5.24) is over 50% in Niger and is greater than 40% in Nigeria, Togo, Cote d'Ivoire, Liberia, Sierra Leone, Mali, Chad and the Central African Republic. Senegal, the Gambia and Benin have rates of 20% or lower.


South Asia



Map 5.25. LGP flip.

Map 5.26. RCGD flip.


Thresholds 1 and 2. As in the other regions, the LGP flip is the most restrictive threshold, with concentrations in the Indo-Gangetic Plains, southern Pakistan, and western India (Map 5.25). The RCGD flip includes more of south India, Nepal, and west India (Map 5.26).



Map 5.27. Tmax flip.

Map 5.28. Tmax growing season flip.



Thresholds 4 and 5. Tmax flip covers central Pakistan, the northern Indo-Gangetic Plains, and parts of east and west India (Map 5.27). Tmax growing season flip is more extensive across central India, Pakistan, the northern Indo-Gangetic Plains, and eastern India (Map 5.28).



Map 5.29. Rainfall per rainy day increase.

Map 5.30. Rainfall per rainy day decrease.



Threshold 6 and 7. Rainfall intensity is likely to increase in western India, Nepal and parts of east India (Map 5.29). Central and north India are likely to see decreases in rainfall amounts per rainy day (Map 5.30). Bangladesh is not highlighted under any of the thresholds.



Map 5.31. Coefficient of rainfall variability.


Map 5.32. Percent area cropped.

India, Pakistan and Bangladesh are all intensively cropped (Map 5.32). Pakistan already has very high CV of rainfall, along with northwest and south India (Map 5.31).




Map 5.33. Economic access: % population on less than USD 2 per day. Map 5.34. Physical access: travel time to nearest town >250K.

Nepal, Bangladesh and much of India have greater than 60% of the population living on less than USD 2 per day (Map 5.33). Bangladesh, India and central Pakistan are highly urbanized (5.34), in contrast to Nepal Map.



Map 5.35. Kilocalories available per person per day.



Map 5.36. Stunting prevalence (% under five).

Bangladesh has the lowest kilocalorie availability in the region (Map 5.35) while Bangladesh and Nepal have stunting prevalence over 50% (Map 5.36).

Price volatility 2005-2009 (real prices)




Figure 9. Price volatility Bangladesh




Figure 10. Price volatility Delhi.

Wheat prices are very stable in Delhi, rice less so. Bangladesh was clearly affected by the 2008 price spike.

Conclusion: Analysis of hotspots and implications for CCAFS research

This project set out to identify hotspots of food insecurity and climate change. We have done this for the global tropics as a whole, as well as within specific regions, including those currently targeted by the CCAFS programme.

Climate change hotspots

Nine climate change thresholds were defined. The major implications of these are:

- Length of growing period declines by 5% or more across a broad area of the tropics, including heavily cropped areas of Mexico, Brazil, Southern and West Africa, the Indo-Gangetic Plains, and Southeast Asia. This suggests that at a minimum, most of the tropics will experience a change in growing conditions that will require adaptation to current agricultural systems.
- Length of growing period flips to less than 120 days in a number of locations across the tropics, notably in Mexico, northeast Brazil, Southern and West Africa and India. This is a critical threshold for a number of crops as well as rangeland vegetation, and hence these areas are important to target for high exposure to climate change.
- Reliable crop growing days decrease to critical levels below which cropping might become too risky to pursue as a major livelihood strategy in a large number of places across the global tropics, including West Africa, East Africa, and the Indo-Gangetic Plains. If we use this threshold as an indicator of where CCAFS might move next, there are large and troubling areas in Southern Africa, south India, northeast Brazil, and south Mexico. There is good consensus from the various GCMs used (i.e., these results are robust) for West Africa, the Indo-Gangetic Plains, parts of East Africa, and much of Southern Africa and south India. These results seem to be less robust for south Mexico, northeast Brazil, and parts of East Africa (Kenya, specifically).
- High temperature stress (above 30°C) will be widespread in East and Southern Africa, north and south India, Southeast Asia, northern Latin America and Central America. During the primary growing season high temperature stress will be a problem for different areas. These are also important high exposure thresholds, but their implications should be considered in conjunction with cropping system type, dependence upon agriculture, and disease threats.
- Much of the tropics already experiences highly variable rainfall, above the median of 21% for cropped areas. Thus any increase in this variability will make agriculture riskier.
- Reduced rainfall per rain event can be compared to the current drought risk map; it appears that a number of current drought prone areas are included here, such as Southern Africa, West Africa and Central America; however, new areas may also become more drought prone, such as central India.
- A third, different type of hotspot relating to threshold 7, is the increase in rainfall intensity, although without more analysis it is not so easy to interpret what this may mean. East Africa, the Indo-Gangetic Plains, and parts of West Africa may be prone to increased erosion and runoff; these

problems may be widespread in Southern Africa and throughout the neotropics. Some of these areas (eastern India and Bolivia) are currently flood prone.

- When eight of the thresholds were combined, Southern Africa emerged as highly exposed under several thresholds, as did selected areas in each of the regions, including northeast Brazil, Mexico, Pakistan, India and Afghanistan.

Food security hotspots

Eight food security indicators were mapped to identify current food insecurity hotspots. At least two indicators were chosen for each component of food security.

- **Availability:** bean, millet and sorghum yields are low throughout the areas of interest, although millet and sorghum are important dryland crops. The story for maize, wheat and rice is more mixed, but most of Africa (except Southern) has low yields relative to Latin America and southern India. The net food production index is stagnant in all areas of interest, with differences between countries rather than by region.
- **Access:** GDP per capita is low in many countries in Africa, as well as Afghanistan, Nepal, Bangladesh, Laos and Cambodia. Hotspots for poverty are West, Central and East Africa, India and Bangladesh and Southeast Asia. These are national level indicators. Travel time to markets indicates population density as well as infrastructure and urban growth. Regional differences emerge in this map.
- **Utilization:** Africa and south Asia are clearly much more chronically food insecure regions than Latin America or China. We have no data for food preferences, nor did we map sub-national data, which would be more helpful for targeting.
- In terms of resource pressure, again Africa is highlighted for population growth rates. Arable land per capita really differs by country.

Vulnerability domains

The implications of the nine vulnerability domains are as follows:

The most vulnerable domain for most exposures is high exposure, high sensitivity and low coping capacity (HHL). Such areas are highly exposed to climate change and have significant agriculture and high levels of food insecurity. However, we have used the same coping capacity threshold (40% stunting) in all domains. Thus the HHH category is also one to watch, as there are places with other types of food insecurity (such as constrained market access and poverty) and hence low coping capacity. Also, if we used 30% stunting instead, a number of other countries would be included, for example Kenya. Areas in the HLL capacity should also be carefully considered; if agriculture expanded or intensified in these areas their sensitivity could increase.

For now, areas in the low exposure domain are not a priority for CCAFS, but there is considerable variation in the definition of these domains by the exposure thresholds. Thus the majority of the tropics are in high exposure under the threshold “LGP changes by more than 5%”, and about half is included in high exposure for “CV rainfall currently greater than 21%”. This contrasts with the most restrictive threshold, “LGP flips from more than 120 to less than 120 days”. The category to scrutinize is LHL, as these regions are highly sensitive due to having a significant amount of cropped area and being chronically food insecure.

Under exposure 1 (LGP decreases more than 5%), HHL is the category with the most people, followed by HHH. For exposure 2 (LGP flips) HHL is very small in terms of people; most people are in the categories LHL or LHH. Exposure 3 (reliable crop growing days - RGCP flips) has about 10 million more people in the HHL category, but again most people are in LHL or LHH. Under exposure 4 (maximum temperature -Tmax flips), the vulnerable population more than triples relative to exposure 3. Under exposure 5 (temperature flips) again more people are in the vulnerable categories. Under exposure 6 (rain per rainy day decrease), the most vulnerable population drops to 27.5 million, while under exposure 7 (rain per rainy day increase) 45.7 million are in the HHL category. This suggests that the choice of domain variables makes a big difference in terms of areas included.

Limitations to this analysis:

As with any exercise of this nature, there are a number of ‘health warnings’ to observe. First, these maps only cover current food security, as we are not working with a model that could give us predictive capacity to estimate future food security in the areas of interest. Second, we have primarily used national level food security data, which is a) not always reliable and b) masks within-country variation as well as within-household variation, the latter important to understand the social dimensions. Third, we have not mapped the links between climate change exposure and other food security variables besides crop yields (directly) and utilization (indirectly).

Next steps: there are several next steps that could be taken to improve on the mapping work done here.

1. Vary the indicators for coping capacity and sensitivity. Sensitivity varies by type of cropping system, just as food security impacts differ depending upon the aspect considered.

Use household food security data, such as that collected by the Food Economy Group or NGOs like Save the Children or CARE, to simulate or evaluate the household level food security impacts of climate change. This is best done as part of regional scenario exercises that include vulnerability analysis both currently and under different scenarios of adaptive capacity.

2. Explore disease interactions with climate change.
3. Map drivers of food security and food system vulnerability rather than food security outcomes, using the IPCC framework of exposure, sensitivity and adaptive capacity.
4. Model food security to 2050 under a couple of key scenarios including one or more of the following key drivers (in addition to climate change):
 - Transformation of the rural economy is a key issue: what will the structure of agriculture look like under high urban migration?
 - High versus low economic growth and impacts on adaptive capacity.
 - Government versus private sector as key actors shaping innovations, etc. in food systems.
 - Regionalization versus globalization of markets.

References

- Barrett CB. 2010. Measuring Food Insecurity. *Science* Vol. 327 p. 825-828
- Christiaensen LJ, Boisvert RN. 2000. Validating operational food security indicators against a dynamic benchmark. Paper presented at American Agricultural Economics Association, Tampa, Florida, August 2000.
- Coates J, Wilde PE, Webb P, Rogers BL, Houser RF. 2006. Comparison of a qualitative and a quantitative approach to developing a household food insecurity scale for Bangladesh. *Journal of Nutrition* 136: 1420S-1430S.
- Devereux S. 2007. The impact of droughts and floods on food security and policy options to alleviate negative effects. *Journal of Agricultural Economics* 37 (issue supplement 1): 47-58.
- Easterling WE, Aggarwal PK, Batima P, Brander KM, Erda L, Howden SM, Kirilenko A, Morton J, Soussana J-F, Schmidhuber J, Tubiello FN. 2007. *Food, fibre and forest products. Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE (eds), Cambridge University Press, Cambridge, UK, 273-313
- Ericksen PJ. 2008a. Conceptualizing food systems for global environmental change research. *Global Environmental Change* 18(1):234-245.
- Ericksen PJ. 2008b. What is the vulnerability of a food system to global environmental change? *Ecology and Society* 13(2): Art 14.
- FAO 2011 *The State of Food and Agriculture 2010-2011*, Rome: FAO
- FAO Statistical databases 2011. <http://faostat.fao.org>.
- GECAFS 2005. Science Plan and Implementation Strategy. *GECAFS Report 1*. Wallingford, UK.
- Herrero, M, Thornton PK, Notenbaert A, Msangi S, Wood S, Kruska R, Dixon J, Bossio D, van de Steeg J, Freeman HA, Li X. Parthasarathy Rao P. 2009. *Drivers of change in crop-livestock systems and their impacts on agro-ecosystems services and human well-being to 2030*. CGIAR Systemwide Livestock Programme. International Livestock Research Institute, Nairobi: 101.
- Jones P, Thornton P. 2008. Croppers to livestock keepers: livelihood transitions to 2050 in Africa due to climate change. *Environmental Science and Policy* 12: 427-437.
- Jones P, Thornton P, Heinke J. 2009. Generating characteristic daily weather data using downscaled climate model data from the IPCC's Fourth Assessment. Unpublished report (downloaded from <http://futureclim.info>).
- McCarthy JJ, Canziani O F, Leary N A, Dokken D J, and White K S, editors. 2001. *Climate change 2001: impacts, adaptation and vulnerability. Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK.
- Misselhorn A, Eakin H, Devereux S, Drimie S, Msangi S, Simelton E, Stafford-Smith M. 2010. Vulnerability to What? In Ingram J, Ericksen P, Liverman D, (ed) *Food Security and Global Environmental Change*. Earthscan, London.
- Nelson A. 2008. *Travel time to major cities: A global map of Accessibility*. Global Environment Monitoring Unit - Joint Research Centre of the European Commission, Ispra Italy. Available at <http://www.tem.jrc.it/accessibility>.
- Nicholls RJ, Wong PP, Burkett VR, Codignotto JO, Hay JE, McLean RF, Ragoonaden S, Woodroffe CD, 2007. Coastal systems and low-lying areas. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE (eds). Cambridge, Cambridge University Press: 315-356.
- Ramankutty N, Evan TA, Monfreda C, Foley JA. 2008. Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* 22: GB 1003, doi: 10.120/2007BG002952.
- Randall DA, Wood RA, Bony S, Colman R, Fichetef T, Fyfe J, Kattsov V, Pitman A, Shukla J, Srinivasan J, Stouffer RJ, Sumi A, Taylor KE, 2007. Climate models and their evaluation, in: Solomon S, Qin D, Manning M, Chen, Marquis M, Averyt KB, Tignor M, Miller HL. (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA.

- Riely F, Mock N, Cogill B, Bailey L, Kenefick E. 1999. *Food security indicators and framework for use in the monitoring and evaluation of food aid programs*. Food and Nutrition Technical Assistance (FANTA) Project, Washington D.C.
- Stamoulis K, Zezza A. 2003. A conceptual framework for national agricultural, rural development and food security strategies and policies. *ESA Working Papers* No. 03-17, FAO, Rome.
- UN 2010 Millennium Development Goals Indicators <http://mdgs.un.org/unsd/mdg/Default.aspx>
- UNEP 2011. Global Bank Data Platform. <http://preview.grid.unep.ch/index.php?preview=home&lang=eng>
- Webb P, Coates J, Frongillo EA, Rogers BL, Swindale A, and Bilinsky P. 2006. Measuring household food insecurity: why it's so important and yet so difficult to do. *Journal of Nutrition* 136:1404S-1408S.
- WHO / UNICEF *Joint Monitoring Programme (JMP) for Water Supply and Sanitation 2010*. www.wssinfo.org/definitions-methods/watsan-categories/ Accessed on 26 May 2011.
- Wood S, Hyman G, Deichmann U, Barona E, Guo R Z, Castaño S, Rivera O, Diaz E, Marin J. Additional data and technical contributions from: Coulombe H, Wodon Q, Muniz M, Benin S, Benson T, Berdegue J, Gonzalez J, Barufi A, Lanjouw P, Simler K, Blankespoor B, Murray S. 2010. *\$1 and \$2 day poverty maps of the developing countries. Input to the CGIAR Strategy and Results Framework*. <http://docs.google.com/viewer?a=v&pid=sites&srcid=Y2d4Y2hhbmdlLm9yZ3xhbGxpYW5jZlZxneDoyZmJhZTVhNzBiY2Q3Y2lw>
- World Bank Databank, *World Development Indicators*. <http://databank.worldbank.org>. Accessed on 1 October and 5 November 2010.
- World Health Organization and the United Nations Children's Fund. 2009. *WHO child growth standards and the identification of severe acute malnutrition in infants and children. A Joint Statement by the World Health Organization and the United Nations Children's Fund*. www.who.int/nutrition/topics/malnutrition/en/index.html.
- You L, Crespo S, Guo Z, Koo J, Ojo W, Sebastian K, Tenorio MT, Wood S, Wood-Sichra U. 2000. *Spatial Production Allocation Model (SPAM) Version 3 Release 2*. <http://MapSPAM.info>. Accessed in April 2010
- www.CGIAR.CSI.fao.org/gIEWS/pricetool/). Accessed on 15 and 16 October 2010.
- www.wssinfo.org/data-estimates/table/. Accessed on 30 October 2010.