

Nutrition and Health Surveillance in Chittagong Division

Key results for the period: February 2001 to January 2002

Rural Bangladesh achieved a steady decline in malnutrition during the last decade but there are still millions of malnourished children and mothers leading impoverished lives. Since its inception in 1990 the HKI/IPHN Nutritional Surveillance Project (NSP) has been a source of high quality data on nutrition, health, socio-demography, food security and poverty in rural areas of all six divisions in the country. This bulletin presents NSP data collected in 2001 from rural areas in Chittagong Division. According to international criteria, malnutrition was a 'very high' or 'serious' public health problem in children and mothers. Over one half (58%) of households had an energy intake <2122 kcal/person/d and anemia was a 'serious' problem in both children (40%) and mothers (43%), which suggest that the diet lacked both energy and micronutrients. Investing in nutrition is crucial because malnutrition contributes to poverty and impedes both social and economic development.

The prevalence of malnutrition in Bangladesh is amongst the highest in the world. Although there is an increasing magnitude of problems among the urban poor, 77% of the population still resides in rural areas, where most lack adequate access to food, health care and other basic needs. The Government of Bangladesh, donors and development agencies recognise that malnutrition is a major barrier to poverty alleviation and development in Bangladesh and warrants effective action. In order to design, monitor and evaluate policies and programs, detailed information on the magnitude and determinants of the problem is needed.

Information generated by the NSP over the past 12 years has helped inform policymakers, program managers and donor organizations on many development concerns in Bangladesh including health, nutrition, food security, gender disparities, and poverty. HKI encourages the use of its surveillance data to facilitate policy and program action for the benefit of the population from whom they were collected.

Following the successful experience of the HKI Nutrition and Health Surveillance System (NSS) in data sharing in Indonesia, the data collected by the NSP in Bangladesh in 2001 on key indicators of nutrition, health and their determinants are included in a CDROM that accompanies this series of bulletins.

Data collection in Chittagong Division

Data are collected every 2 mo in rural areas of four sub-districts (*upazila*) in Chittagong Division: Nabinagar, Hathazari, Chaudagram and Cox's Bazar. These four sub-districts were randomly selected when the new sampling design was introduced in February 1998, and provide data that are statistically representative of the division. A new sample of 1,500 households is randomly selected from the total of four sub-districts during each round of data collection as follows. Fifteen *mauza* (administrative unit within a sub-district) are randomly selected from each sub-district. One village is randomly selected from each *mauza*, and 25 households with at least one child aged <5 years are systematically sampled from each village. A precoded questionnaire is used to record data from each household on the health

and nutrition of one mother and all her children aged <5 years; household demography and socio-economic status; household food consumption; household food production; natural disasters, household crises and coping strategies; and participation in NGO programs.

Data are also collected from the five other divisions of the country: Barisal, Dhaka, Khulna, Rajshahi and Sylhet. The data from all six rounds of data collection in the six divisions are aggregated to provide a nationally representative sample of 54,000 rural households each year.

Findings presented

In this bulletin, data are presented on a selection of indicators of household demographic and socio-economic status, household food security (loans for food and homestead gardening), caring practices (breastfeeding), the performance of national health programs (vitamin A capsule distribution), food consumption (household energy intake and consumption of non-cereal foods), health status (child and maternal diarrhea) and nutritional status (child and maternal nightblindness, anemia and anthropometry). The definition and method of data collection of each indicator are described together with the general findings for rural Bangladesh and the specific findings for Chittagong Division. The figures show the data collected in Chittagong Division. While the scope of this bulletin is limited to presenting the findings, they serve to facilitate discussion on the immediate and underlying causes of malnutrition and on ways to address it.

Demographic and socio-economic status (SEE TABLE)

What is indicated. The demographic and socio-economic status of a household determines the extent to which the household unit can adequately feed and care for all members, provide a healthy environment and gain access to health services. These factors are therefore important determinants of the health and nutritional status of household members.

Data collection method. A series of questions is asked to obtain the information shown in the Table.

Findings. *Rural Bangladesh* - Demographic and socio-economic conditions varied considerably between the six divisions. *Chittagong Division* - The level of household crowding was lower than all other divisions, despite having the highest number of household members (with Sylhet Division). The percentage of functionally landless households (75%) was also highest in this division, and the percentage

of households with a female decision-maker (12%) was three times the national rural average.

Loan for food in the last month (SEE FIG 1)

What is indicated. A loan for food is a good indicator of household food insecurity in Bangladesh because it is a coping strategy used when a household is unable to produce or purchase sufficient food for consumption.

Data collection method. The respondent is asked whether the household took a loan to obtain food in the last month, either in cash or in kind.

Findings. *Rural Bangladesh* - Overall, 12% of households took a food loan in 2001, ranging from 9% in Jun/Jul to 14% in Feb/Mar and Oct/Nov. *Chittagong Division* - The percentage was below the national rural average throughout most of the year.

Homestead gardening (SEE FIG 2)

What is indicated. Homestead gardening in Bangladesh has been shown to lower the risk of vitamin A deficiency in preschool children and mothers. Households that practice improved gardening techniques consume non-cereal foods more frequently and have a more diverse diet than other households.

Data collection method. Three types of homestead gardens are distinguished: 'traditional' homestead gardens which produce only gourd and traditional types of vegetables seasonally in scattered plots; 'improved' homestead gardens which produce a wider range of vegetables seasonally in fixed plots; and 'developed' homestead gardens which grow a wide variety of vegetables all year in fixed plots.

Findings. *Rural Bangladesh* - In 2001, 76% of households cultivated traditional gardens, 8% improved gardens and 1% developed gardens. *Chittagong Division* - The percentage of households with improved gardens (11%) was higher than the national rural average, but a smaller percentage had traditional and developed gardens.

Breastfeeding of children aged <60 mo (SEE FIG 3)

What is indicated. Infants and young children should be exclusively breastfed for the first 6 mo of life because of nutritional and health benefits. Breastfeeding should continue well into the second year of life, complemented with nutritious foods from 6 mo of age.

Data collection method. The mother is asked whether her child is currently breastfed.

Findings. *Rural Bangladesh* - Throughout rural Bangladesh almost all mothers breastfed their

children well into the second year of life (97% of children aged 12-14 mo and 88% of children aged 21-23 mo), and many for much longer. *Chittagong Division* - The percentage of breastfed children was similar to the national rural average for the first year of life, but was lower than all other divisions towards the end of the second year, with only 74% of children aged 21-23 mo breastfed.

Vitamin A capsule receipt in children aged 12-59 mo (SEE FIG 4)

What is indicated. Preschool children in Bangladesh need vitamin A supplements because their diet does not supply enough vitamin A and so they are at high risk of illness, blindness and dying due to vitamin A deficiency. The Government of Bangladesh currently aims to give every child aged 12-59 mo a high-dose vitamin A capsule (VAC) containing 200,000 IU twice a year at six-monthly intervals during the National Immunization Days for polio. IVACG/WHO also recommend that children aged 6-11 mo be given 100,000 IU vitamin A twice a year. However, these children are currently not included in the national VAC distribution campaigns because the polio vaccine has a target age of 12-59 mo.

Data collection method. The NSP monitors the coverage of the VAC program during a round of NSP data collection that follows a VAC distribution. During the year 2001, the NSP collected data on coverage of the VAC program in Feb/Mar for the November 2000 VAC distribution and in Jun/Jul for the May 2001 VAC distribution. The mother is asked whether her child received a VAC during the most recent distribution.

Findings. *Rural Bangladesh* - Coverage of the VAC program among children aged 12-59 mo increased from around 50% in the early 1990s to almost 90% in the late 1990s. The coverage in November 2000 and May 2001 was 97% and 96% respectively, exceeding all previous years. *Chittagong Division* - Coverage of the VAC program in November 2000 (97%) was the same as the national rural average, while coverage in May 2001(93%) was lower than all other divisions, albeit still very high.

Household energy intake (SEE FIG 5)

What is indicated. Household energy intake is used as an indicator of household poverty: households with an energy intake <1805 kcal/person/d are considered 'extreme' or 'hard-core' poor, and households with an energy intake of 1805-2122 kcal/person/d are considered 'moderate' or 'absolute' poor¹. Household energy intake is also used as an indicator of household

Table Household demography and socio-economic status in 2001

	Chittagong Division	Rural Bangladesh
No. household members (mean)	6.6	5.5
Household crowding (median no. household members per 100 sq ft)	2.0	2.2
Female decision-maker (%)	12	4
Functionally landless ^a (%)	75	68
Manual labor as main source of income (%)	19	22
Clean source of drinking water ^b (%)	99	98
Closed latrine (%)	59	39
Parent's education ^c	Mother	49
	Father	50

^a Own <50 decimals (2000 m²) cultivable land

^b Water obtained from hand pump, deep tube well or tap

^c At least one year of formal education

Fig 1. Percentage of households that took a loan for food in the last month in 2001




Fig 2. Percentage of households that participated in homestead gardening in 2001




Fig 3. Percentage of breastfed children by age in 2001




Fig 4. VAC coverage in children aged 12-59 mo in 2001





Fig 5. Household energy intake in Dec 2001/Jan 2002



food security. However, this indicator neither incorporates the social or environmental dimensions of poverty, nor does it give an indication of the quality of the household diet.

Data collection method. Since December 2001, the NSP has collected data on household energy intake using a 7-day list-recall method. The respondent is asked to list all food and beverage items consumed by household members in the previous week and to estimate the quantity of each item consumed. These data are converted to a value of daily energy intake per household member.

Findings. Rural Bangladesh - In Dec 2001/Jan 2002, 24% of rural households had an energy intake <1805 kcal/person/d and 23% had an energy intake of 1805-2122 kcal/person/d. **Chittagong Division** - The percentage of households with a low energy intake (<2122 kcal/person/d) was, with Barisal Division, higher than all other divisions (58%). About one-third (32%) of households had an intake <1805 kcal/person/d and were considered 'extremely' poor.

Household consumption of non-cereal foods (SEE FIG 6)

What is indicated. A high quality diet, which contains a diverse range of non-cereal foods that are good sources of micronutrients, reduces the risk of micronutrient deficiencies. One way to assess the quality and diversity of the household diet is to examine how often household members eat non-cereal foods.

Data collection method. The mother is asked to recall on how many days in the last 7 days the household members ate five common non-cereal foods: *dal* (lentils), eggs, green leafy vegetables, yellow/orange fruit or vegetables, and fish. A household is considered to have eaten a non-cereal food 'regularly' if household members ate it on ≥ 4 days in the last 7 days. As data on this indicator were not collected in Dec 2001/Jan 2002, data from the previous year are included in Fig 6 in order to show seasonal variation throughout the year.

Findings. Rural Bangladesh - During most of the year <10% of households ate *dal*, yellow/orange fruits or vegetables, eggs and green leafy vegetables regularly, and 40-70% households ate fish regularly. **Chittagong Division** - Regular consumption of non-cereal foods was similar to the national rural average.

Child and maternal diarrhea (SEE FIG 7)

What is indicated. Diarrhea is a form of morbidity that is relatively easy to monitor, because it occurs relatively frequently and respondents easily

understand its definition. It is also a major cause of undernutrition in developing countries, particularly among young children. The prevalence of diarrhea reflects hygiene conditions both inside the house and in the neighborhood.

Data collection method. The mother is asked a series of questions to determine whether she and her child had diarrhea in the last 24 hr. Diarrhea is defined as ≥ 3 loose, watery or mucoid stools in 24 hr.

Findings. *Rural Bangladesh* - The prevalence of diarrhea in mothers ranged from 0.6% in Aug/Sep to 1.3% in Apr/May. The prevalence was about five times higher among children aged 6-59 mo, ranging from 5.5% in Dec/Jan to 7.6% in Apr/May. *Chittagong Division* - The prevalence of diarrhea in mothers was similar to the national rural average, but the prevalence in children was up to 4.3% higher. The prevalence peaked in Dec/Jan in both mothers and children.

Child and maternal nightblindness (SEE FIG 8)

What is indicated. Vitamin A deficiency is associated with an increased risk of illness and dying and is a leading cause of blindness among children and mothers. Nightblindness is the first clinical sign of vitamin A deficiency and is considered a public health problem in areas where the prevalence in children aged 18-59 mo is $\geq 1\%$. For each person affected by nightblindness in a population, there are many more with a low vitamin A level, which increases the risk of illness and dying.

Data collection method. The mother is asked whether she or her child have difficulty seeing and therefore moving around when there is insufficient light, for example, at dusk. Fieldworkers verify that this is due to nightblindness and not to any other sight defect.

Findings. *Rural Bangladesh* - The prevalence of nightblindness in children fell from 3.6% in 1983 to 0.6% in 1997. The data collected in 2001 show that the prevalence (0.21%) has been sustained below the level that signals a public health problem. The prevalence in non-pregnant mothers was also low (0.40%). *Chittagong Division* - The prevalence of nightblindness in both children and non-pregnant mothers was lower than all other divisions.

Child and maternal anemia (SEE FIG 9)

What is indicated. The main cause of anemia in most of Bangladesh is iron deficiency. A mother who is iron deficient or anemic during her pregnancy has a greater risk of dying during childbirth and of giving birth to a small baby with low iron stores. Iron deficiency and anemia also impair the growth and

Fig 6. Percentage of households that consumed non-cereal foods regularly (≥ 4 d in previous week) in 2001. GLV = green leafy vegetables; YOFV = yellow/orange fruits or vegetables




Fig 7. Percentage of children aged 6-59 mo and mothers with diarrhea in the previous 24 hr in 2001




Fig 8. Percentage of children aged 18-59 mo and non-pregnant mothers with nightblindness in 2001




Fig 9. Percentage of children aged 6-59 mo and non-pregnant mothers with anemia in Oct/Nov 2001




Fig 10. Percentage of wasted non-pregnant mothers (BMI < 18.5 kg/m²) in 2001





Fig 11. Percentage of wasted children aged 0-23 mo and 24-59 mo (WHZ < -2 SD) in 2001



learning ability of children, lower immunity to infectious diseases and reduce work capacity and productivity in adults. Anemia is considered a ‘severe’ public health problem if the prevalence is $\geq 40\%$ and a ‘moderate’ public health problem if the prevalence is 20-39%.

Data collection method. A national anemia survey was conducted by the NSP during routine data collection in Oct/Nov 2001. A drop of capillary blood was taken from the finger of a random sample of 1,148 children and 1,093 mothers, and the hemoglobin concentration (Hb, g/dL) was estimated using a portable hemoglobinometer (HemoCue®). Anemia is defined as Hb < 11 g/dL for children aged 6-59 mo and Hb < 12 g/dL for non-pregnant women.

Findings. Rural Bangladesh - Almost one-half of children (47%) and one-third of mothers (33%) in rural Bangladesh were anemic in Oct/Nov 2001. **Chittagong Division** - The prevalence of anemia in mothers (43%) was the highest of all divisions, whilst the prevalence in children (40%) was lowest of all divisions except Khulna Division.

Maternal wasting (SEE FIG 10)

What is indicated. Wasting among mothers threatens both their health and survival because it increases their susceptibility to life-threatening diseases and their risk of dying, especially during childbirth. Wasted mothers are more likely to give birth to small infants and to be physically weak. The prevalence of adult wasting indicates a ‘critical’ food insecurity situation in areas where it is $\geq 40\%$ and a ‘serious’ food insecurity situation in areas where it is 20-39%. **Data collection method.** Wasting in non-pregnant women is defined as a body mass index (BMI) below 18.5 kg/m², which is calculated by dividing body weight by the square of height.

Findings. Rural Bangladesh - The prevalence of maternal wasting fluctuated seasonally in rural Bangladesh, ranging from 35% in Aug/Sep to 41% in Dec/Jan. **Chittagong Division** - The prevalence of maternal wasting was lower than all other divisions throughout the year, but nevertheless indicates ‘severe’ food insecurity throughout the year.

Child wasting (SEE FIG 11)

What is indicated. Child wasting (low weight for height) results directly from an inadequate intake of food and/or from diseases. It reflects recent or current nutritional status. The prevalence of child wasting indicates a ‘critical’ problem in areas where it is $\geq 15\%$ and a ‘serious’ problem in areas where it is 10-14%.

Data collection method. Child wasting is defined as a weight for height z-score <-2 standard deviations (SD) of the median of the reference population (NCHS).

Findings. *Rural Bangladesh* - Between 9-15% of children aged 0-23 mo and 8-14% of children aged 24-59 mo were wasted. The prevalence was highest in Jun/Jul and lowest in Dec-Mar. *Chittagong Division* - The prevalence of child wasting was similar to the national rural average. Child wasting was a 'serious' problem throughout most of the year.

Child stunting (SEE FIG 12)

What is indicated. Stunting (low height for age) results from consumption of a diet of inadequate quality for a prolonged period of time. As stunting takes time to develop, it reflects past nutritional status or chronic undernutrition. A prevalence of ≥40% is considered to be 'very high' and a prevalence of 30-39% is considered 'high'.

Data collection method. Child stunting is defined as a height for age z-score <-2 SD of the median of the reference population (NCHS).

Findings. *Rural Bangladesh* - In rural Bangladesh, 40-45% of children aged 0-23 mo and 50-54% of children aged 24-59 mo were stunted. The prevalence was highest in Feb-Mar and lowest in Aug-Sep. *Chittagong Division* - The prevalence of child stunting was 'very high' in children aged 24-59 mo and 'very high' or 'high' in children aged 0-23 mo throughout the year. Compared with the national rural average, the prevalence of child stunting was on average 5% lower in children aged 0-23 mo but 2% higher in children aged 24-59 mo.

Child underweight (SEE FIG 13)

What is indicated. Underweight (low weight for age) can be the result of wasting and/or stunting. A prevalence of ≥30% is considered 'very high'.

Data collection method. Child underweight is defined as weight for age z-score <-2 SD of the median of the reference population (NCHS).

Findings. *Rural Bangladesh* - Between 41-46% of children aged 0-23 mo and 53-61% of children aged 24-59 mo were underweight. The prevalence was highest in Jun/Jul and lowest in Dec/Jan, and was 'very high' in all divisions throughout all the year. *Chittagong Division* - Compared with the national rural average, the prevalence of child underweight was on average 2% lower in both age groups. The peak prevalence was in Apr/May.

Fig 12. Percentage of stunted children aged 0-23 mo and 24-59 mo (HAZ <-2 SD) in 2001





Fig 13. Percentage of underweight children aged 0-23 mo and 24-59 mo (WAZ <-2 SD) in 2001



CONCLUSIONS

The NSP recorded substantial improvements in child nutritional status in rural Bangladesh during the 1990s², and surveillance in 2001 has shown that these encouraging trends have continued. However, the prevalence of malnutrition in children and mothers is still amongst the highest in the world. Many millions of children and mothers in rural Bangladesh continue to lack sufficient access to food, health care services and a clean environment.

The NSP findings for Chittagong Division in 2001 show that 32% of households had a energy intake <1805 kcal/person/d and were considered 'extremely' poor and a further 26% had an intake of 1805-2122 kcal/person/d and were considered 'moderately' poor. The high prevalence of wasting in mothers (28-37%) was consistent with a 'serious' food insecurity

situation throughout the year. Child wasting (6-15% in children aged 0-59 mo) was a 'serious' problem throughout most of the year. The prevalence of child stunting (35-40% in children aged 0-23 mo and 50-58% in children aged 24-59 mo) and underweight (38-46% in children aged 0-23 mo and 51-63% in children aged 24-59 mo) were also 'very high'.

Anemia was a 'serious' problem in children (40%) and mothers (43%), and suggests that the diet lacked micronutrients as well as energy. The VAC program had very high coverage (93-97%) and the prevalence of nightblindness in both children and mothers was below the 1% threshold that indicates a public health problem. However, the high prevalence of stunting and anemia suggests that households in this division had a low dietary intake of micronutrients.

All necessary resources should be mobilized at all levels to ensure that faster progress is made in reducing malnutrition in Bangladesh in the coming decade. Direct nutrition interventions are needed to assist those affected by malnutrition, including nutritional rehabilitation and direct feeding programs

for the severely malnourished; micronutrient supplementation to prevent and control anemia and vitamin A deficiency among those at highest risk, particularly young children and women of child-bearing age; and food fortification to improve the micronutrient status of the population as a whole. All food, agriculture and development policies and programs should place greater emphasis on interventions to improve household food security, including crop diversification, homestead food production, employment and income creation, credit facilities, and food assistance. Infectious diseases such as diarrhea are still a major cause of malnutrition, and so households need a sanitary environment and better access to preventive and curative health services. These interventions should be complemented with poverty alleviation strategies that support labor-intensive economic growth, sound macroeconomic management, good governance and social development, including the empowerment of women, and by global trade policies that stimulate the growth of Bangladesh's economy. Surveillance should be used to monitor the implementation of policies and programs and to assess their impact.

References

1. BBS (1998). Poverty lines and poverty measurements. In: *Household Expenditure Survey 1995-1996*, pp. 115-8. Bangladesh Bureau of Statistics, Dhaka.
2. HKI/IPHN (2001). Progress in Bangladesh towards the goals of the 1990 World Summit for Children. *Nutritional Surveillance Project Bulletin No. 3*. HKI, Dhaka.



Helen Keller
WORLDWIDE

Helen Keller International
A division of
Helen Keller Worldwide

Helen Keller International, Bangladesh

P. O. Box 6066 Gulshan, Dhaka 1212
Bangladesh
Telephone: 880-2-811 6156 /912 0028
Fax: 880-2-811 3310

Contacts:

Dora Panagides, Country Director
E-mail: dpanagides@hkidhaka.org
Dr. Harriet Torlesse, Nutrition Advisor
E-mail: htorlesse@hkidhaka.org
Quazi Imad Uddin Ibrahim, Statistician
E-mail: imad@hkidhaka.org
Nasreen Huq, Sr. Policy Advisor
E-mail: nasreenh@hkidhaka.org
Md. Ruhul Amin, Publications Officer
E-mail: amin@hkidhaka.org

Helen Keller International Asia-Pacific Regional Office

P. O. Box 4338, Jakarta Pusat, Indonesia
Telephone: 62-21-719 7861 / 719 9163
Fax: 62-21-719 8148

Contacts:

Dr. Martin W. Bloem, Regional Director
E-mail: mwbloem@compuserve.com
Dr. Regina Moench-Pfanner, Regional Coordinator
E-mail: remoench@cbn.net.id
Dr. Saskia de Pee, Regional Scientific Advisor
E-mail: sdepee@compuserve.com
Md. Aminuzzaman Talukder, Regional Agriculture Advisor
E-mail: zaman@hkinpal.wlink.com.np

© 2002 Helen Keller Worldwide

Reprints or reproductions of portions or all of this document are encouraged provided due acknowledgement is given to the publication and publisher.

This publication was made possible through support provided by the United States Agency for International Development Mission to Bangladesh under the terms of Cooperative Agreement No. 388-A-00-99-00060-00. The opinions expressed herein are those of the authors and do not necessarily reflect the views of USAID.