


Peru

Epidemiological Fact Sheet

on HIV/AIDS
and sexually
transmitted
infections




2000 Update



2 – Peru

Country Information

Population pyramid, 1999



UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the working group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the working group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

The working group and its partners have established a framework standardizing the collection of data deemed important for a thorough understanding of the current status and trends of the epidemic, as well as patterns of risk and vulnerability in the population. Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed-upon indicators was not available for many countries in 1999. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the working group would like to encourage all programme managers as well as national and international experts to communicate additional information to the working group whenever such information becomes available. The working group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Indicators	Year	Estimate	Source
Total Population (thousands)	1999	25,230	UNPOP
Population Aged 15-49 (thousands)	1999	13,267	UNPOP
Annual Population Growth	1990-1998	1.7	UNPOP
% of Population Urbanized	1998	71	UNPOP
Average Annual Growth Rate of Urban Population	1990-1998	2.0	UNPOP
GNP Per Capita (US\$)	1997	2,610	World Bank
GNP Per Capita Average Annual Growth Rate	1996-1997	5.4	World Bank
Human Development Index Rank (HDI)	1999	80	UNDP
% Population Economic Active		45.6	ILO
Unemployment Rate	1997	7.7	ILO
Total Adult Literacy Rate	1995	89	UNESCO
Adult Male Literacy Rate	1995	95	UNESCO
Adult Female Literacy Rate	1995	83	UNESCO
Male Secondary School Enrollment Ratio	1996	72.5	UNESCO
Female Secondary School Enrollment Ratio	1996	67.6	UNESCO
Crude Birth Rate (births per 1,000 pop.)	1999	24	UNPOP
Crude Death Rate (deaths per 1,000 pop.)	1999	6	UNPOP
Maternal Mortality Rate (per 100,000 live births)	1990	280	WHO
Life Expectancy at Birth	1998	68	UNPOP
Total Fertility Rate	1998	2.9	UNPOP
Infant Mortality Rate (per 1,000 live births)	1999	42	UNICEF/UNPOP

Contact address:
 UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance
 20, Avenue Appia
 CH-1211 Geneva 27
 Switzerland
 Fax: +41 22 791 4878
 email: surveillance@UNAIDS.org
<http://www.who.ch/emc/diseases/hiv>
<http://www.unaids.org>

Estimated number of people living with HIV/AIDS

In 1999 and during the first quarter of 2000, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1997 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalized epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 age range was used as the denominator in calculating adult HIV prevalence.

□ **Estimated number of adults and children living with HIV/AIDS, end of 1999**

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 1999:

Adults and children	48000		
Adults (15-49)	47000	Adult rate (%)	0.35
Women (15-49)	12000		
Children (0-15)	640		

□ **Estimated number of deaths due to AIDS**

Estimated number of adults and children who died of AIDS during 1999:

Deaths in 1999	4100
-----------------------	-------------

□ **Estimated number of orphans**

Estimated number of children who have lost their mother or both parents to AIDS (while they were under the age of 15) since the beginning of the epidemic:

Cumulative orphans	8900
---------------------------	-------------

Estimated number of children who have lost their mother or both parents to AIDS and who were alive and under age 15 at the end of 1999:

Current living orphans	6261
-------------------------------	-------------

Assessment of epidemiological situation – Peru

A study conducted in Lima, the major urban center, in 1996-97 found less than 1 percent of antenatal clinic women tested were HIV positive.

Limited information on HIV prevalence among sex workers is available since the mid-1980s. In 1986-88, less than 1 percent of sex workers tested were HIV positive. In 1989-90, nearly 1 percent of sex workers tested were HIV positive. By 1995-96, 5 percent of sex workers tested were HIV positive. Outside of the major urban areas, HIV prevalence among sex workers in Callao increased from no evidence of HIV infection in 1986 to 1 percent in 1988-89. Among sex workers tested in Ica in 1997, 2 percent were HIV positive.

HIV prevalence among IVDU increased from 1 percent in 1986-88 to 28 percent in 1989-90.

Information from all 24 geographic departments shows HIV prevalence among STD clinic patients tested increasing from 2 percent in 1986-88 to 19 percent in 1989-90. A study conducted in one site in 1995 found 7 percent of STD clinic patients tested HIV positive.

4 - Peru

HIV sentinel surveillance

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV data base maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences is compiled. To provide for a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study on which the medians were calculated are printed at the end of this fact sheet.



The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and – where applicable – other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

□ HIV prevalence in selected populations in percent (for blood donors: 1/100 000)

Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	
Pregnant women	Major Urban Areas	N-sites																1	
		Minimum																0.5	
		Median																	0.5
		Maximum																	0.5
Pregnant women	Outside Major Urban Areas	N-sites																	
		Minimum																	
		Median																	
		Maximum																	
Sex workers	Major Urban Areas	N-sites				1			1							1			
		Minimum				0.3			0.7							5			
		Median				0.3			0.7							5			
		Maximum				0.3			0.7							5			
Sex workers	Outside Major Urban Areas	N-sites			1	1	1				1		1		1	1			
		Minimum			0	0.3	0.6					0.6		0.8		0.8	1.9		
		Median			0	0.3	0.6					0.6		0.8		0.8	1.9		
		Maximum			0	0.3	0.6					0.6		0.8		0.8	1.9		
Injecting drug users	Major Urban Areas	N-sites				1			1										
		Minimum				1.2			28.1										
		Median				1.2			28.1										
		Maximum				1.2			28.1										
Injecting drug users	Outside Major Urban Areas	N-sites																	
		Minimum																	
		Median																	
		Maximum																	
STI patients	Major Urban Areas	N-sites				1			1										
		Minimum				1.7			18.7										
		Median				1.7			18.7										
		Maximum				1.7			18.7										
STI patients	Outside Major Urban Areas	N-sites												1					
		Minimum												7					
		Median												7					
		Maximum												7					
Blood Donors	National	N-sites																	
		Minimum																	
		Median																	
		Maximum																	
Blood Donors	Major Urban Areas	N-sites																	
		Minimum																	
		Median																	
		Maximum																	
Men having sex with men	Major Urban Areas	N-sites																	
		Minimum																	
		Median																	
		Maximum																	

Maps of HIV sentinel sites

Mapping the geographical distribution of HIV sentinel sites for different population groups may assist interpreting both the national coverage of the HIV surveillance system and explaining differences in levels and trends of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the UNICEF/WHO HealthMap Programme, has produced maps showing the location and HIV prevalence of HIV sentinel sites in relation to population density, major urban areas and communication routes. Maps illustrate separately the most recent results from HIV sentinel surveillance in pregnant women and in sub-populations at higher risk of HIV infection.



The boundaries and names shown and the designations used on these maps do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2000, all rights reserved.

6 - Peru

Reported AIDS cases

AIDS cases by year of reporting

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	Total	Unkn
0	0	0	0	1	4	5	22	104	177	257	372	398	646	669	789	1070	1183	1058	954	310	8940	921

Date of last report: 30-09-1999

Aids cases by age and sex

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases is aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of HAART (Highly Active Anti-Retroviral Therapy).

Sex	Age	<96	1996	1997	1998	1999	Unkn.	Total	%
All	All	4514	1183	1058	954	310	921	8019	100.0
	0-4	98	29	16	18	22	39	183	2.3
	5-9	13	5	2	3	1	7	24	0.3
	10-14	11	3	2	2	3	2	21	0.3
	15-19	149	42	32	21	12	20	256	3.2
	20-24	645	170	154	169	48	120	1186	14.8
	25-29	962	308	232	205	63	163	1770	22.1
	30-34	923	209	202	179	50	175	1563	19.5
	35-39	629	137	154	148	40	123	1108	13.8
	40-44	369	108	91	98	26	84	692	8.6
	45-49	235	60	62	43	21	61	421	5.3
	50-54	150	44	40	36	15	25	285	3.6
	55-59	94	22	27	6	2	21	151	1.9
	60+	97	18	35	16	4	23	170	2.1
NS	139	28	9	10	3	58	189	2.4	
Male	All	3865	922	861	729	234	725	6611	100.0
	0-4	53	15	9	11	14	24	102	1.5
	5-9	6	1	2	2	1	5	12	0.2
	10-14	7	2	1	1	1	1	12	0.2
	15-19	119	26	25	13	9	12	192	2.9
	20-24	534	126	121	114	30	95	925	14.0
	25-29	834	239	198	166	50	130	1487	22.5
	30-34	820	174	165	133	42	141	1334	20.2
	35-39	545	109	124	115	28	97	921	13.9
	40-44	317	91	74	77	22	65	581	8.8
	45-49	214	50	51	35	18	51	368	5.6
	50-54	127	34	31	32	12	20	236	3.6
	55-59	84	16	22	6	1	19	129	2.0
	60+	84	17	31	15	4	20	151	2.3
NS	121	22	7	9	2	45	161	2.4	
Female	All	631	249	183	217	73	184	1353	100.0
	0-4	44	13	6	6	8	14	77	5.7
	5-9	7	4	0	1	0	2	12	0.9
	10-14	4	1	1	1	2	1	9	0.7
	15-19	29	15	6	8	3	8	61	4.5
	20-24	105	42	29	54	18	24	248	18.3
	25-29	126	63	31	37	12	30	269	19.9
	30-34	100	35	36	43	8	30	222	16.4
	35-39	82	27	28	33	11	25	181	13.4
	40-44	51	17	17	21	3	18	109	8.1
	45-49	21	10	11	8	3	9	53	3.9
	50-54	23	9	9	3	3	5	47	3.5
	55-59	10	6	3	0	1	2	20	1.5
	60+	13	1	4	1	0	3	19	1.4
NS	16	6	2	1	1	13	26	1.9	
NS	All	18	12	14	8	3	12	55	100.0
	0-4	1	1	1	1	0	1	4	7.3
	5-9	0	0	0	0	0	0	0	0.0
	10-14	0	0	0	0	0	0	0	0.0
	15-19	1	1	1	0	0	0	3	5.5
	20-24	6	2	4	1	0	1	13	23.6
	25-29	2	6	3	2	1	3	14	25.5
	30-34	3	0	1	3	0	4	7	12.7
	35-39	2	1	2	0	1	1	6	10.9
	40-44	1	0	0	0	1	1	2	3.6
	45-49	0	0	0	0	0	1	0	0.0
	50-54	0	1	0	1	0	0	2	3.6
	55-59	0	0	2	0	0	0	2	3.6
	60+	0	0	0	0	0	0	0	0.0
NS	2	0	0	0	0	0	2	3.6	

AIDS cases by mode of transmission

Hetero: Heterosexual contacts.

Homo/Bi: Homosexual contacts between men.

IDU: Injecting drug use. This transmission category also includes cases in which other high-risk behaviours were reported, in addition to injection of drugs.

Blood: Blood and blood products.

Perinatal: Vertical transmission during pregnancy, birth or breastfeeding.

NS: Not specified/unknown.

Sex	Trans. Group	<96	1996	1997	1998	1999	Unkn	Total	%
All	Total	4514	1183	1058	954	310	921	8940	100.0
	Hetero	1345	621	564	553	197	407	3687	41.2
	Homo/Bi	2300	401	347	298	65	308	3719	41.6
	IDU	4	0	0	0	1	2	7	0.1
	Blood	100	10	4	1	1	16	132	1.5
	Perinatal	85	23	14	19	24	27	192	2.1
	Other Known	49	57	67	46	13	56	288	3.2
	Unknown	631	71	62	37	9	105	915	10.2
Male	Total	3865	922	861	729	234	725	7336	100.0
	Hetero	878	413	411	354	134	274	2464	33.6
	Homo/Bi	2300	401	347	298	65	308	3719	50.7
	IDU	4	0	0	0	1	1	6	0.1
	Blood	67	6	2	1	0	8	84	1.1
	Perinatal	43	8	8	11	15	16	101	1.4
	Other Known	45	50	49	37	13	44	238	3.2
	Unknown	534	47	45	28	7	77	738	10.1
Female	Total	631	249	183	217	73	184	1537	100.0
	Hetero	462	202	148	196	62	128	1198	77.9
	IDU	0	0	0	0	0	1	1	0.1
	Blood	32	4	2	0	1	8	47	3.1
	Perinatal	42	14	5	7	9	10	87	5.7
	Other Known	3	5	15	7	0	12	42	2.7
	Unknown	92	24	13	7	1	25	162	10.5
	NS	Total	18	12	14	8	3	12	67
Hetero		5	6	5	3	1	5	25	37.3
IDU		0	0	0	0	0	0	0	0.0
Blood		1	0	0	0	0	0	1	1.5
Perinatal		0	1	1	1	0	1	4	6.0
Other Known		1	2	3	2	0	0	8	11.9
Unknown		5	0	4	2	1	3	15	22.4

Curable Sexually Transmitted Infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STI are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STI facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Significant also is the observation of a sharp decline in the concentration of HIV in the genital secretions when the infection is treated. Monitoring trends in STI can provide valuable information on the sexual transmission of HIV as well as the impact of behavioural interventions, such as promotion of condom use.

Clinical services offering STI care are an important access point for people at high risk for both AIDS and STI, not only for diagnosis and treatment but also for information and education. Therefore, control and prevention of STI have been recognized as a major strategy in the prevention of HIV infection and ultimately AIDS. One of the cornerstones of STI control is adequate management of patients with symptomatic STIs. This includes diagnosis, treatment and individual health education and counselling on disease prevention and partner notification. Consequently, monitoring different components of STI control can also provide information on HIV prevention within a country.

Estimated incidence and prevalence of curable STIs

STI's	Year	Incidence			Year	Prevalence		
		Male	Female	All		Male	Female	All
Chlamydia trach.								
Gonorrhoea								
Syphilis								
Trichomonas								

Comments:

Source:

STI Incidence, men

Prevention Indicator 9: Proportion of men aged 15-49 years who reported episodes of urethritis in the last 12 months.

Year	Area	Age	Rate	N=
1996	All		9.0	2123

Comments:

Sources: DHS - 1996

STI Prevalence, women

Prevention Indicator 8: Proportion of pregnant women aged 15-24 years attending antenatal clinics whose blood has been screened with positive serology for syphilis.

Year	Area	Age	Rate	N=
1996	Lima		12.0	3566
1997	Lima		17.0	3450
1998	Lima		8.0	3990

Comments:

Sources: PROCETSS

STI Case management (counselled)

Prevention Indicator 7: Proportion of people presenting with STI or for STI care in health facilities who received basic advice on condoms and on partner notification.

Year	Area	Age	Rate	N=
1998	Lima		26.0	190

Comments:

Sources: Study for the determination of the quality of care for STD's in Peru.

STI Case management (treatments)

Prevention Indicator 6: Proportion of people presenting with STI in health facilities assessed and treated in an appropriate way (according to national standards).

Year	Area	Age	Rate	N=
1998	All		71.0	184

Comments:

Sources: Study for the determination of the quality of care for STD's in Peru

8 – Peru

Health service indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. These efforts may range from reaching out to vulnerable communities through large-scale educational campaigns or interpersonal communication; provision of treatment for STIs; distribution of condoms and needles; creating and enabling environment to reduce risky behaviour; providing access to voluntary testing and counselling; home or institutional care for persons with symptomatic HIV infection; and preventing perinatal transmission and transmission through infected needles or blood in health care settings. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators – such as the percentage of a population with access to health care services; the percentage of women covered by antenatal care; or the percentage of immunized children – may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS – related issues.

Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services – total:			
% of population with access to health services – urban:			
% of population with access to health services – rural:			
Contraceptive prevalence rate (%):	1990-1999	64	UNICEF/UNPOP
% of births attended by trained health personnel:	1990-1999	56	UNICEF
% of 1-yr-old children fully immunized – DPT:	1995-1998	99	UNICEF
% of 1-yr-old children fully immunized – Polio:	1995-1998	99	UNICEF
% of 1-yr-old children fully immunized – Measles:	1995-1998	90	UNICEF
Proportion of blood donations tested:			
% of ANC clinics where HIV testing is available:			
HIV/AIDS Hospital Occupancy Rate (Days):			

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programmes implement activities to increase both availability of and access to condoms. The two condom availability indicators below are intended to highlight areas of strength and weakness at the beginning and end of the distribution system so that programmatic resources can be directed appropriately to problem areas.

Condom availability (central level)

Prevention Indicator 2: Availability of condoms in the country over the last 12 months (central level).

Year	Area	N	Rate

Comments:
Sources:

Condom availability (peripheral level)

Prevention Indicator 3: Proportion of people who can acquire a condom (peripheral level).

Year	Area	N	Rate

Comments:
Sources:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, intravenous drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of regular behavioural surveys in order to monitor trends in behaviours and target interventions.

Knowledge of HIV- related preventive practices

Prevention Indicator 1: Proportion of people citing at least two acceptable ways of protection from HIV infection.

Year	Area	Age Group	Male	Female	All
1996	All	15-49	37.6	32.9	

Comments:

Sources: DHS, 1996

Reported non-regular sexual partnerships

Prevention Indicator 4: Proportion of sexually active people having at least one sex partner other than a regular partner in the last 12 months.

Year	Area	Age Group	Male	Female	All
1996	All	15-49	13.6	1.5	

Comments:

Sources: DHS, 1996

Reported condom use in risk sex (gen pop)

Prevention Indicator 5: Proportion of people reporting the use of a condom during the most recent intercourse of risk.

Year	Area	Age Group	Male	Female	All
1996	All	15-49	32.6	35.3	

Comments:

Sources: DHS, 1996

10 – Peru

Knowledge and behaviour

Ever use of condom

Percentage of people who ever used a condom.

Year	Area	Age Group	Male	Female	All
1992	All	15-19		1.8	
1996	All	15-19		3.6	
1992	All	20-24		7.5	
1996	All	20-24		15.0	
1992	All	25-29		14.5	
1996	All	25-29		22.3	
1992	All	30-34		20.2	
1996	All	30-34		24.4	
1992	All	35-39		17.3	
1996	All	35-39		23.2	
1992	All	40-44		14.5	
1996	All	40-44		17.9	
1992	All	45-49		13.2	
1996	All	45-49		12.1	
1992	All	Total		11.3	
1996	All	Total		16.2	

Comments:

Sources: Demographic and Health Survey

Median age at first sexual experience

Median age of people at which they first had sexual intercourse.

Year	Area	Age Group	Male	Female	All
1996	All	20-24		19.7	
1996	All	45-49		18.8	

Comments:

Sources: DHS/1996

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

Year	Area	Age Group	Rate	N
1992	All	15	693	2.2
1996	All	15	1308	2.6
1992	All	16	761	5.2
1996	All	16	1294	7.8
1992	All	17	712	9.4
1996	All	17	1228	11.6
1992	All	18	660	17.5
1996	All	18	1218	17.9
1992	All	19	651	24.7
1996	All	19	1090	29.8

Comments:

Sources: DHS/1992 – DHS/1996

Sources

Data presented in this Epidemiological Fact Sheet come from several different sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations Agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

-
- Aguero, G., F. S. Wignall, W. Alexander, et al., 1988, HIV Infections in Peru, IV International Conference on AIDS, Stockholm, 6/13-14, Poster 5078.
- Alarcon, J., O. Palacios, A. Tejada, et al., 1989, Incidence d'Infection HIV-1 dans Prostituees de Callao, Perou. Etude de Cohorte., V International Conference on AIDS, Montreal, 6/4-9, Poster Th.H.P. 17.
- Alarcon, J., C. Rodriguez, K. Johnson, et al., 1997, Determinants and Prevalence of HIV and Syphilis in Pregnant Women in Lima, Peru, V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract P033.
- Golenbock, D. T., J. Guerra, J. Pfister, et al., 1988, Absence of Infection with Human Immunodeficiency Virus in Peruvian Prostitutes, AIDS Research and Human Retroviruses, vol. 4, no. 6, pp. 493-499.
- Getaneh, A., J. Sanchez, D. Watts, et al., 1997, Sero Prevalence of Sexually Transmitted Disease and Infections Related to Sexual Behavioral Characteristics among the . . ., V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCV264.
- McCarthy, M. C., F. S. Wignall, J. Sanchez, et al., 1996, The Epidemiology of HIV-1 Infection in Peru, 1986-1990, AIDS vol. 10, no. 10, pp. 1141-1145.
- Quijano, E., M. Cairampoma, A. Ramirez, et al., 1997, Impacto de las Intervenciones de Salud Sobre la Infeccion por VIH en Trabajadoras Sexuales Callao, Peru 1987 - Junio 1997, V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCJ176.
- Revata, C., P. Quispe, E. Nino de Guzman, et al., 1997, Seroprevalencia de Infeccion por VIH en 2 Grupos de Frecuente Transmision en la Provincia de ICA, Peru 1995 - 1996, V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCV256.
-

Websites:

National AIDS Programme (Spanish only):

<http://www.minsa.gob.pe/procetss/index.htm>

12 – Peru

Annex: HIV Surveillance data by site

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Pregnant women	Major Urban Areas	Lima															0.5
----------------	-------------------	------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	-----

Pregnant women	Outside Major Urban Areas																
----------------	---------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Sex workers	Major Urban Areas	National			0.3			0.7									5
-------------	-------------------	----------	--	--	-----	--	--	-----	--	--	--	--	--	--	--	--	---

Sex workers	Outside Major Urban Areas	Callao		0	0.3	0.6					0.6	0.8		0.8			1.9
		Not specified															
		Ica															

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Injecting drug users	Major Urban Areas	National			1.2			28.1									
----------------------	-------------------	----------	--	--	-----	--	--	------	--	--	--	--	--	--	--	--	--

Injecting drug users	Outside Major Urban Areas																
----------------------	---------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

STI patients	Major Urban Areas	National			1.7			18.7									
--------------	-------------------	----------	--	--	-----	--	--	------	--	--	--	--	--	--	--	--	--

STI Patients	Outside Major Urban Areas	Not specified												7			
--------------	---------------------------	---------------	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Blood Donors	National																
--------------	----------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Blood Donors	Major Urban Areas																
--------------	-------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Group	Area	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
-------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Men having sex with men	National																
-------------------------	----------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--