



**IOCH**  
**Immunization and Other Child Health Project**

**Vaccination Coverage Survey of**  
**Naogaon Municipality**

**February 2001**

**Survey Report No. 32**

**This survey was conducted by IOCH, a project of Management Sciences for Health,  
funded by USAID under AID contract No. HRN-I-01-98-00033-00, Task Order No. 01**

House 1, Road 23, Gulshan 1, Dhaka 1212, Bangladesh  
Tel: 8828596, 8829279, 8813611, 8813410  
Fax: 880-2-8826229  
E-mail: [ioch@citechco.net](mailto:ioch@citechco.net)

July 2001

## Table of Contents

List of tables and charts	2
Acronyms	3
Terminology	4
<b>Summary Results</b>	<b>5</b>
Background	5
Objectives	5
Methodology	5
Routine immunization coverage levels of children	6
Routine TT immunization coverage levels of women	6
Coverage levels in the 8 <sup>th</sup> NID campaign	7
Suggested solutions	7
Map showing the locations of the selected clusters	8
Tables and Charts	9
<b>Annexures</b>	
Annex A: EPI Cluster Survey design (extracts from an article written by Anthony G Turner, Robert J Magnani and Muhammed Shuaib)	15
Annex B: List of selected clusters for the survey	16
Annex C: List of never vaccinated children by clusters	17
Acknowledgements	18
List of IOCH reports	19

## **List of Tables**

**Table 1:** Routine immunization coverage levels of the children

**Table 2:** Invalid doses of immunization provided to the children

**Table 3:** Missed opportunities by antigens

**Table 4:** EPI Card availability and retention

**Table 5:** Distance between the child's home and the vaccination site

**Table 6:** Reasons for non-immunization and partial immunization of the children

**Table 7:** Reasons for non-immunization and partial immunization for TT of the women

**Table 8:** Coverage of the 1<sup>st</sup> Round of the 8<sup>th</sup> NID Campaign

**Table 9:** Reasons for non-immunization of OPV during the 1<sup>st</sup> Round of the 8<sup>th</sup> NID campaign

## **List of Charts**

**Chart 1:** Drop-out rate for childhood immunization

**Chart 2:** Sources of childhood immunization services

**Chart 3:** Respondent's knowledge about required visits to immunization Centers

**Chart 4:** Routine immunization coverage levels for TT of the women (15-49 years)

**Chart 5:** Drop-out rates for TT immunization

**Chart 6:** Providers of TT immunization

**Chart 7:** Sources of information about the eighth NID campaign

## Acronyms

BCC	Behavior Change Communication
BCG	Bacillus of Calmette and Guerin
CES	Coverage Evaluation Survey
COSAS	Coverage Survey Analysis System
DPT	Diphtheria, Pertussis and Tetanus
EPI	Expanded Program on Immunization
FWC	Family Welfare Center
IOCH	Immunization and Other Child Health
Mahallah	Smaller localities (smaller than a village)
MOHFW	Ministry of Health and Family Welfare
Mouza	Smallest administrative locality in an Upazila
MSH	Management Sciences for Health
NGO	Non Governmental Organization
NID	National Immunization Day
OPV	Oral Polio Vaccine
TT	Tetanus Toxoid
WHO	World Health Organization

## Terminology

This provides the meaning of some of the more technical terms used in this report and a brief explanation of their use.

**By card:** An immunization given to a child is termed as by card if the date of the dose is entered on an immunization card. Only doses recorded by card are treated as valid data in this survey.

**By history:** Immunization history collected from a parent's recall is termed as by history. Often no date will be mentioned. This information is only included in crude data.

**Crude coverage** rate is calculated from the doses recorded by card and/or by history. It is not ascertained whether the doses were given at the correct age and/or following the correct interval (where applicable). Crude data however, helps us to understand how much additional coverage could be achieved if all vaccines were given at the optimum age for the child and following the optimum interval. It also provides useful information on access to the EPI program and on the operational aspects of the provision of health services.

**Valid coverage** rate is calculated from the vaccinations recorded by card. Valid data includes only the doses of vaccines that were given after the minimum date of eligibility and/or after the minimum interval necessary to be effective and to protect the child. There is no maximum interval for a dose and therefore a dose administered after 52 weeks is still regarded as valid. By comparing crude coverage with valid coverage data of any particular antigen, one can determine how much coverage was lost due to the inability to give vaccine at the appropriate time.

**Invalid doses** are those administered at the wrong age and/or at the wrong interval. Doses administered before the minimum age in the case of DPT/Polio 1<sup>st</sup> doses and Measles vaccine or with less than four weeks interval in the case of DPT or Polio vaccines are classified as "invalid" doses.

The **criteria for a valid dose** used in this survey is the criteria recognized by the Bangladesh EPI program: minimum age for DPT/Polio 1<sup>st</sup> dose - 6 weeks old; minimum DPT/Polio interval - 4 weeks; minimum age for Measles vaccine - 38 weeks old.

**Program access** is measured by the percentage of children surveyed who received DPT 1<sup>st</sup> dose (crude data – by card and history) in the routine immunization session.

**Fully immunized** means the child has received all the doses it requires (BCG, OPV 1-3, DPT 1-3 and measles).

**Missed opportunity** refers to a visit of a child to a vaccination center for a dose that he received. However at that time he was also eligible for another dose of antigen that he did not receive. If the missed dose was provided at a later date, it is a *corrected missed opportunity*. If not, it is an *uncorrected missed opportunity*.

## SUMMARY RESULTS

### Background

The routine EPI program in the municipalities is carried out by a variety of private and public providers at fixed (hospitals, clinics, dispensaries, etc.) and at outreach sites. NGOs and private practitioners also provide immunization services in many places. The municipal authorities are primarily responsible for providing and/or coordinating primary health care including routine EPI services in municipal areas. However, in the absence of an effective management information system and reliable service statistics at municipal level, it is often difficult to assess the level of immunization coverage of the municipalities. In view of this situation, the IOCH decided to conduct a series of coverage evaluation surveys in the selected municipal areas to assess the level of immunization coverage in these municipal areas. As a part of this effort, the IOCH conducted a coverage evaluation survey in Naogaon Municipality in February 2001.

### Objectives

The overall objective of the survey was to assess the level of immunization coverage in Naogaon Municipality. The specific objectives were to:

- a) assess the level of routine immunization coverage of the children (12-23 months) and find out the reasons for non-immunization and partial immunization.
- b) assess the level of TT immunization coverage among women of 15-49 years of age, irrespective of their marital status, and find out the reasons for non-immunization and partial immunization.
- c) assess the coverage levels of OPV and vitamin A administered during the 8<sup>th</sup> NID campaign.

### Methodology

The survey employed the WHO recommended 30-cluster survey methodology that has been widely used in many developing countries to assess immunization coverage (the survey methodology and its limitations are presented in Annex A). In all, 30 clusters were randomly selected from Naogaon Municipality following PPS sampling procedures. A list of the selected clusters is provided in Annex- B and their locations are shown on the maps in page 8. From each cluster, 7 children 12 – 23 months and 7 women of reproductive age (15 – 49 years) irrespective of their marital status were selected following 30 cluster survey methodology to ascertain their routine immunization coverage. Also, 7 children < 5 years (0 – 59 months) were selected to assess the immunization coverage of the 8<sup>th</sup> National Immunization Campaign.

The WHO standard questionnaires were used for documenting the routine immunization status of the children and women. Also, separate questionnaires were used to collect data on NIDs and reasons for non-immunization and dropouts. The data were collected by the experienced Field Investigators of the Survey Team of the IOCH. Data processing and analysis were done by the Monitoring & Evaluation Unit of the IOCH using COSAS 4.41<sup>1</sup> and EpiInfo.

---

<sup>1</sup> COSAS (Coverage Survey Analysis System) is a dedicated software for analyzing coverage evaluation survey data.

### **Coverage levels for the routine immunization of children**

*Access:* Based on crude data (card plus history), 99% children received at least one dose of antigen (DPT 1<sup>st</sup> dose in this case) from routine immunization sessions. Only 1% children did not receive a dose of any antigen.

*Crude coverage between 12-23 months:* 99% children received BCG, 97% received three doses of OPV, 97% received three doses of DPT and 94% received measles vaccine. 93% children were fully immunized.

*Valid coverage between 12-23 months:* 99% children received BCG, 80% received three doses of OPV, 80% received three doses of DPT and 93% received measles vaccine. 76% children were fully immunized.

*Valid coverage by 12 months:* 99% children received BCG, 80% received three doses of OPV, 80% received three doses of DPT and 92% received measles vaccine. 76% children were fully immunized.

*Source of immunization and distance of immunization center:* Childhood immunization in this area was provided mostly by the GOB EPI outreach centers (78%), followed by hospitals and clinics.

*Reasons for non-immunization and partial immunization of children:* In this area, only two children were found non-immunized. One of them never received a dose of any antigen because of sickness. The reason for non-immunization of the other child was not specified. However, 12 children dropped out, and majority of them dropped out because of illness. The other reasons for dropout included: a) the parents did not know that their children needed to receive the 2<sup>nd</sup>/3<sup>rd</sup> doses to get full immunization (17%); b) wrong knowledge of adverse reaction to immunization (17%); and c) parents were too busy to take their children to EPI centers (17%).

*Problems detected:* Although access to immunization was quite high, there was 2% dropout from DPT1 to DPT3 and 5% from DPT1 to Measles vaccine. There were also a number of invalid doses due to early immunization (8% for DPT1, 7% for DPT2 and 5% for DPT3 vaccine). However, prevalence of uncorrected missed opportunity was very low (ranging 0 for DPT3 to 1% for measles). Child immunization cards (EPI Cards) were ever given to 99% of the children; but they were available at the time of interview in 66% of the cases. EPI Card retention rate was 67%, which means that 33% of the EPI cards were lost. Over one-fourth of the parents (29%) could mention correctly the number of times (i.e. 4 times) their children needed to visit the immunization centers to get fully immunized.

### **Coverage levels for the routine TT immunization of women**

The survey of coverage levels for immunization against tetanus toxoid showed that about 74% of women of reproductive age (15-49 years) received a first dose of TT; 71% women received two doses; 50% received 3 doses, and 22% received 5 doses of the TT vaccine. About one-fourth of the women never received any TT vaccine.

*Dropout rate for TT immunization:* The dropout rate for TT immunization among the women of reproductive age was quite high. The dropout rate for TT1 to TT2 was 3% and for TT1 to TT 3 was 20%.

*Reasons for non-immunization and partial immunization of women:* One of the major reasons cited for non-immunization was that the women were unaware of the need for immunization (33%). 46% women mentioned that there was no provision of TT vaccine when they had their pregnancies, while 17% were afraid of the pain of TT injection. The major reasons for partial immunization or dropout were that the women were either unaware of the need to return for subsequent doses of TT immunization (48%) or the health worker did not specify when to return for the next dose of TT vaccine (12%).

### **Coverage levels for the 8<sup>th</sup> NID campaign**


During the 8<sup>th</sup> NIDs, 99% of the children <5 years received OPV in both the rounds. The coverage for OPV in either round was also 99%. Vitamin A capsules were given to 96% of the eligible children during the 8<sup>th</sup> NID campaign.

Majority of the parents learned about the NID campaign from IPC by the GOB health workers (83%) and miking (44%). The other sources of information included family members and neighbors (14%) and television/radio (21%).

### **Suggested solutions**

The survey indicates the need for appropriate BCC activities and counseling to make the parents/caretakers understand the importance of full immunization of their children. The information must stress that each child, irrespective of sex, must be fully immunized before 12 months of age, and how to achieve full immunization (the time and place of the immunization sessions and the number of doses required). All women of childbearing age must be informed about how to prevent tetanus with 5 doses of TT vaccination to protect herself and her newborn child.

The poor quality of services, as reflected by the percentage of invalid doses (8% for DPT1, 7% for DPT2 and 5% for DPT3) and dropout rates (2% for DPT1 to DPT3 and 5% for DPT1 to measles) could be further improved by proper screening of eligibility of clients for each antigen at the time of vaccination, and by adequate counseling of mothers. There is also a need to further train the service providers to help them keep up with EPI policies and guidelines and increase their capacity for counseling parents and women of reproductive age about EPI.




## TABLES AND CHARTS

**Table 1: Routine immunization coverage levels of the children**

Name of the vaccine	Coverage (%) Immunization of 12-23 months age group		Coverage (%) Immunized by 12 months of age
	Crude data (Access)	Valid data	Valid data
BCG	99	99	99
Polio-1	99	91	91
Polio-2	98	94	84
Polio-3	97	80	80
DPT-1	99	91	91
DPT-2	98	84	84
DPT-3	97	80	80
Measles	94	93	92
Fully immunized	93	76	76
Zero Dose	1	-	-

**Chart-1: Drop-out rate for childhood immunization**



**Table 2: Invalid doses of immunization provided to the children**

Invalid doses	Percentage
DPT-1	8
DPT-2	7
DPT-3	5
Measles	1


**Table 3: Missed opportunities by antigens**

Name of the vaccine	Uncorrected		Corrected	
	Number	Percent	Number	Percent
BCG	1	1	0	-
DPT-1	1	1	9	4
DTP-2	1	1	8	4
DPT-3	0	-	9	4
OPV-1	1	1	9	4
OPV-2	1	1	8	3
OPV-3	0	-	9	4
Measles	1	1	2	1

**Table 4: EPI card availability and retention**

Card Status	Number	Percentage
EPI card available	139	66
EPI card ever given	208	99
EPI card retention	139	67

**Chart 2: Sources of childhood immunization services**




**Table 5: Distance between the child's home and the vaccination site**

Time (Minutes)	Number	Percentage
1-5 Min.	132	63
6-10 Min.	60	29
11-30 Min.	18	9
Don't know	-	-


**Table 6: Reasons for non-immunization and partial immunization of the children**

Reasons	Non- immunized (%) (N=2)	Partially immunized (%) (N=12)
Did not know that the child needs to receive the 2 <sup>nd</sup> /3 <sup>rd</sup> doses	-	17
Wrong knowledge about adverse reaction	-	17
Mother too busy	-	17
Child sick, was not taken to site	50	33
Mother was sick/family problem	-	8
Others	50	8


**Chart 3: Respondents' knowledge about required visits to immunization centers**




**Chart 4: Routine immunization coverage levels for TT of women (15-49 years)**



**Chart 5: TT Immunization drop-out rate among women 15-49 years**



**Chart 6: Providers of TT immunization**



**Table 7: Reasons for non-immunization and partial immunization for TT of the women**

<b>Reasons</b>	<b>Non- immunized (%) (N=24)</b>	<b>Partially immunized (%) (N=116)</b>
Next dose is not yet due	-	19
Don't feel need for immunization	33	-
Health worker did not specify the next dose	-	12
According to health worker 2/3 doses of TT is enough during pregnancy	-	12
Unaware of need to return for subsequent doses	-	48
In our times TT immunization was not in practice	46	-
Postponed until another time	-	2
Fear of side reactions	4	-
Fear of injection	17	3
Others		4


**Table 8: Coverage of the 8<sup>th</sup> NID Campaign**

<b>Round</b>	<b>OPV %</b>	<b>Vit "A" %</b>
1 <sup>st</sup> round	99	96
2 <sup>nd</sup> round	99	-
Both rounds	99	-
Any round	100	-

**Table 9: Reasons for non-immunization of OPV during 8<sup>th</sup> NID campaign**

<b>Reasons</b>	<b>1<sup>st</sup> Round %</b>	<b>2<sup>nd</sup> Round %</b>
Child way from home	100	-
Others	-	100

**Chart 7: Source of information about the 8<sup>th</sup> NID campaign**



The following are extracts from **Anthony G Turner, Robert J Magnani and Muhammad Shuaib's** article entitled **"A not quick as quick but much cleaner alternative to the Expanded Programme on Immunization (EPI) cluster survey design"** published in the *International Journal of Epidemiology* in 1996, volume 25, Issue No. 1, pages 198-203.

#### The standard EPI Cluster Survey Design

"The sample design for the EPI Cluster Survey is a two-stage design involving the selection of 30 primary sampling units or 'clusters' (usually village or other area units), from which 210 children with a target age range (usually 12-23 months) are chosen, seven children per cluster. The sample size of 210 children (per domain or stratum) is mandated by the desire to estimate the level of immunization coverage to within +/- 10 percentage points of the true population proportion with 95% statistical confidence, assuming a design effect (i.e. *deff*) of 2.0. Based upon prior experience with immunization coverage surveys (primarily in the US), 30 clusters are generally thought to be necessary to yield sufficiently reliable estimate."

"In the standard design, clusters are chosen from a list of primary sampling units (i.e. villages, urban communities, census enumeration areas etc.) through systematic random sampling with probability proportional to estimated size (*ppes*). The latest estimates of cluster population sizes, which are assumed to be proportional to the number of children in the target age group in each cluster, are typically used as measures of size. The 30 clusters so chosen are then visited by survey field staff who carry out the second stage of sample selection and conduct the household interviews. "

"The original EPI design called for sample children to be chosen randomly from a list of all eligible children in each sample cluster. However, because the creation of lists of households and children tends to be time-consuming, costly, and unfeasible in some settings, this procedure is only infrequently used in actual practice. Instead, one of several simplified second stage sampling procedures is commonly used. In one variant, children are selected by first choosing a random direction from a central location in a village or community (e.g. by spinning a bottle). The number of households in that direction to the edge of the community is then counted, and one household is randomly chosen to be the first sample household. Subsequent households are chosen by visiting the nearest neighboring households until information has been gathered on seven children. In a yet simpler variant, a direction from a central starting point is randomly chosen as described above and households are contacted as the interviewer moves in the chosen direction until the required information has been gathered for seven children."

"The second stage sampling methods described above are 'quota sampling procedures' and some of the problems resulting from the use of this approach have been noted over the years."

"First, quota sampling does not ensure that every eligible member of the target population has a known, non-zero chance of being selected. Hence, the standard EPI design, as it is usually applied, is not a true probability sample design. ...."

"A second problem concern sampling weights. .... However, since measures of size in sampling frames are often inaccurate due to census errors and changes in population since the census was taken, application of the standard EPI Cluster Survey method does not automatically result in a self weighting sample. The survey data must be weighed in order to yield unbiased estimates. .... However, since selection probabilities are not known in most EPI Cluster Survey applications, sampling weights can not be calculated."

"Thirdly, a computer simulation study demonstrates that the EPI Cluster Survey based upon quota sampling at the second stage of sample selection is considerably more prone to sampling bias than conventional cluster sampling, particularly where immunized children are 'pocketed' within clusters. "

"Finally, there is the issue of how second stage sample selection should proceed in surveys with multiple measurement objectives."

## List of selected cluster for the survey

<b>Mahallah name</b>	<b>Total Population</b>	<b>Cluster Number</b>
Baghbari	539	1
Chak Dev Para	4395	2
Chak Elam	922	3
Chak Panchi	794	4
Chak Muktar	2449	5
Daptari Para	1087	6
Kumaigari	2029	7
Mirpur	444	8
Refugee Colony (Bihari)	1700	9
Uttar Para Baganbaria	2179	10
Ukil Para	2120	11
Chak Prasad	4216	12
Chak Pran	1236	13
Dakshin Math Naogaon	4690	14
Dakshin Arzi Naogaon	4842	15,16
Khash Naogaon Mugree	1852	17
Sherpur	2155	18
Uttar Path Naogaon	5337	19
Uttar Arzi Naogaon	3584	20
Boalia (Part)	3177	21
Bhabanipur	2067	22
Jagat Singapur	1080	23
Khalishakuri	1462	24
Par Naogaon	3278	25
Pirozpur	3425	26
Purba Par Naogaonm	3604	27
Razzakpur	2791	28
Sultanpur	3927	29
Sultanpur Master Para	975	30

## List of never vaccinated children by cluster

Mahallah name	Total Population	Cluster Number	Never Vaccinated Children
Baghbari	539	1	-
Chak Dev Para	4395	2	-
Chak Elam	922	3	-
Chak Panchi	794	4	-
Chak Muktar	2449	5	1
Daptari Para	1087	6	-
Kumaigari	2029	7	-
Mirpur	444	8	-
Refugee Colony (Bihari)	1700	9	-
Uttar Para Baganbaria	2179	10	1
Ukil Para	2120	11	-
Chak Prasad	4216	12	-
Chak Pran	1236	13	-
Dakshin Math Naogaon	4690	14	-
Dakshin Arzi Naogaon	4842	15,16	-
Khash Naogaon Mugree	1852	17	-
Sherpur	2155	18	-
Uttar Path Naogaon	5337	19	-
Uttar Arzi Naogaon	3584	20	-
Boalia (Part)	3177	21	-
Bhabanipur	2067	22	-
Jagat Singapur	1080	23	-
Khalishakuri	1462	24	-
Par Naogaon	3278	25	-
Pirozpur	3425	26	-
Purba Par Naogaonm	3604	27	-
Razzakpur	2791	28	-
Sultanpur	3927	29	-
Sultanpur Master Para	975	30	-
<b>Total</b>			<b>2</b>

## **Acknowledgements**

**Survey coordination:**

Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

**Survey management:**

Mr. Jagadindra Majumder, Field Survey Manager, IOCH/MSH

**Data analysis:**

Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

Mr. Biplob Banerjee, Surveillance Data Manager, IOCH/MSH

**Report preparation:**

Md. Mafizur Rahman, Monitoring & Evaluation Specialist, IOCH/MSH

**Report review:**

Dr. Pierre Claquin, Chief of Party, IOCH/MSH

**Digital map preparation:**

Mr. Din Mohammed, Monitoring and Evaluation Assistant, IOCH/MSH

**Cover photo:**

Moumina Dorgabekova/Image Jinn

**Survey team members, IOCH/MSH:**

Md. Abdul Hamid, Field Investigator

Md. Saiful Islam, Field Investigator

Ms. Krishna Rani Shil, Field Investigator

Ms. Khaleda Akhter, Field Investigator

Ms. Mahamuda Parveen, Field Investigator

Ms. Aung Ma Ching Marma, Field Investigator

## List of IOCH Survey/Research/Technical Reports

### Survey Reports

1. Vaccination Coverage Survey of the Slums of Rajshahi City Corporation- January 2000. Survey Report No. 1. May 2000
2. Vaccination Coverage Survey of the Selected Unions along the North-western Border of Bangladesh- February 2000. Survey Report No. 2. June 2000
3. Vaccination Coverage Survey of the Selected Unions along the South-west Border of Bangladesh- February 2000. Survey Report No. 3. July 2000
4. Vaccination Coverage Survey of the Slums of Khulna City Corporation- January 2000. Survey Report No. 4. July 2000
5. Vaccination Coverage Survey of the Slums of Chittagong City Corporation- January 2000. Ward Number 1 to 18. Survey Report No. 5. July 2000
6. Vaccination Coverage Survey of the Slums of Chittagong City Corporation- January 2000. Ward Number 19 to 41. Survey Report No. 6. July 2000
7. Vaccination Coverage Survey of the Dinajpur Municipality- January 2000. Survey Report No. 7. July 2000
8. Vaccination Coverage Survey of the Noakhali Municipality- January 2000. Survey Report No. 8. July 2000
9. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 1, 2 & 4. Survey Report No. 9. July 2000
10. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 5, 6 & 7. Survey Report No. 10. July 2000
11. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 3, 8, 9 & 10. Survey Report No. 11. July 2000
12. Vaccination Coverage Survey of the Tribal and Non-tribal Populations in the North-east Border Areas of Bangladesh. Survey Report No. 12. August 2000
13. Vaccination Coverage Survey of the Sylhet Municipality – January 2000. Survey Report No. 13. August 2000.
14. Vaccination Coverage Survey of the Kishoreganj Municipality – April 2000. Survey Report No. 14. September 2000.
15. Vaccination Coverage Survey of the Rangpur Municipality – May 2000. Survey Report No. 15. September 2000.
16. Vaccination Coverage Survey of the Greater Faridpur Municipalities – June 2000. Survey Report No. 16 September 2000.
17. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Chittagong, Khulna and Rajshahi City Corporation Slums – September 2000. Survey Report No. 17. November 2000.
18. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Dhaka City Corporation Slums – September 2000. Survey Report No. 18. November 2000
19. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Slums of selected 27 Municipalities– September 2000. Survey Report No. 19. November 2000
20. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Hard-to-reach and High-risk rural areas – September 2000. Survey Report No. 20. November 2000
21. Vaccination Coverage and Other Health Care Practices Survey in the Pabna Char Areas—August 2000. Survey Report No. 21. January 2001.
22. Vaccination Coverage Survey of the Mymensingh Municipality – April 2000. Survey Report No. 22. February 2001.
23. Vaccination Coverage Survey of the Jessore Municipality – October 20-30, 2000. Survey Report No. 23. February 2001.
24. Vaccination Coverage Survey of the Comilla Municipality - October 1-9, 2000. Survey Report No. 24. February 2001
25. Vaccination Coverage Survey of the Pabna Municipality - July 2000. Survey Report No. 25. February 2001
26. Vaccination Coverage Survey of the Sirajganj Municipality - July 2000. Survey Report No. 26. February 2001
27. Vaccination Coverage Survey of the Bogra Municipality - February 2001. Survey Report No. 27. April 2001
28. Vaccination Coverage Survey of the Peri-urban Areas of DCC- Nov. 2000 (Peri-urban Area Survey-1). Survey Report No. 28. July 2001
29. Vaccination Coverage Survey of the Peri-urban Areas of DCC- Nov. 2000 (Peri-urban Area Survey-2). Survey Report No. 29. July 2001
30. Vaccination Coverage Survey of the Patuakhali, Jhalokathi and Pirojpur Municipalities- Aug. 2000. Survey Report No. 29. July 2001

### **Unicef & IOCH Survey Reports**

1. Vaccination Coverage Survey of the Teknaf and Ukhaia Upazilas- February 2000. Survey Report No. 01, August 2000
2. Vaccination Coverage Survey of the Brahmanbaria Sadar Upazila- February 2000. Survey Report No. 02, August 2000
3. Vaccination Coverage Survey of the Debidwar Upazila- February 2000. Survey Report No. 03, August 2000
4. Vaccination Coverage Survey of the Madaripur Upazila- February 2000. Survey Report No. 04, August 2000
5. Vaccination Coverage Survey of the Maulvi Bazar District- February 2000. Survey Report No. 05, August 2000
6. Vaccination Coverage Survey of the Raumari Upazila - February 2000. Survey Report No. 06, August 2000
7. Vaccination Coverage Survey of the Gangachara Upazila - February 2000. Survey Report No. 07, August 2000
8. Vaccination Coverage Survey of Chittagong Hill Tracts - February 2000. Survey Report No. 08, October 2000

### **Technical Report**

1. Joint National/International Review of EPI Program in Urban Areas of Bangladesh—23 January – 3 February 2000. Technical Report No. 01, July 2000

---

*Additional copies of any of these reports, if needed, will be provided free of cost on request to:*

Mamunul Haque, Communications Advisor, IOCH. E-mail: [mh@citechco.net](mailto:mh@citechco.net)