



IOCH
Immunization and Other Child Health Project

**Vaccination Coverage Survey of the
Kushtia Municipality**

October 2001

Survey Report No. 49

**This survey was conducted by IOCH, a project of Management Sciences for Health,
funded by USAID under AID contract No. HRN-I-01-98-00033-00, Task Order No. 01**

House 1, Road 23, Gulshan 1, Dhaka 1212, Bangladesh
Tel: 8828596, 8829279, 8813611, 8813410
Fax: 880-2-8826229
E-mail: ioch@citechco.net

December 2001

Table of Contents

List of tables and charts	2
Acronyms	3
Terminology	4
Summary Results	5
Background	5
Objectives	5
Methodology	5
Routine immunization coverage levels of children	6
Routine TT immunization coverage levels of women	6
Coverage levels of the 9 th NID campaign	7
Suggested solutions	7
Map showing the locations of the selected clusters	8
Tables and Charts	9
Annexures	
Annex A: EPI Cluster Survey design (extracts from an article written by Anthony G Turner, Robert J Magnani and Muhammed Shuaib)	15
Annex B: List of selected clusters for the survey	16
Annex C: List of never vaccinated children by clusters	17
Acknowledgements	18
List of IOCH reports	19

List of Tables

Table 1: Routine immunization coverage levels of the children

Table 2: Invalid doses of immunization provided to the children

Table 3: Missed opportunities by antigens

Table 4: EPI Card availability and retention

Table 5: Distance between the child's home and the vaccination site

Table 6: Reasons for non-immunization and partial immunization of the children

Table 7: Reasons for non-immunization and partial immunization for TT of the women

Table 8: Coverage of the 9th NID Campaign

Table 9: Reasons for non-immunization of OPV during the 9th NID campaign

List of Charts

Chart 1: Drop-out rate for childhood immunization

Chart 2: Sources of childhood immunization services

Chart 3: Respondent's knowledge about required visits to immunization centers

Chart 4: Routine immunization coverage levels for TT of the women (15-49 years)

Chart 5: Drop-out rate for TT immunization

Chart 6: Providers of TT immunization

Chart 7: Sources of information about the 9th NID campaign

Acronyms

BCC	Behavior Change Communication
BCG	Bacillus of Calmette and Guerin
CES	Coverage Evaluation Survey
COSAS	Coverage Survey Analysis System
DPT	Diphtheria, Pertussis and Tetanus
EPI	Expanded Program on Immunization
FWC	Family Welfare Center
IOCH	Immunization and Other Child Health
Mahallah	Smaller localities (smaller than a village)
MOHFW	Ministry of Health and Family Welfare
Mouza	Smallest administrative locality in an Upazila
MSH	Management Sciences for Health
NGO	Non Governmental Organization
NID	National Immunization Day
OPV	Oral Polio Vaccine
TT	Tetanus Toxoid
WHO	World Health Organization

Terminology

This provides the meaning of some of the more technical terms used in this report and a brief explanation of their use.

By card: An immunization given to a child is termed as by card if the date of the dose is entered on an immunization card. Only doses recorded by card are treated as valid data in this survey.

By history: Immunization history collected from a parent's recall is termed as by history. Often no date will be mentioned. This information is only included in crude data.

Crude coverage rate is calculated from the doses recorded by card and/or by history. It is not ascertained whether the doses were given at the correct age and/or following the correct interval (where applicable). Crude data however, helps us to understand how much additional coverage could be achieved if all vaccines were given at the optimum age for the child and following the optimum interval. It also provides useful information on access to the EPI program and on the operational aspects of the provision of health services.

Valid coverage rate is calculated from the vaccinations recorded by card plus history. In the calculation process, first the rate of validity is calculated based on cards only, then this rate is applied to history cases too. Valid data includes only the doses of vaccines that were given after the minimum date of eligibility and/or after the minimum interval necessary to be effective and to protect the child. There is no maximum interval for a dose and therefore a dose administered after 52 weeks is still regarded as valid. By comparing crude coverage with valid coverage data of any particular antigen, one can determine how much coverage was lost due to the inability to give vaccine at the appropriate time.

Invalid doses are those administered at the wrong age and/or at the wrong interval. Doses administered before the minimum age in the case of DPT/Polio 1st doses and Measles vaccine or with less than four weeks interval in the case of DPT or Polio vaccines are classified as "invalid" doses.

The **criteria for a valid dose** used in this survey is the criteria recognized by the Bangladesh EPI program: minimum age for DPT/Polio 1st dose - 6 weeks old; minimum DPT/Polio interval - 4 weeks; minimum age for Measles vaccine - 38 weeks old.

Program access is measured by the percentage of children surveyed who received DPT 1st dose (crude data – by card and history) in the routine immunization session.

Fully immunized means the child has received all the doses it requires (BCG, OPV 1-3, DPT 1-3 and measles).

Missed opportunity refers to a visit of a child to a vaccination center for a dose that he received. However at that time he was also eligible for another dose of antigen that he did not receive. If the missed dose was provided at a later date, it is a *corrected missed opportunity*. If not, it is an *uncorrected missed opportunity*.

SUMMARY RESULTS

Background

The routine EPI program in the municipalities is carried out by a variety of private and public providers at fixed (hospitals, clinics, dispensaries, etc.) and at outreach sites. NGOs and private practitioners also provide immunization services in many places. The municipal authorities are primarily responsible for providing and/or coordinating primary health care including routine EPI services in municipal areas. However, in the absence of an effective management information system and reliable service statistics at municipal level, it is often difficult to assess the level of immunization coverage of the municipalities. In view of this situation, the IOCH decided to conduct a series of coverage evaluation surveys in the selected municipal areas to assess the level of immunization coverage in these municipal areas. As a part of this effort, the IOCH conducted a coverage evaluation survey in the Kushtia Municipality in October 2001.

Objectives

The overall objective of the survey was to assess the level of immunization coverage in the Kushtia Municipality. The specific objectives were to:

- a) assess the level of routine immunization coverage of the children (12-23 months) and find out the reasons for non-immunization and partial immunization.
- b) assess the level of TT immunization coverage among women of 15-49 years of age, irrespective of their marital status, and find out the reasons for non-immunization and partial immunization.
- c) assess the coverage levels of OPV and Vitamin A administered during the 9th NID campaign.

Methodology

The survey employed the WHO recommended 30-cluster survey methodology that has been widely used in many developing countries to assess immunization coverage (the survey methodology and its limitations are presented in Annex A). In all, 30 clusters were randomly selected from the Kushtia Municipality following PPS sampling procedures. A list of the selected clusters is provided in Annex- B and their locations are shown on the maps in page 8. From each cluster, 7 children 12 – 23 months and 7 women of reproductive age (15 – 49 years) irrespective of their marital status were selected following 30 cluster survey methodology to ascertain their routine immunization coverage. Also, 7 children < 5 years (0 – 59 months) were selected to assess the immunization coverage of the 9th National Immunization Campaign.

The WHO standard questionnaires were used for documenting the routine immunization status of the children and women. Also, separate questionnaires were used to collect data on NIDs and reasons for non-immunization and dropouts. The data were collected by the experienced Field Investigators of the Survey Team of the IOCH. Data processing and analysis were done by the Monitoring & Evaluation Unit of the IOCH using COSAS 4.41¹ and EpiInfo.

¹ COSAS (Coverage Survey Analysis System) is a dedicated software for analyzing coverage evaluation survey data.

Coverage levels for the routine immunization of children

Access: Based on crude data (card plus history), 99% children received at least one dose of antigen (DPT 1st dose in this case) from routine immunization sessions. Only 1% children did not receive a dose of any antigen.

Crude coverage between 12-23 months: 99% children received BCG, 98% children received three doses of OPV, 98% received three doses of DPT and 95% received measles vaccine. 94% children were fully immunized.

Valid coverage between 12-23 months: 99% children received BCG, 87% children received three doses of OPV, 87% received three doses of DPT and 94% received measles vaccine. 86% children were fully immunized.

Valid coverage by 12 months: 99% children received BCG, 85% children received three doses of OPV, 85% received three doses of DPT and 84% received measles vaccine. 75% children were fully immunized.

Source of immunization and distance of immunization center: Childhood immunization in this area was provided mostly by the GOB EPI outreach centers (50%), followed by the GOB hospitals and clinics (37%). NGO clinics provided immunization to 13% cases only. All the EPI outreach centers were located within half an hour walking distance from the homes of the children.

Reasons for non-immunization and partial immunization or dropout of children: Only one child out of 210 were never immunized. The primary reasons for his/her non-immunization, as cited by his/her parents, was lack of awareness of need and importance of immunization. As regards reasons for partial immunization or dropout, over half of the parents (58%) reported sickness of their children. The other reasons for partial immunization or dropout cited by the parents included fear of pain of injection (17%), fear of adverse reaction (8%) and other family problems (8%).

Problems detected: The dropout rate for childhood immunization in Kustia Municipality was low, compared to many other municipalities in Bangladesh. There was 2% dropout from DPT1 to DPT3 and 4% from DPT1 to Measles vaccine. There was, however, a number of invalid doses due to early immunization (3% for DPT1, 3% for DPT2, 3% for DPT3 and 1% for Measles). Prevalence of uncorrected missed opportunity for different antigens was also low (1%). Child immunization cards (EPI Cards) were ever given to 99% of the children; but they were available at the time of interview in 56% of the cases. EPI Card retention rate was 57%, which means that 43% of the EPI cards were lost. About 42% of the parents could mention correctly the number of times (i.e. 4 times) their children needed to visit the immunization centers to get fully immunized.

Coverage levels for the routine TT immunization of women

The survey of coverage levels for immunization against tetanus toxoid showed that about 91% of women of reproductive age (15-49 years) received a first dose of TT; 86% received two doses; and 32% received 5 doses of the TT vaccine. About 9% women never received any TT vaccine.

Dropout rate for TT immunization: The dropout rate for TT immunization among the women of reproductive age was quite high. The dropout rate for TT1 to TT2 was 6%, TT2 to TT3 was 14%, TT3 to TT4 was 28% and TT1 to TT5 was 65%. It implies that 65% of those women who received TT1 dose did not get fully immunized for life-long protection against tetanus.

Sources of TT immunization: Majority of the women (56%) received TT vaccine from GOB EPI outreach centers, followed by GOB hospitals and clinics (33%). NGO clinics provided TT immunization to 9% cases and private clinics to 2% cases only.

Reasons for non-immunization and partial immunization of women: Half of the non-immunized women reported that there was no provision of TT vaccine when they had their pregnancies. The other reasons cited for non-immunization were that the women were unaware of the need for immunization (39%) and fear of pain of injection (11%). The primary reasons for partial immunization or dropout were that the women did not know that they needed subsequent doses to get fully immunized (31%) or they were told by the health workers that 2 or 3 doses were enough for them for their pregnancies, and they were not advised (by health worker) for full immunization (22%).

Coverage levels for the 9th NID campaign

During the 9th NIDs, 98% of the children <5 years received OPV in both the rounds. The coverage for OPV in each of the Rounds was 99%. Vitamin A capsules were given to 96% of the eligible children during the 9th NID campaign

The primary reasons for not receiving OPV during the 9th NIDs included lack of awareness of NID campaign, inconvenient timing of NID session and absence of children from homes on the day of NIDs.

Majority of the parents learned about the 9th NID Campaign from the IPC during home visits by the GOB/municipal field workers (72%), followed by miking (40%). Television/radio as a source of information was cited by 27% of the parents; while 13% came to know about the NIDs from other family members/ neighbors.

Suggested solutions

The survey indicates the need for appropriate BCC activities and counseling to make the parents/caretakers understand the importance of full immunization of their children. The information must stress that each child, irrespective of sex, must be fully immunized before 12 months of age, and how to achieve full immunization (the time and place of the immunization sessions and the number of doses required). All women of childbearing age must be informed about how to prevent tetanus with 5 doses of TT vaccination to protect herself and her newborn child.

Although the quality of services was not very bad, as reflected by the percentage of invalid doses (3% for DPT1, 3% for DPT2, 3% for DPT3 and 1% for Measles) and dropout rates (2% for DPT1 to DPT3 and 4% for DPT1 to measles), there is still room for further improvement of quality of services by proper screening of eligibility of clients for each antigen at the time of vaccination, and by adequate counseling of mothers. There is also a need to further train the service providers to help them keep up with EPI policies and guidelines and increase their capacity for counseling parents and women of reproductive age about EPI.



TABLES AND CHARTS

Table 1: Routine immunization coverage levels of the children

Name of the vaccine	Coverage (%) Immunization of 12-23 months age group		Coverage (%) Immunized by 12 months of age
	Crude data (Access)	Valid data	Valid data
BCG	99	99	99
OPV1	100	96	96
OPV2	99	92	92
OPV3	98	87	85
DPT1	99	96	96
DPT2	99	92	92
DPT3	98	87	85
Measles	95	94	84
Fully immunized	94	86	75
Zero Dose	1	-	-

Chart-1: Drop-out rate for childhood immunization




Table 2: Invalid doses of immunization provided to the children

Antigens	Percentage
DPT1	3
DPT2	3
DPT3	3
Measles	1

Table 3: Missed opportunities by antigens

Name of the vaccine	Uncorrected		Corrected	
	Number	Percent	Number	Percent
BCG	2	1	3	1
DPT1	2	1	7	3
DTP2	1	1	6	3
DPT3	1	1	4	2
OPV1	2	1	7	3
OPV2	1	1	6	3
OPV3	1	1	4	2
Measles	1	1	1	.5

Table 4: EPI card availability and retention

Card Status	Number	Percentage
EPI card available	118	56
EPI card ever given	208	99
EPI card retention	118	57

Chart 2: Sources of childhood immunization services




Table 5: Distance between the child’s home and the vaccination site

Time (Minutes)	Number	Percentage
1-5 Min.	130	62
6-10 Min.	78	37
11-30 Min.	2	1
Total	210	100

Table 6: Reasons for non-immunization and partial immunization of the children

Reasons	Non-immunized (%) (N=1)	Partially immunized (%) (N=12)
Did not know about need of immunization	100	-
Fear of adverse reaction	-	8
Family problem/mother sick	-	8
Child sick, was not taken to site	-	58
Fear of pain of injection	-	17
Health worker's behavior was not good	-	9

Chart 3: Respondents' knowledge about required visits to immunization centers




Chart 4: Routine immunization coverage levels for TT of women (15-49 years)




Chart 5: TT Immunization drop-out rate among women 15-49 years




Chart 6: Providers of TT immunization




Table 7: Reasons for non-immunization and partial immunization for TT of the women

Reasons	Non-immunization (%) (N=18)	Partially immunization (%) (N=125)
Next dose is not yet due	-	29
Don't feel need for immunization	39	-
Health worker did not specify the next dose	-	10
As per HW advice 2/3 TT is enough during the pregnancy	-	22
Unaware of need of next dose	-	31
In our times TT immunization was not in practice	50	-
Fear of injection	11	6
Health worker was not available at the site	-	1
Family problem	-	1


Table 8: Coverage of the 9th NID Campaign

Round	OPV %	Vit "A" %
1 st round	99	-
2 nd round	99	96
Both rounds	98	-
Any round	99	-

Table 9: Reasons for non-immunization of OPV during 9th NID campaign

Reasons	1 st Round %	2 nd Round %
Did not know about NID	67	-
Time inconvenient	-	50
Away from home	-	50
Others	33	-

Chart 7: Source of information about the 9th NID campaign



The following are extracts from **Anthony G Turner, Robert J Magnani and Muhammad Shuaib's** article entitled **"A not quick as quick but much cleaner alternative to the Expanded Programme on Immunization (EPI) cluster survey design"** published in the *International Journal of Epidemiology* in 1996, volume 25, Issue No. 1, pages 198-203.

The standard EPI Cluster Survey Design

"The sample design for the EPI Cluster Survey is a two-stage design involving the selection of 30 primary sampling units or 'clusters' (usually village or other area units), from which 210 children with a target age range (usually 12-23 months) are chosen, seven children per cluster. The sample size of 210 children (per domain or stratum) is mandated by the desire to estimate the level of immunization coverage to within +/- 10 percentage points of the true population proportion with 95% statistical confidence, assuming a design effect (i.e. *deff*) of 2.0. Based upon prior experience with immunization coverage surveys (primarily in the US), 30 clusters are generally thought to be necessary to yield sufficiently reliable estimate."

"In the standard design, clusters are chosen from a list of primary sampling units (i.e. villages, urban communities, census enumeration areas etc.) through systematic random sampling with probability proportional to estimated size (*ppes*). The latest estimates of cluster population sizes, which are assumed to be proportional to the number of children in the target age group in each cluster, are typically used as measures of size. The 30 clusters so chosen are then visited by survey field staff who carry out the second stage of sample selection and conduct the household interviews. "

"The original EPI design called for sample children to be chosen randomly from a list of all eligible children in each sample cluster. However, because the creation of lists of households and children tends to be time-consuming, costly, and unfeasible in some settings, this procedure is only infrequently used in actual practice. Instead, one of several simplified second stage sampling procedures is commonly used. In one variant, children are selected by first choosing a random direction from a central location in a village or community (e.g. by spinning a bottle). The number of households in that direction to the edge of the community is then counted, and one household is randomly chosen to be the first sample household. Subsequent households are chosen by visiting the nearest neighboring households until information has been gathered on seven children. In a yet simpler variant, a direction from a central starting point is randomly chosen as described above and households are contacted as the interviewer moves in the chosen direction until the required information has been gathered for seven children."

"The second stage sampling methods described above are 'quota sampling procedures' and some of the problems resulting from the use of this approach have been noted over the years."

"First, quota sampling does not ensure that every eligible member of the target population has a known, non-zero chance of being selected. Hence, the standard EPI design, as it is usually applied, is not a true probability sample design."

"A second problem concern sampling weights. However, since measures of size in sampling frames are often inaccurate due to census errors and changes in population since the census was taken, application of the standard EPI Cluster Survey method does not automatically result in a self weighting sample. The survey data must be weighed in order to yield unbiased estimates. However, since selection probabilities are not known in most EPI Cluster Survey applications, sampling weights can not be calculated."

"Thirdly, a computer simulation study demonstrates that the EPI Cluster Survey based upon quota sampling at the second stage of sample selection is considerably more prone to sampling bias than conventional cluster sampling, particularly where immunized children are 'pocketed' within clusters. "

"Finally, there is the issue of how second stage sample selection should proceed in surveys with multiple measurement objectives."

List of Selected Clusters for the Survey

Ward No.	Mahalla Name	Total HH	Total Population	Cluster No.
1	Kamalapur	457	2583	1 – 2
	Thana para	1400	7806	3 – 6
2	Amla para	879	4816	7 – 8
	Pashim Arua para	1070	5977	9 – 11
	Pashim Mil para	154	945	12
3	Chourash (part)	246	1406	13
	Court para	1504	8779	14 – 16
	C & B staff Quarter	474	2467	17
	Gour Gopal Gali	216	1286	18
	Housing Estate (Block A)	421	2185	19
	Housing Estate (Block B)	436	2428	20
	Housing Estate (Block C)	369	2148	21
	Kali sankar pur	1130	6523	22 – 23
4	Bahadur Khali	293	1307	24
	Bahadur Khan gali (part)	542	3090	25
	Hari sankar pur	1065	5784	26 – 28
	Purba mill para	978	5220	29 – 30

List of Never Vaccinated Children Identified by Clusters

Ward No.	Mahalla Name	Total HH	Total Population	Cluster No.	Never Vaccinated Children
1	Kamalapur	457	2583	1 - 2	-
	Thana para	1400	7806	3 - 6	-
2	Amla para	879	4816	7 - 8	-
	Pashim Arua para	1070	5977	9 - 11	-
	Pashim Mil para	154	945	12	-
3	Chourash (part)	246	1406	13	-
	Court para	1504	8779	14 - 16	1
	C & B staff Quarter	474	2467	17	-
	Gour Gopal Gali	216	1286	18	-
	Housing Estate (Block A)	421	2185	19	-
	Housing Estate (Block B)	436	2428	20	-
	Housing Estate (Block C)	369	2148	21	-
4	Kali sankar pur	1130	6523	22 - 23	-
	Bahadur Khali	293	1307	24	-
	Bahadur Khan gali (part)	542	3090	25	-
	Hari sankar pur	1065	5784	26 - 28	-
	Purba mill para	978	5220	29 - 30	-

Acknowledgements

Survey coordination:

Mr. Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

Survey management:

Mr. Jagadindra Majumder, Field Survey Manager, IOCH/MSH

Data analysis:

Mr. Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

Mr. Biplob Banerjee, Surveillance Data Manager, IOCH/MSH

Report writing:

Mr. Md. Mafizur Rahman, Monitoring & Evaluation Specialist, IOCH/MSH

Report review:

Dr. Pierre Claquin, Chief of Party, IOCH/MSH

Digital map preparation:

Mr. Din Mohammed, Monitoring and Evaluation Assistant, IOCH/MSH

Cover photo:

Ms. Moumina Dorgabekova/Image Jinn

Survey Team members, IOCH/MSH:

Mr. Md. Abdul Hamid, Field Investigator

Mr. Md. Saiful Islam, Field Investigator

Ms. Krishna Rani Shil, Field Investigator

Ms. Khaleda Akhter, Field Investigator

Ms. Mahamuda Parveen, Field Investigator

Ms. Aung Ma Ching Marma, Field Investigator

Ms. Niva Rani Taju, Field Investigator

Ms. Sultana Parvin Maya, Field Investigator

List of IOCH Survey/Research/Technical Reports

Survey Reports

1. Vaccination Coverage Survey of the Slums of Rajshahi City Corporation- January 2000. Survey Report No. 1. May 2000
2. Vaccination Coverage Survey of the Selected Unions along the North-western Border of Bangladesh- February 2000. Survey Report No. 2. June 2000
3. Vaccination Coverage Survey of the Selected Unions along the South-west Border of Bangladesh- February 2000. Survey Report No. 3. July 2000
4. Vaccination Coverage Survey of the Slums of Khulna City Corporation- January 2000. Survey Report No. 4. July 2000
5. Vaccination Coverage Survey of the Slums of Chittagong City Corporation- January 2000. Ward Number 1 to 18. Survey Report No. 5. July 2000
6. Vaccination Coverage Survey of the Slums of Chittagong City Corporation- January 2000. Ward Number 19 to 41. Survey Report No. 6. July 2000
7. Vaccination Coverage Survey of the Dinajpur Municipality- January 2000. Survey Report No. 7. July 2000
8. Vaccination Coverage Survey of the Noakhali Municipality- January 2000. Survey Report No. 8. July 2000
9. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 1, 2 & 4. Survey Report No. 9. July 2000
10. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 5, 6 & 7. Survey Report No. 10. July 2000
11. Vaccination Coverage Survey of the Slums of Dhaka City Corporation- January 2000. Dhaka Slums of Zones 3, 8, 9 & 10. Survey Report No. 11. July 2000
12. Vaccination Coverage Survey of the Tribal and Non-tribal Populations in the North-east Border Areas of Bangladesh. Survey Report No. 12. August 2000
13. Vaccination Coverage Survey of the Sylhet Municipality – January 2000. Survey Report No. 13. August 2000.
14. Vaccination Coverage Survey of the Kishoreganj Municipality – April 2000. Survey Report No. 14. September 2000.
15. Vaccination Coverage Survey of the Rangpur Municipality – May 2000. Survey Report No. 15. September 2000.
16. Vaccination Coverage Survey of the Greater Faridpur Municipalities – June 2000. Survey Report No. 16. September 2000.
17. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Chittagong, Khulna and Rajshahi City Corporation Slums – September 2000. Survey Report No. 17. November 2000.
18. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Dhaka City Corporation Slums – September 2000. Survey Report No. 18. November 2000
19. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Slums of selected 27 Municipalities– September 2000. Survey Report No. 19. November 2000
20. Results of Coverage Evaluation Survey of Routine EPI and August 2000 OPV + TT (NNT) Campaign, Hard-to-reach and High-risk rural areas – September 2000. Survey Report No. 20. November 2000
21. Vaccination Coverage and Other Health Care Practices Survey in the Pabna Char Areas—August 2000. Survey Report No. 21. January 2001.
22. Vaccination Coverage Survey of the Mymensingh Municipality – April 2000. Survey Report No. 22. February 2001.
23. Vaccination Coverage Survey of the Jessore Municipality – October 20-30, 2000. Survey Report No. 23. February 2001.
24. Vaccination Coverage Survey of the Comilla Municipality - October 1-9, 2000. Survey Report No. 24. February 2001
25. Vaccination Coverage Survey of the Pabna Municipality - July 2000. Survey Report No. 25. February 2001
26. Vaccination Coverage Survey of the Sirajganj Municipality - July 2000. Survey Report No. 26. February 2001
27. Vaccination Coverage Survey of the Bogra Municipality - February 2001. Survey Report No. 27. April 2001
28. Vaccination Coverage Survey of the Peri-urban Areas of DCC- Nov. 2000 (Peri-urban Area Survey-1). Survey Report No. 28. July 2001
29. Vaccination Coverage Survey of the Peri-urban Areas of DCC- Nov. 2000 (Peri-urban Area Survey-2). Survey Report No. 29. July 2001
30. Vaccination Coverage Survey of the Patuakhali, Jhalokathi and Pirojpur Municipalities- Aug. 2000. Survey Report No. 30. July 2001
31. Vaccination Coverage Survey of the Bandarban, Rangamati and Khagrachari Municipalities- January 2001. Survey Report No. 31. July 2001
32. Vaccination Coverage Survey of the Naogaon Municipality- February 2001. Survey Report No. 32. July 2001

33. Vaccination Coverage Survey of the Tongi Municipality- March 2001. Survey Report No. 33. July 2001
34. Vaccination Coverage Survey of the Tangail Municipality- March 2001. Survey Report No. 34. July 2001
35. Vaccination Coverage Survey of the Savar Municipality- May 2001. Survey Report No. 35. July 2001
36. Vaccination Coverage Survey of the Rangpur District- June 2001. Survey Report No. 36. July 2001
37. Vaccination Coverage Survey of the Nawabganj Municipality- June 2001. Survey Report No. 37. July 2001
38. Vaccination Coverage Survey of the Expanded Areas of the Proposed Barisal City Corporation- June 2001. Survey Report No. 38. July 2001
39. Vaccination Coverage Survey of the Barisal Municipality- June 2001. Survey Report No. 39. July 2001
40. Vaccination Coverage Survey of the Kadam Rasul Municipality- July 2001. Survey Report No. 40. August 2001
41. Vaccination Coverage Survey of the Jamalpur Municipality- July 2001. Survey Report No. 41. August 2001
42. Vaccination Coverage Survey of the Brahmanbaria Municipality- June 2001. Survey Report No. 42. September 2001
43. Vaccination Coverage Survey of the Gazipur Municipality- Vaccination Coverage Survey of the Hard-to-Reach Unions of Dewanganj, Raumari and Rajibpur Upalizas- July 2001. Survey Report No. 45. September 2001
44. Vaccination July 2001. Survey Report No. 43. September 2001
45. Vaccination Coverage Survey of the Narsingdi Municipality- August 2001. Survey Report No. 44. September 2001
46. Coverage Survey of the Satkhira Municipality- September 2001. Survey Report No. 46. September 2001
47. Coverage Survey of the Barguna Municipality- October 2001. Survey Report No. 47. November 2001
48. Coverage Survey of the Bhola Municipality- October 2001. Survey Report No. 48. November 2001

Unicef & IOCH Survey Reports

1. Vaccination Coverage Survey of the Teknaf and Ukhia Upazilas- February 2000. Survey Report No. 01, August 2000
2. Vaccination Coverage Survey of the Brahmanbaria Sadar Upazila- February 2000. Survey Report No. 02, August 2000
3. Vaccination Coverage Survey of the Debidwar Upazila- February 2000. Survey Report No. 03, August 2000
4. Vaccination Coverage Survey of the Madaripur Upazila- February 2000. Survey Report No. 04, August 2000
5. Vaccination Coverage Survey of the Maulvi Bazar District- February 2000. Survey Report No. 05, August 2000
6. Vaccination Coverage Survey of the Raumari Upazila - February 2000. Survey Report No. 06, August 2000
7. Vaccination Coverage Survey of the Gangachara Upazila - February 2000. Survey Report No. 07, August 2000
8. Vaccination Coverage Survey of Chittagong Hill Tracts - February 2000. Survey Report No. 08, October 2000

Technical Report

1. Joint National/International Review of EPI Program in Urban Areas of Bangladesh—23 January – 3 February 2000. Technical Report No. 01, July 2000

Additional copies of any of these reports, if needed, will be provided free of cost on request to:

Mamunul Haque, Communications Advisor, IOCH. E-mail: mh@citechco.net