



**IOCH**  
**Immunization and Other Child Health Project**

**Vaccination Coverage Survey of the  
Manikganj Municipality**

**January 2003**

**Survey Report No. 100**

**This survey was conducted by IOCH, a project of Management Sciences for Health,  
funded by USAID under AID contract No. HRN-I-01-98-00033-00, Task Order No. 01**

House 1, Road 23, Gulshan 1, Dhaka 1212, Bangladesh  
Tel: 8828596, 8829279, 8813611, 8813410  
Fax: 880-2-8826229  
E-mail: [ioch@citechco.net](mailto:ioch@citechco.net)

**February 2003**

## Table of Contents

List of tables and figures	2
Acronyms	3
Terminology	4
<b>Summary Results</b>	<b>5</b>
Background	5
Objectives	5
Methodology	5
Routine immunization coverage levels of children	6
Routine TT immunization coverage levels of women	7
Coverage levels of the SNID Campaign- 2002	8
Conclusions and recommendations	9
Map showing the locations of the selected clusters	12
Tables and figures	13
<b>Annexures</b>	
Annex A: List of selected clusters for the survey	22
Annex B: List of never vaccinated children by clusters	22
Acknowledgements	23
List of IOCH reports	24

## **List of Tables**

- Table 1:** Routine immunization coverage levels of the children  
**Table 2:** Routine immunization coverage levels of the children by gender  
**Table 3:** Invalid doses of immunization provided to the children  
**Table 4:** Missed opportunities by antigens  
**Table 5:** EPI cards availability and retention  
**Table 6:** Time required to reach the EPI center by mode of transportation  
**Table 7:** Reasons for non-immunization and partial immunization of the children  
**Table 8:** Age distribution of women never receiving TT vaccine  
**Table 9:** Interval between TT1 and TT2, TT2 and TT3, TT3 and TT4 and TT4 and TT5  
**Table 10:** Children born protected against tetanus  
**Table 11:** Knowledge about number of TT doses required for life-long protection against tetanus  
**Table 12:** TT cards availability and retention  
**Table 13:** Reasons for non-immunization and partial immunization for TT of the women  
**Table 14:** OPV and Vitamin A coverage during the SNIDs- 2002  
**Table 15:** Sources of OPV during the SNIDs- 2002  
**Table 16:** Time required to reach the SNID sites by mode of transportation  
**Table 17:** Households visited during the child-to-child search of the 2<sup>nd</sup> round of the SNIDs  
**Table 18:** Reasons for not receiving OPV from the SNID sites

## **List of Figures**

- Figure 1:** Drop-out rate for childhood immunization  
**Figure 2:** Knowledge about required visits to immunization centers for full immunization  
**Figure 3:** Sources of child immunization services  
**Figure 4:** Routine immunization coverage levels for TT of the women 15-49 years  
**Figure 5:** TT immunization dropout rate among women 15 – 49 years  
**Figure 6:** Providers of TT immunization  
**Figure 7:** Sources of information about the SNID Campaign- 2002

## Acronyms

BCC	Behavior Change Communication
BCG	Bacillus of Calmette and Guerin
CES	Coverage Evaluation Survey
COSAS	Coverage Survey Analysis System
DPT	Diphtheria, Pertussis and Tetanus
EPI	Expanded Program on Immunization
FWC	Family Welfare Center
IOCH	Immunization and Other Child Health
Mahallah	Smaller localities (smaller than a village)
MOHFW	Ministry of Health and Family Welfare
Mouza	Smallest administrative locality in an Upazila
MSH	Management Sciences for Health
NGO	Non Governmental Organization
NID	National Immunization Day
OPV	Oral Polio Vaccine
SNID	Sub-national Immunization Day
TT	Tetanus Toxoid
WHO	World Health Organization

## Terminology

This provides the meaning of some of the more technical terms used in this report and a brief explanation of their use.

**By card:** An immunization given to a child is termed as by card if the date of the dose is entered on an immunization card. Only doses recorded by card are treated as valid data in this survey.

**By history:** Immunization history collected from a parent's recall is termed as by history. Often no date will be mentioned. This information is only included in crude data.

**Crude coverage** rate is calculated from the doses recorded by card and/or by history. It is not ascertained whether the doses were given at the correct age and/or following the correct interval (where applicable). Crude data however, helps us to understand how much additional coverage could be achieved if all vaccines were given at the optimum age for the child and following the optimum interval. It also provides useful information on access to the EPI program and on the operational aspects of the provision of health services.

**Valid coverage** rate is calculated from the vaccinations recorded by card plus history. In the calculation process, first the rate of validity is calculated based on cards only, then this rate is applied to history cases too. Valid data includes only the doses of vaccines that were given after the minimum date of eligibility and/or after the minimum interval necessary to be effective and to protect the child. There is no maximum interval for a dose and therefore a dose administered after 52 weeks is still regarded as valid. By comparing crude coverage with valid coverage data of any particular antigen, one can determine how much coverage was lost due to the inability to give vaccine at the appropriate time.

**Invalid doses** are those administered at the wrong age and/or at the wrong interval. Doses administered before the minimum age in the case of DPT/Polio 1<sup>st</sup> doses and Measles vaccine or with less than four weeks interval in the case of DPT or Polio vaccines are classified as "invalid" doses.

The **criteria for a valid dose** used in this survey is the criteria recognized by the Bangladesh EPI program: minimum age for DPT/Polio 1<sup>st</sup> dose - 6 weeks old; minimum DPT/Polio interval - 4 weeks; minimum age for Measles vaccine - 38 weeks old.

**Dropout** cases refer to the children/women who have initially received at least one dose of any antigen and then failed to receive the subsequent doses to get fully immunized. Dropout rate implies the inability of the EPI to follow-up and protect the cohort of children initially reached out.

**Program access** is measured by the percentage of children surveyed who received DPT 1<sup>st</sup> dose (crude data – by card and history) in the routine immunization session.

**Fully immunized** means the child has received all the doses it requires (BCG, OPV 1-3, DPT 1-3 and measles).

**Missed opportunity** refers to a visit of a child to a vaccination center for a dose that he received. However at that time he was also eligible for another dose of antigen that he did not receive. If the missed dose was provided at a later date, it is a *corrected missed opportunity*. If not, it is an *uncorrected missed opportunity*.

## SUMMARY RESULTS

### Background

The routine EPI program in the municipalities is carried out by a variety of private and public providers at fixed (hospitals, clinics, dispensaries, etc.) and at outreach sites. NGOs and private practitioners also provide immunization services in many places. The municipal authorities are primarily responsible for providing and/or coordinating primary health care including routine EPI services in municipal areas. However, in the absence of an effective management information system and reliable service statistics at municipal level, it is often difficult to assess the level of immunization coverage of the municipal areas. In view of this situation, the IOCH decided to conduct a series of coverage evaluation surveys in the selected municipal areas to assess the level of immunization coverage in these areas. As a part of this effort, the IOCH conducted a coverage evaluation survey in the Manikganj Municipality in January 2003.

### Objectives

The overall objective of the survey was to assess the level of immunization coverage in the Manikganj Municipality. The specific objectives were to:

- a) assess the level of routine immunization coverage of the children (12-23 months) and find out the reasons for non-immunization and partial immunization;
- b) assess the level of TT immunization coverage among women of 15-49 years of age, irrespective of their marital status, and find out the reasons for non-immunization and partial immunization; and
- c) assess the coverage levels of OPV and Vitamin A administered during the SNID Campaign- 2002.

### Methodology

The survey employed the WHO recommended 30-cluster survey methodology that has been widely used in many developing countries to assess immunization coverage. In all, 30 clusters were randomly selected from the Manikganj Municipality following PPS sampling procedures. A list of the selected clusters is provided in Annex- A and their locations are shown on the maps in page 12. From each cluster, 7 children 12 – 23 months and 7 women of reproductive age (15 – 49 years) irrespective of their marital status were selected following 30 cluster survey methodology to ascertain their routine immunization coverage. Also, 7 children < 5 years (0 – 59 months) were selected to assess the immunization coverage of the SNIDs- 2002

The WHO standard questionnaires were used for documenting the routine immunization status of the children and women. Also, separate questionnaires were used to collect data on SNIDs and reasons for non-immunization and dropouts. The data were collected by the experienced Field Investigators of the Survey Team of the IOCH. Data processing and analysis were done by the Monitoring & Evaluation Unit of the IOCH using COSAS 4.41<sup>1</sup> and EpiInfo.

---

<sup>1</sup> COSAS (Coverage Survey Analysis System) is a dedicated software for analyzing coverage evaluation survey data.

## **Coverage levels for the routine immunization of children**

***Access to child immunization:*** Based on crude data (card plus history), 98% children received at least one dose of antigen (DPT 1<sup>st</sup> dose in this case) from routine immunization sessions. 2% children did not receive a dose of any antigen.

***Crude coverage of 12-23 months age group:*** 98% children received BCG, 95% children received three doses of OPV, 95% received three doses of DPT and 91% received measles vaccine. 91% children were fully immunized.

***Valid coverage of 12-23 months age group:*** 98% children received BCG, 86% children received three doses of OPV, 86% received three doses of DPT and 87% received measles vaccine. 81% children were fully immunized.

***Valid coverage by 12 months:*** 98% children received BCG, 86% children received three doses of OPV, 86% received three doses of DPT and 84% received measles vaccine. 78% children were fully immunized.

***Routine immunization coverage by sex:*** There was no significant sex difference in accessing routine child immunization services. Boys' access to immunization, as measured by the crude coverage of DPT1, was two percentage points higher than that of the girls (99% for boys vs. 97% for girls). But, the measles coverage of the boys was one percentage point lower than that of the girls, resulting in one percentage point lower crude FIC for boys than that of the girls (90% crude FIC for boys vs. 91% crude FIC for girls). The proportions of invalid doses for DPT and OPV were higher among the girls than the boys, resulting in higher valid FIC for the boys than that for the girls (83% valid FIC for boys vs. 79% valid FIC for girls).

***Dropout rates:*** Although access to child immunization was quite good (98% for DPT1), there were 3% dropouts from DPT1 to DPT3 and 8% from DPT1 to measles.

***Invalid doses:*** 4% of the DPT1 doses were administered before 6 weeks of age of the children, and 3% of the measles doses were administered before 38 weeks of age of the children. Besides, 2% of the DPT2 doses 1% of the DPT3 doses were given before 4 weeks interval between the doses.

***Missed opportunities:*** Total missed opportunities (uncorrected plus corrected) for different antigens ranged from 0 to 3%. The prevalence of uncorrected missed opportunities for different antigens ranged 1% - 2%. The composite index for total missed opportunities was 25, reflecting the need for improving the quality of screening during vaccination sessions.

***EPI card retention:*** 98% of the children interviewed were ever given EPI cards; however, EPI cards were available with 60% of the respondents at the time of interview. EPI card retention rate was 61% only, as 39% of the EPI cards were lost.

***Knowledge about required visit to immunization center for full immunization:*** 39% of the mothers interviewed did not have any idea about how many times a child was required to be taken to an immunization center to get fully immunized; while 12% had wrong idea about it.

Only 49% of the mothers could mention correctly the number of times (i.e. 4 times) a child is required to visit immunization center to get fully immunized.

**Sources of immunization services and distance of vaccination centers:** Childhood immunization in this area was provided mostly by the GOB hospitals (36%), followed by the municipal EPI outreach centers (28%). NGO clinics and GOB EPI outreach centers provided EPI services to 22% and 6% of the cases respectively. All the EPI outreach centers could be reached within 15 minutes travel time, and 92% of them were located within 10 minutes walking distance from the homes of the children.

**Reasons for non-immunization and partial immunization or dropout of children:** The primary reasons for non-immunization of children were parents lack of awareness of need and importance of immunization (50%), and fear of pain of vaccination (50%). The primary reasons for partial immunization or dropout included lack of knowledge about the need of subsequent doses or measles vaccination for getting fully immunized (38%) and sickness of the children (31%).

### **Coverage levels for the routine TT immunization of women**

**TT immunization coverage:** Access to TT immunization for the women 15 – 49 years was fairly good. 87% of the women received TT1. The corresponding figures for TT2, TT3 and TT4 were 82%, 58% and 42% respectively. Only 24% of the women received TT5, which provide lifelong protection against tetanus. 13% of the women never received any TT vaccine.

**Age distribution of women never receiving TT immunization:** The younger women, as well as older ones are less likely to receive TT vaccine. 39% of the women under 20 years and 37% of the women over 35 years of age had never received any dose of TT vaccine; while the corresponding figures ranged from 5% to 7% for the women of other age-groups.

**TT immunization dropout rates:** TT immunization dropout rates were high. The dropout rate for TT2 to TT3 was 30%. The corresponding rates for TT3 to TT4 and TT4 to TT5 were 26% and 44% respectively. The dropout rate for TT1 to TT5 was as high as 73%, indicating that 73% of the women who received first dose of TT did not complete 5 doses TT immunization schedule.

**Invalid TT doses:** A significant proportion of TT doses were invalid as they were administered before the minimum required interval between the doses. 19% of the TT3 doses were given before 6 months interval between TT2 and TT3, and as such were invalid. 72% of the TT4 doses were invalid, since they were given before one year interval between TT3 and TT4; similarly, 47% of the TT5 doses were invalid for the same reason.

**Protection against tetanus at birth:** 98% of the newborn babies were found protected against tetanus, indicating that 2% newborn babies were still not protected against tetanus at birth.

**Knowledge about full TT immunization:** 90% of the women did not have correct knowledge about the number of TT doses required for a woman for full immunization. Only 10% women could correctly mention that a woman was required to receive 5 doses of TT vaccine for full immunization for lifelong protection against tetanus.

***TT card retention:*** 86% of the women were ever given TT cards; however, TT cards were available with 22% of the women only at the time of interview. TT card retention rate was 25% only, as 75% of the TT cards were lost.

***Sources of TT immunization:*** Majority of the women received TT vaccine from the hospitals (26%), followed by the NGO clinics (24%). Municipal EPI outreach centers and GOB EPI outreach centers provided TT immunization to 23% and 12% of the cases respectively. Private clinics provided immunization services to 5% of the cases only.

***Reasons for non-immunization and partial immunization of women:*** The primary reasons for non-immunization of TT cited by majority of the women were lack of awareness of need and importance of TT immunization (32%) and fear of injection (36%). The primary reasons for partial TT immunization or dropout were that the women were not aware of the need of subsequent doses to get fully immunized against tetanus (60%), or the health workers did not advise for 5 dose TT schedule for full TT immunization for lifelong protection (15%).

### **Coverage levels for the SNIDs- 2002**

***OPV and Vitamin A coverage:*** 98% of the children <5 years received OPV in both the rounds of the SNIDs- 2002. The coverage for OPV was also 99% in each of the rounds. Vitamin A capsules were given to 99% of the eligible children (12 – 59 months of age).

***Sources of OPV during the SNIDs:*** Most of the children received OPV from the SNID sites (99% in each round). None received OPV during child-to-child search.

***Distance of SNID sites and mode of transportation used:*** All the SNID sites were located within 10 minutes walking distance from the homes of the children.

***Households visited during child-to-child search:*** 88% of the households were visited by the health workers/volunteers during child-to-child search in the 2<sup>nd</sup> round of the SNIDs, as reported by the parents/respondents.

***Sources of information of the SNID:*** Majority of the parents learned about the SNIDs from miking (58%), followed by relatives and neighbors (34%). About one-third of the parents (33%) came to know about the NIDs from the municipal worker. Television as a source of information was cited by 26% of the parents.

***Reasons for not receiving vaccines from the SNID sites:*** The primary reason for not receiving OPV from the SNID sites included: a) lack of information of NIDs (50% in the 1<sup>st</sup> round and 33% in the 2<sup>nd</sup> round); b) preoccupation of the parents on NID day (33% in the 2<sup>nd</sup> round) and b) children were away from home on NID day (34% in the 2<sup>nd</sup> round).

## **Conclusions and recommendations**

Access to child immunization was quite good (98% for DPT1); but this high access dropped to 91% for fully immunized children because of dropouts of different antigens. (Dropout rate for DPT1 to DPT3 was 3% and DPT1 to measles was 8%). Similarly, access to TT immunization for the women 15 – 49 years (irrespective of their marital status) was also good. 87% of the women received the first dose of TT; but TT dropout rate was very high (73% for TT1 to TT5), resulting in very low coverage of TT5 (24%), which provides lifelong protection against tetanus. Such high dropout rates, especially for TT immunization, imply inability of the EPI program to follow-up and protect the cohort of children and women initially reached out.

- ***Programmatic strategies must be undertaken to reduce the existing high dropout rates in both child immunization and TT immunization. The program should focus on quality of counseling of mothers/women (on immunization) by the health workers. The health worker at the first contact must counsel the mother/woman properly to motivate her to return and to get herself and/or her child fully immunized. He/she must:***
  - *explain to mother/woman the importance of full immunization, and concept of 8 doses and 4 visits required for full immunization of a child;*
  - *tell mother/woman of reproductive age the benefit and importance of full TT immunization and the concept of 5 dose TT schedule for lifelong protection against tetanus;*
  - *issue an EPI card/TT card dully filled out and explain the mother/woman the importance of keeping the EPI card/TT card in safe and bringing it on the next due date;*
  - *tell the mother/woman clearly when she should come back for next doses, and inform her that the date is written in the EPI card/ TT card so that she can check it if forgets the date;*
  - *inform the mother/woman of possible side effects of injection and how it should be handled;*
  - *pay attention to the hospitality at the clinic/EPI center, and to supportive environment.*
- ***Program managers and field supervisors should ensure that EPI sessions are held as per plan, and at a regular and adequate interval***
- ***The service providers should be given refresher training to improve their technical skills on counseling of mothers/women on immunization***

There were also too many invalid doses in child immunization (4% for DPT1, 2% for DPT2, 1% for DPT3 and 3% for measles), which further reduced the crude full immunization coverage of 91% to 81% when validity of doses was taken into account. Total missed opportunities for different antigens ranged 0-3%. Like child immunization, a very high proportion of TT doses were invalid (19% for TT3, 72% for TT4 and 47% for TT5). These high rates of invalid doses and missed opportunities reflect service providers' inability to screen the clients properly.

- ***Emphasis should be given on screening of clients for immunization to avoid or reduce invalid doses and missed opportunities. The service providers must:***
  - *screen properly each and every child/ woman to decide his/her eligibility for a specific dose of specific antigen;*

- *check EPI card/ EPI register/ TT card or any other record to decide on the eligibility of a particular dose of specific antigen(s);*
- *if a child/woman is found not eligible for a dose today, ask him/her to return at a specified date and explain the reason to her/mother clearly and patiently.*
- *check if there is any missed opportunity for other antigens.*
- *The service providers should be given refresher training to improve their technical skills on screening of clients for immunization.*

EPI card (child immunization card) and TT card play an important role in ensuring good quality of immunization services. It helps the mothers to adhere to immunization schedule, as well as assists the service providers to screen the children for specific doses of specific antigens. Unfortunately, the retention rates of both the EPI card and TT card were very low, 61% and 25% respectively.

- *Mothers/women should be explained the benefits and importance of EPI cards/ TT cards for immunization of themselves and their children.*
- *Mothers should be asked to preserve the EPI card safely until the child is 5 years old, and to bring the card with them whenever they come to the clinic/ EPI center for immunization.*
- *Women should also be asked to carefully preserve their TT cards, and to bring their TT cards with them whenever they come to clinic/ EPI center for TT immunization.*
- *In the case of loss of EPI card/TT card, it should be provided over and over, and the history of the earlier vaccinations accurately recorded again and again, if necessary.*

Mothers have a poor understanding of full immunization. 51% of the mothers could not mention how many times a child was required to be taken to EPI center to get fully immunized. Similarly, 90% of the women 15 – 49 years did not know how many TT doses were required for a woman for lifelong protection against tetanus.

- *During IPC between the mother/woman and the service provider and/or at the first contact, the mother/woman should be clearly explained the importance of full immunization of children and women, and of the immunization schedule of full immunization for both children and women.*

Lack of awareness of need and importance of immunization and fear of pain of vaccination were cited as primary reasons for non-immunization of child and women; while lack of knowledge about the need and importance of subsequent doses to get fully immunized resulted in high drop out rates for child and TT immunization.


- *The program should strengthen BCC activities to inform the community of importance and benefits of immunization in general, and to motivate the mothers/women to get themselves and their children fully immunized in particular. Very selective and focused mass media campaign, in addition to IPC by health workers, may also be conducted to achieve this end.*

2% of the children 0 –59 months did not receive OPV in both the rounds of the SNIDs- 2002. 12% of the households were not visited by the health workers/volunteers during child-to-child search in the 2<sup>nd</sup> round of the SNIDs.

- ***Area specific innovative strategies suitable to local situation have to be undertaken during the next SNIDs/NIDs to reach to the left out children. These may include, but not limited to, the following:***
  - *detailed microplanning for each activity;*
  - *use of updated map in microplanning and child-to-child search;*
  - *adequate orientation training of volunteers and workers;*
  - *adequate number of SNID/NID sites with required number of health workers and volunteers;*
  - *using masque miking, as well as Imam of the masque during Jumma Pray;*
  - *special team at railway station, bus stand, ferry-ghat etc. for traveling children*
  - *evening SNID/NID sites for working mothers; and*
  - *special mobile teams at night to vaccinate homeless and floating children.*
- ***Supervision of field workers during child-to-child search needs to be further strengthened to ensure that each and every household is visited and properly marked by the workers***
- ***More attention should be given to high risk areas and traveling/homeless/floating population***

1% of the eligible children (12 – 59 months) did not receive Vitamin A during the 1<sup>st</sup> round of the SNIDs- 2002. Besides, Vitamin A capsules were administered wrongly to 4 ineligible children (out of 210) who were <1 year of age on the day of 1<sup>st</sup> round SNID.

- ***Special attention should be given to:***
  - *ensure that each and every eligible child 12–59 months receives Vitamin A;*
  - *the exact age group to limit shortage (better screening for age); and*
  - *no Vitamin A capsules should be given to the parents to administer them to their children either at NID site or in their homes.*



## TABLES AND FIGURES


**Table 1: Routine immunization coverage levels of the children**

Name of the Vaccine	Coverage (%) Immunization of 12-23 months age group		Coverage (%) Immunized by 12 months of age
	Crude data (Access)	Valid data	Valid data
BCG	98	98	98
OPV1	98	94	94
OPV2	97	89	89
OPV3	95	86	86
DPT1	98	94	94
DPT2	97	89	89
DPT3	95	86	86
Measles	91	87	84
Fully immunized	91	81	78
Zero Dose	2	-	-

**Table 2: Routine immunization coverage levels by gender**

Name of the vaccine	Coverage % Immunization of 12-23 months age group				Coverage % Immunized by 12 months	
	Crude data (Access)		Valid data		Valid data	
	Male (%)	Female (%)	Male (%)	Female (%)	Male (%)	Female (%)
BCG	99	97	99	97	99	97
OPV1	99	97	99	89	99	89
OPV2	98	96	92	85	92	85
OPV3	95	94	90	82	90	82
DPT1	99	97	99	89	99	89
DPT2	98	96	92	85	92	85
DPT3	95	94	90	82	90	82
Measles	90	91	86	88	80	87
Fully immunized	90	91	83	79	76	79
Zero dose	1	3	-	-	-	-

**Figure 1: Drop-out rate for child immunization**



**Table 3: Invalid doses of immunization provided to the children**

Antigens	Percents
DPT1	4
DPT2	2
DPT3	1
Measles	3

**Table 4: Missed opportunities by antigens**


Name of the vaccine	Uncorrected		Corrected		Total		
	Number	Percent	Number	Percent	Number	Percent	
BCG	5	2	3	1	8	3	
DPT1	5	2	1	1	6	3	
DTP2	0	0	0	0	0	0	
DPT3	0	0	0	0	0	0	
OPV1	5	2	1	1	6	3	
OPV2	0	0	0	0	0	0	
OPV3	0	0	0	0	0	0	
Measles	1	1	4	2	5	3	
*Index						25	

*\* The idea is to propose one composite index reflecting the quality of screening during vaccination sessions.*


**Table 5: EPI cards availability and retention**

Card Status	Number	Percentage
EPI card available	125	60
EPI card ever given	206	98
EPI card retention	125	61

**Figure 2: Knowledge about required visits to immunization centers for full immunization**



**Figure 3: Sources of child immunization services**




**Table 6: Time required to reach the EPI center by mode of transportation**

Time required	Mode of transport					
	Walking		Rickshaw		Total	
	#	%	#	%	#	%
1-5 Minutes	97	47	7	3	104	50
6-10 Minutes	94	45	7	3	101	48
11-15 Minutes	5	2	-	-	5	2
Total	196	94	14	6	210	100

**Table 7: Reasons for non-immunization and partial immunization of the children**

Reasons for non-immunization or partial immunization	Non-immunized (%) (N=4)	Partially immunized (%) (N=16)
Did not know about need of immunization	50	-
Did not know about need of next dose	-	19
Did not know about importance of measles vaccine	-	19
Child was sick and not taken to immunization center	-	31
Painful for the children	50	19
Others	-	12


**Figure 4: Routine immunization coverage levels for TT among women 15-49 years**



**Table 8: Age distribution of women who never received TT vaccine**

Age group	Received		Not received		Total	
	#	%	#	%	#	%
15-19	17	61	11	39	28	100
20-25	62	94	4	6	66	100
26-30	54	95	3	5	57	100
31-35	37	93	3	7	40	100
36-45	12	63	7	37	19	100
Total	182	87	28	13	210	100

**Figure 5: TT Immunization drop-out rate among women 15-49 years**



**Table 9: Interval between TT1 and TT2, TT2 and TT3, TT3 and TT4, TT4 and TT5 doses**

Interval between dose	<1 months		1 months+		<6 months		6 months+		<1 year		1 year +		Total	
	#	%	#	%	#	%	#	%	#	%	#	%	#	%
TT1-TT2	4	3	141	97	-	-	-	-	-	-	-	-	145	100
TT2-TT3	-	-	-	-	19	19	79	81	-	-	-	-	98	100
TT3-TT4	-	-	-	-	-	-	-	-	49	72	19	28	68	100
TT4-TT5	-	-	-	-	-	-	-	-	16	47	18	53	34	100

**Table 10: Children born protected against tetanus**

Status of children born protected	Number	Percentage
Protected	205	98
Not Protected	5	2


**Table 11: Knowledge about number of TT doses required for life time protection against tetanus**

Answers	Number	Percentage
5 doses	22	10
Don't know/ no idea	188	90

**Table 12: TT cards availability and retention**

Card Status	Number	Percentage
TT card available	46	22
TT card ever given	181	86
TT card retention	46	25

**Figure 6: Providers of TT immunization**



**Table 13: Reasons for non-immunization and partial immunization for TT of the women**

<b>Reasons</b>	<b>Non-immunized (%) (N=28)</b>	<b>Partially immunized (%) (N=132)</b>
Next dose is not yet due	-	14
Don't feel need for immunization	32	-
Health worker did not specify the next dose	-	7
As per HW's advice, 2/3 TT is enough during the pregnancy	-	15
Unaware of need of next dose	-	60
In our times, TT immunization was not introduced	32	-
Fear of injection	36	4

**Table 14: OPV and Vitamin A Coverage during the SNIDs-2002  
N=210**

<b>Round</b>	<b>OPV (%)</b>	<b>Vitamin A (%)</b>
1 <sup>st</sup> round	99	99
2 <sup>nd</sup> round	99	-
Both rounds	98	-
Any round	99	-

**Table 15: Sources of OPV during the SNIDs-2002**

<b>Sources of OPV</b>	<b>1<sup>st</sup> Round</b>		<b>2<sup>nd</sup> Round</b>	
	<b>#</b>	<b>%</b>	<b>#</b>	<b>%</b>
NID site	208	99	207	99
Child to child search	-	-	-	-
Mobile Team	-	-	-	-
Not received	2	1	3	1
Total	210	100	210	100


**Table 16: Time required to reach the SNID sites by mode of transportation**

Time required	Mode of transport			
	Walking		Total	
	#	%	#	%
1-5 Min.	148	70	148	70
6-10 Min.	62	30	62	30
Total	210	100	210	100

**Table 17: Households visited during the child-to-child search of the 2<sup>nd</sup> round of the SNIDs-2002**

Variable	Number	Percents
Households visited	184	88
Households not visited	26	12
Total	210	100

**Figure 7: Sources of information about the SNID campaign**



**Table 18: Reasons for not receiving of OPV from SNID sites**

<b>Reasons</b>	<b>1<sup>st</sup> Round (%) (N=2)</b>	<b>2<sup>nd</sup> Round (%) (N=3)</b>
Did not know about NIDs	50	33
Forgot the date of SNID days	50	-
Parents were too busy	-	33
Child was away from home	-	34

**Annex-A****List of Selected Clusters for the Survey**

<b>Word no.</b>	<b>Mouza name</b>	<b>Mahalla name</b>	<b>Total HH</b>	<b>Total Pop.</b>	<b>Cluster No.</b>
1	Boitara	Boitara	248	1242	1
	Bhul Joyra	Bhul Joyra	85	557	2
	Gangadhar Patt	Gangadhar Patt	170	791	3
	Kandarkandi	Kandarkandi	69	376	4
	Nowakhanda (part-2)	Nowakhanda (part-2)	292	1608	5
	Nowakhanda (part-1)	Nowakhanda (part-1)	274	1445	6
	Narangi	Narangi	201	1027	7
	Nabagram	Nabagram	126	709	8
	Paschim Sheota	Paschim Sheota	316	1728	9
	Uttar Sheota	Uttar Sheota	448	2053	10,11
2	Char Hijuliy	Char Hijuliy	197	994	12
	Dhakua Para	Dhakua Para	144	714	13
	Dashora	Dashora	2505	12094	14-21
	Hijuliy	Hijuliy	238	1300	22
3	Andharmanik	Andharmanik	160	839	23
	Barashurundi	Barashurundi	329	1712	24
	Bewtha	Bewtha	123	612	25
	Joynagar	Joynagar	107	489	26
	Pouli (puratan)	Pouli (puratan)	380	1882	27,28
	Paschim Bandutia	Paschim Bandutia	707	3486	29,30

**Annex-B****List of Never Vaccinated Children Identified by Clusters**

<b>Word no.</b>	<b>Mouza name</b>	<b>Mahalla name</b>	<b>Total HH</b>	<b>Total Pop.</b>	<b>Cluster No.</b>	<b>Never Vaccinated children</b>
1	Nowakhanda (part-2)	Nowakhanda (part-2)	292	1608	5	1
	Nowakhanda (part-1)	Nowakhanda (part-1)	274	1445	6	1
	Nabagram	Nabagram	126	709	8	1
3	Pouli (puratan)	Pouli (puratan)	380	1882	27*, 28	1*

## **Acknowledgements**

### **Survey coordination:**

Mr. Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

### **Survey management:**

Mr. Jagadindra Majumder, Field Survey Manager, IOCH/MSH

### **Data analysis:**

Mr. Md. Mafizur Rahman, Monitoring and Evaluation Specialist, IOCH/MSH

Ms. Shahida Akhtar Ripa, PEF Monitoring, IOCH/MSH

### **Report writing:**

Mr. Md. Mafizur Rahman, Monitoring & Evaluation Specialist, IOCH/MSH

### **Report review:**

Dr. Pierre Claquin, Chief of Party, IOCH/MSH

### **Digital map preparation:**

Mr. Din Mohammed, Monitoring and Evaluation Assistant, IOCH/MSH

### **Cover photo:**

Ms. Moumina Dorgabekova/Image Jinn

### **Survey Team members, IOCH/MSH:**

Mr. Md. Abdul Hamid, Field Investigator

Mr. Md. Saiful Islam, Field Investigator

Ms. Krishna Rani Shil, Field Investigator

Ms. Khaleda Akhter, Field Investigator

Ms. Mahamuda Parveen, Field Investigator

Ms. Aung Ma Ching Marma, Field Investigator

Ms. Niva Rani Taju, Field Investigator