

Annual Report 1997



IITA


 The International Institute of Tropical Agriculture (IITA) was founded in 1967 as an international agricultural research institute with a mandate for major food crops, and with ecological and regional responsibilities to develop sustainable production systems in tropical Africa. It became the first African link in the worldwide network of agricultural research centers supported by the Consultative Group on International Agricultural Research (CGIAR), formed in 1971.

IITA is governed by an international board of trustees and is staffed by approximately 80 scientists and other professionals from over 30 countries, and approximately 1,300 support staff. A large proportion of the staff are located at the Ibadan campus, while others are at stations in other parts of Nigeria, and in Benin, Cameroon, and Uganda. Others are located at work sites in Côte d'Ivoire, Ghana, Malawi, Mozambique, Tanzania, Zambia, and Zimbabwe. Funding for IITA comes from the CGIAR and bilaterally from national and private donor agencies.

IITA conducts research, training, and germplasm and information exchange activities in partnership with regional bodies and national programs in many parts of sub-Saharan Africa. The research agenda addresses crop improvement, plant health, and resource and crop management within a farming systems framework. Research focuses on smallholder cropping systems in the humid and subhumid tropics of Africa and on the following major food crops: cassava, maize, plantain and banana, yam, cowpea, and soybean.

The goal of IITA's research and training mission is to "improve the nutritional status and well-being of low-income people of the humid and subhumid tropics of sub-Saharan Africa."

Global links Cosponsored by the World Bank, the Food and Agriculture Organization of the United Nations (FAO), the United Nations Development Programme (UNDP), and the United Nations Environment Programme (UNEP), the CGIAR is an informal association of over 40 governments and about 15 international organizations and private foundations. The CGIAR provides the main financial support for IITA and 15 other international centers around the world, whose collective goal is to improve food security, reduce poverty, and protect the environment in developing countries.



INTERNATIONAL INSTITUTE
OF
TROPICAL AGRICULTURE

**ANNUAL
REPORT
1997**

CONTENTS

1	From the Director General
4	Margaret Quin acclaims 1997 “A Year of Progress”
6	Maize wild relatives get a stranglehold on <i>Striga</i>
8	IITA spurs doubling of cassava use in Nigeria
10	Optimizing crop rotation sustains productivity in moist savanna
12	Landraces winning the race against cassava mosaic disease
16	Common goals drive RCMD’s research agenda
18	Restoring nutrients to sub Saharan soils
20	Alley farming’s future brightens
22	IITA and ILRI collaborating for future generations
24	Simultaneous cropping with <i>Pueraria</i>
26	Farmers’ perceptions of soil degradation
28	Mimicking the forest multistrata systems for the humid tropics
32	Peter Neuenschwander finds that success breeds success
34	Integrated strategies to reverse plantain losses in Ghana
36	IPM helps farmers increase cowpea production
38	IITA’s insect pathology program applying LUBILOSA’s lessons
40	BioNet International and IITA’s insect taxonomy role
42	Partnership for food the Technology Testing & Transfer Unit
47	Michael Bassey praises training in technology transfer
48	El Niño, NARS, and IITA news on the Web
51	IITA research highlights
71	Graduate research completed at IITA in 1997
73	Financial statement
74	Statement of activity
75	Statement of cash flow
75	Donors 1997
76	Publications by IITA staff
91	Board of trustees
92	Professional staff
93	Abbreviations and acronyms used in the text

From the Director General

For IITA 1997 was a year of extremes. The Institute's excellent research is benefiting more farmers in Africa, who, with improved technologies, get more income through production systems that minimize environmental impact.

At the same time, the Institute had to face potentially serious financial shortfalls: there were continuous cost increases and a reduction in unrestricted core funding. Notwithstanding these difficulties, we have decided that, for the time being, the Institute will maintain the existing broad crop portfolio and decentralized research structure. This approach is needed so that we can effectively address the research needs of our ecoregion.


In particular, our Humid Forest Station is carrying out extensive research in the humid forest benchmark area in Cameroon. This serves both the Systemwide Alternatives to Slash-and-Burn Program and the Ecoregional Program for the Humid and Subhumid Tropics of sub-Saharan Africa (EPHTA).

The station in Namulonge, Uganda is working closely with the African Highlands Initiative, while our station in Kano, Nigeria links into the moist savanna benchmark area of EPHTA and serves also for joint IITA/ILRI/ICRISAT research activities. Support to the benchmark area in Benin is provided from our station at Cotonou.


This year was also marked by a Center Commissioned External Review (CCER) of our cereal/legume projects. The CCER team of five experts was complemented by a national agricultural research and extension systems team. Extensive field visits and discussions allowed for an excellent analysis of the work undertaken. The team highlighted the progress made and, particularly, the successful development of a systems approach to crop improvement, plant health and production problems. The review noted that for the three main projects excellent progress is being made in reaching the objectives. It also made constructive recommendations on further integration of biotechnology and plant genetic resource research.

Towards the end of the year, we had a two-day planning meeting with the Conference des responsables de la recherche agronomique africains (CORAF). This allowed us to streamline our respective activities and to avoid duplication. A memorandum of understanding was developed, including a strategy adopted to integrate the various research networks in West and Central Africa.

IITA existed 30 years in 1997. Given the overall financial situation, the festivities were limited to social activities for all the staff. At 30, IITA has become somewhat leaner than it was, for example, 10 years ago, but it is also more experienced and has learned to do more with less. Thus, the Institute is in good health, and I am confident that it will continue to evolve and produce excellent results in a changing environment.



Dr. Lukas Brauer



IIITA works across three agroecological zones of sub-Saharan Africa: the savannas, the humid forests, and the mid-altitude regions. Each zone poses unique challenges to food production, which require multidisciplinary technology development.





Margaret Quin Acclaims 1997 "A Year Of Progress"

The Crop Improvement Division (CID), headed by Margaret Quin, continued its well-focused research agenda in 1997

During the late 1980s IITA developed maize that yielded well in *Striga*-infested fields. Since 1993 the Institute has focused on reducing *Striga* emergence.

"The aim was to cause *Striga* to be less able to develop on maize. This isn't easy, but we stuck to it, and by 1997 we had really made progress. This has significantly contributed to maize's sustainability potential," Quin explained.

Adapting cassava to dry areas progressed. IITA now has cassava green mite-resistant varieties which stay green and retain a good leaf canopy during the dry season.

Adapting soybean to African soil ranked high in 1997. "There was definite progress in selection for acceptable grain and fodder yields using no phosphorus fertilizer. This built on IITA's nitrogen fixation record. Because of IITA, soybean is becoming farmer friendly, especially when considering crop rotations," Quin said.

In 1997 IITA's yam program expanded into Tanzania and Uganda, where many of the growers are women. Farmer participatory evaluation of improved yam germplasm is ongoing.

"Last but not least, cowpea transformation with candidate genes for insect resistance was achieved," she said.



"In 1994 IITA won the King Baudouin Award for producing black sigatoka resistant plantain hybrids, but they lacked qualities that consumers like. We now have hybrids with high yields, larger fruit, and better pulp color that resemble, more closely, preferred plantain. It's a major advance with high potential."



Maize wild relatives get a stranglehold on *Striga*

Striga species, a flowering plant indigenous to Africa, has a terrible reputation. The Food and Agriculture Organization of the United Nations estimates that the parasite causes annual losses in excess of US\$7 billion, adversely affecting over 100 million African people in West Africa alone.

The Striga problem in Africa is intimately associated with intensification of land use.

It is no wonder that *Striga* is a major threat to agriculture. One plant can produce 50,000 to 500,000 seeds, which remain viable for up to 14 years in the soil. These seeds are virtually microscopic. After germination, the seedlings can grow three to seven days in the absence of a host. If they attach to a host, such as the root of a maize plant, the parasite grows underground for up to seven weeks prior to breaking through the soil. The scenario fits a science-fiction thriller, except this is for real.

Most of the damage to the unfortunate host plant occurs underground, making it difficult for farmers to determine the actual cause of the damage. Was it a pest or a disease? Symptoms of parasitism resemble drought stress accompanied by severe plant stunting, and they can all add up to 100% yield loss.

Resembling a battlefield the struggle between *Striga* and a maize crop shows the victor *Striga*

Elite tropical maize has low resistance to *Striga*. A maize wild relative though *Zea diploperennis* is highly resistant. IITA scientists backcrossed the resistant genes thus incorporating them into adapted maize genotypes.



The *Striga* problem in Africa is intimately associated with intensification of land use. Traditional African cropping systems have included prolonged fallow, rotations, and intercropping, which kept infestations at tolerable levels. However, as population pressures increase and markets develop, demand for food production increases and land use intensifies.

Integrated control

Components of successful control are to prevent the influx of *S. hermonthica* seeds into fields and reduce their density in the soil, reduce parasitism, and prevent parasite reproduction. Perhaps easier

said than done, one might think, but IITA has developed an integrated control strategy that starts with the use of parasite-free planting material, and then focuses on crop rotation and nitrogen-fixing legume cultivars that have been selected for efficacy in reducing the density of *Striga* seeds in the soil. In short, the legume, such as soybean, causes the microscopic seeds to abort in their development. This focus is reinforced by the use of managed biological control and high-yielding adapted maize with resistance to *Striga*.


This last element has proven difficult to achieve in the past because elite tropical maize has a low resistance to *Striga*. This caused scientists to begin looking for wild relatives of maize that were resistant to *Striga*. They found it in *Zea diploperennis*, which has high resistance to the parasite. Through a process of backcrossing, the resistant genes were incorporated into adapted maize genotypes.

Unfortunately, *Striga* resistance is a quantitatively inherited trait, involving many genes, and it is difficult to transfer this trait efficiently to locally adapted cultivars through backcrossing.

To accelerate progress in incorporating the resistance genes from *Z. diploperennis* into adapted maize, a marker-assisted selection project was developed in 1997 by IITA, the Centro Internacional de Mejoramiento de Maize y Trigo (CIMMYT), Mexico, the Kenya Agricultural Research Institute (KARI), and the University of Hohenheim. This project is funded by the Rockefeller Foundation.

Through the use of molecular markers to identify resistant backcross material at the gene level, scientists are optimistic that they are on their way to getting highly resistant, adapted maize into an integrated control program. Such a program would provide a major advance in the control of *Striga*.

Pink *Striga hermonthica* carpets a farmer's field. But the beauty of the flowering plant belies its devastating effects on maize, a staple in sub-Saharan Africa. The weed parasitizes maize roots, thereby injuring them, leading to extensive yield loss.



IITA spurs doubling of cassava use in Nigeria


Per capita food production in Africa has been declining partly due to rapid population growth, soil degradation, periodic droughts, and a weak technological base. Despite this, Nigeria, the most populous country on the continent, produces more cassava than any country in the world.

Increase in cassava production will benefit farmers only if the demand is maintained or increased.

Per capita cassava production in Nigeria has increased from 130 kg in 1980 to 260 kg in 1994. This rise has been credited to the adoption of improved, high-yielding varieties by cassava farmers. These varieties, many

attributable to IITA, are mostly adapted to the humid forest and moist savanna zones of Nigeria. IITA continues to develop germplasm with drought tolerance, good leaf retention in order to provide fodder for livestock, and low cyanide content requiring minimal processing.

But the fact remains, increase in cassava production will benefit farmers only if the demand is maintained or increased. In order for this to happen, cassava growers and product-makers must take advantage of existing markets or create new ones. Only when cassava is linked to these markets will increased production translate into increased income and better welfare for the people.



Workers at a cassava processing plant near their farms in Ogun State, Nigeria, spread cassava grits in the sun to evaporate moisture in preparation for milling it into flour.



High quality cassava flour produced in a village packed in 50 kg bags is loaded for transport to a factory where it will be made into bakery products. Because of cassava flour's long shelf life as opposed to fresh roots (two to three days) it can be stored for up to six months.

Creating new markets

IITA is addressing this by developing technologies to improve cassava processing and utilization. These technologies have made it possible to expand cassava processing in rural areas and produce products with a high market value for urban users. These technologies and processes have been introduced to women's groups, small-scale cassava farmers, and processors.

Industrial users of cassava flour are substituting between 10% and 50% cassava for wheat flour in the making of products such as biscuits and noodles. One factory has switched from the use of sugarcane molasses to cassava flour for the production of ethanol. This reduces production costs.

Home caterers use cassava flour to make confectioneries such as cakes and doughnuts. Some home caterers use up to 100% cassava flour for making traditional baked products.

Between October 1995 and September 1996, three biscuit and one noodle manufacturer surveyed used an average of 60 metric tons of cassava flour and 612 metric tons of wheat flour monthly. The cassava flour cost is half that of wheat flour, which is often imported, expending foreign exchange.

IITA has determined the functionality of cassava flour and starch across a range of cassava varieties. One practical outcome of this research is that a number of recipes have been developed by IITA to demonstrate the flexibility of cassava flour in baking. These recipes are starting points for caterers and home bakers to develop products of their own.

Still, there are problems in the cassava flour production business. An IITA survey in several Nigerian states showed the top five problems to be the high cost of cassava, the lack of capital, drying in the rainy season, shortage of labor, and an inadequate supply of fresh roots.

The trend, though, is toward more use. IITA continues to develop the cultivars and technologies needed to make cassava production and processing vital parts of the economies of Nigeria and other countries in sub-Saharan Africa.

Optimizing crop rotation sustains productivity in moist savanna

The moist savannas of West and Central Africa have high potential for crop and livestock production. The region may be the emerging breadbasket of sub-Saharan Africa.

If the aim is to reduce dependence on fertilizer, the use of soybean in management systems shows potential.

Savanna soils are generally low in organic matter and nitrogen, therefore nitrogen deficiencies in maize are common. Genotypes with improved nitrogen-use efficiency can increase productivity of maize-based cropping systems when they are combined with improved soil fertility, crop rotation, organic manure, and inorganic fertilizers.

IITA scientists had to develop low nitrogen-tolerant maize that performed at least as well as or better than available varieties when grown under higher level of nitrogen.

Selected families (34) from the low N-tolerant pool, grown with no nitrogen inputs, had a mean grain yield of 3.2 t/ha, compared with 2.4 t/ha for the commercial hybrid.

The role of soybean

Improved soybean varieties can also increase the productivity and sustainability of maize-grain legume cropping systems because grain legumes can reduce reliance of maize on nitrogen fertilizer.

The new traits developed maximize nitrogen derived from the atmosphere (Ndfa) and optimize fodder production. By combining the traits, soybean varieties were developed that have a high and stable grain yield which maximize nitrogen and organic matter contribution to the soil and benefit the subsequent crop.


The other new trait increases *Striga hermonthica* germination, reducing the parasite's seed bank for the next maize crop.

Trials compared the performance of the six best soybean breeding lines with that of a check variety. Tgx 1829-1E had a Ndfa of 52.5 compared to the check variety's 36. Variety Tgx 1892-1E gave vastly improved yields over the check variety: 1809 kg/ha vs 1193 kg/ha for grain, and 2312 kg/ha vs 1750 kg/ha for fodder.

The improvement in both grain and fodder yield indicates that this new variety will give more soybean grain yield (thus quick income and greater nutrition) and show more return of organic matter and nitrogen to the soil through fodder.

The sun provides energy for plant growth and it is also used after harvest to evaporate moisture from cowpea pods allowing them to be stored.





The bottom line

Statistics show that different soybean residue management techniques and varieties can improve subsequent maize grain yield, although they are not as effective as small amounts of nitrogen fertilizer. For example, with no fertilizer, a maize crop following previous maize yielded 686 kg/ha, maize crops planted after five different soybean varieties yielded 1174, 1328, 1330, 1232, and 1381 kg/ha, and maize crops with 20, 40, and 60 kg/ha added nitrogen yielded 1937, 2070, and 2203 kg/ha, respectively.

Maize grown after soybean had significantly higher shoot dry weight and grain yield than that growing in former maize plots. This depended on the soybean cultivars.

Compared to maize, maize grain yield increase following soybean was variable, but the main effect of previous soybean crops was positive, even though the soybean was not inoculated and the above-ground soybean residues, except harvest litter, were removed from the fields.

In trials conducted in farmers' fields, the yield increase following the medium-duration soybean variety was similar to that from 40 kg of nitrogen per ha as urea applied four weeks after planting to maize preceded by maize. Thus, if the aim is to reduce dependence on fertilizer, the use of soybean in management systems shows potential.

Cowpea is another source of soil nitrogen. There are substantial increases in maize yield if it is grown immediately after cowpea, as the nutrients in cowpea roots, litter, and vines become available during the subsequent season.

The benefits of double cropping cowpea with early maize accumulate over time, but soil phosphorus must be adequate for the legume to have an effect.

As is common in sub-Saharan Africa, animal traction provides motive power in agriculture. In Kaduna, Nigeria, a farmer uses cattle in plowing to control weeds in his soybean field (above). Soybean seeds are separated from pods by threshing. Soybean and soymilk are rapidly becoming major food staples as well as industrial commodities in West Africa largely as a result of IITA soybean varieties (below).



Landraces winning the race against cassava mosaic disease

The most widespread and economically damaging crop disease in Africa is cassava mosaic disease (CMD). The virus causing this disease is spread by the whitefly. It costs the continent an estimated US\$2 billion a year in crop losses. IITA is responding.

Viruses are insidious. In order to infect either plants or people, they have to reproduce themselves, which they can't do on their own. Molecularly speaking, they have figured out how to replicate themselves, thereby causing chaos in cells that are not resistant to them.

Viruses commandeer a host's DNA, which is the molecular basis of heredity of both people and plants, and use the DNA's reproductive capacity to produce an assortment of systemic diseases.

IITA helps its partners

Today, thanks to more than a quarter century of research by IITA scientists, farmers throughout Africa are planting new cassava varieties with

Sustainable multiplication and distribution of certified cassava planting material from improved genotypes is fundamental to the acceleration of impact at the farm level.



better resistance to highly destructive viruses. Over the years, this was accomplished by deploying a source of resistance that was derived from an interspecific cross between cassava and a wild relative.

More recently, African cassava landraces have also contributed new genes for resistance to the disease. For the past seven years, IITA scientists and national program partners from Benin, Ghana, Nigeria, and Togo have collected resistant landraces in West Africa to further diversify and expand the gene pool of African-adapted germplasm. These were incorporated into IITA's breeding populations, which produced offspring with heightened resistance to CMD and some other diseases. In addition, several CMD-resistant landraces were also resistant to the cassava green mite.

Cassava farmers in Africa, using these improved materials, could obtain yields up to five times those of many local susceptible cultivars when diseases are severe. However, sustainable multiplication and distribution of certified cassava planting material from improved genotypes is fundamental to the acceleration of impact at the farm level.

To facilitate this, IITA is distributing improved genotypes to national partners for testing under local conditions. In addition, the Institute has engaged in a project with Centro Internacional de Agricultura Tropical (CIAT), Colombia, to map the genes conferring CMD resistance, and to develop molecular markers for the trait.

The outcome will assist the incorporation and enhancement of resistance to the disease in African, Central and South American, and Asian cassava gene pools. It will also provide a means, through indirect selection, for developing CMD-resistant germplasm, adapted to Latin America and Asia, which could be deployed should the virus and vector come together on these continents.



A young woman clutching freshly harvested cassava leaves with ease and grace balances a yam on her head. Cassava leaves contain protein, vitamins, and minerals (above). Making garri, done almost exclusively by women in Nigeria, involves peeling and grating storage roots, then fermenting the mash, pressing to remove liquid, sieving, and then frying (below).



In Africa,
human beings
think holistically
among all things







Common Goals Drive RCMD's Research Agenda

The Resource and Crop Management Division, headed by Horst Grimme, has four projects under its research umbrella: Short Fallow Stabilization, Agroecosystem Development Strategies, Improving Postharvest Systems, and Farming Systems Diversification.


The Short Fallow Stabilization project's humid forest component is concentrated in areas where population growth and intensified land use have forced farmers to reduce fallow periods.

"Soil fertility must be regenerated. But any fallow technology to regenerate fertility has to be well adapted and acceptable to farmers. Growing herbaceous legumes shows a lot of promise in this regard," Grimme explained.

The Agroecosystem Development Strategies project works closely with EPHTA, the Ecoregional Program for the Humid and Subhumid Tropics of sub-Saharan Africa. Like EPHTA, it assists farmers to increase productivity and food security through sustainable production technologies.

"Our postharvest work also increases food security by enhancing crop handling, product development, and reducing loss. Some 60 postharvest devices have been engineered by the unit," Grimme added.

Farmers are much the focus of the Resource and Crop Management Division. "Our projects are aimed at increasing smallholder productivity and cash income. To get farmer acceptance, though, the farmer has to be involved from the beginning," he said.







Intercropped sorghum and millet struggle to survive in nutrient depleted soil reflecting the breakdown of nature's self-sustaining system. This is compounded by drought and diseases.

Restoring nutrients to sub-Saharan soils

Land-use intensification can be sustained only if the nutrients depleted during cultivation are replenished

Population growth and the rising aspirations for the well-being of rural populations lead to clearance of large areas of forest, an extension of the cropping phase, a shortening of the fallow period, and subsequent soil fertility decline. This nutrient mining results in the breakdown of nature's self-sustaining system. Beside nutrient mining, topsoil removal, due to water and wind erosion, is one of the greatest threats to sustainable agricultural production in sub-Saharan Africa.

ITA scientists are applying biological processes to restore soil fertility and protect the soil by adapting germplasm to adverse soil conditions, enhancing soil biological activity, and optimizing nutrient cycling to minimize external inputs and maximize efficiency of their use.

In this paradigm, soil organic matter (SOM) plays a substantial role in maintaining soil fertility by providing nutrients through decomposition, retaining moisture, helping maintain soil structure, buffering available

nutrients, and binding toxic substances. Crop residues play an important role in maintaining soil fertility because they provide nutrients through decomposition.

However, soil organic inputs alone cannot sustain agricultural production. Sustainable crop production in the majority of soils in West Africa requires the continuous addition of organic materials, minimal soil disturbance, judicious fertilizer and lime use, multiple cropping that includes perennial crops, fallowing, and rotation of crops.

Nutrient management systems

Related to the use of soil organic matter to restore fertility is another factor which balances nutrients

The bottom line is to maintain improved soil nutrient balance by promoting utilization of locally available sources of plant nutrients.

Population growth in Africa has precipitated the clearance of great areas that were formerly forest and balanced ecosystems. The challenge is to preserve this land against nutrient depletion and to make it productive for food crops and raising livestock.



Called balanced nutrient management systems (BNMS), it has proven potential to raise cereal yields in sub-Saharan Africa five times. At present, yields are on average less than one tonne per ha, about one-fifth what could be obtained with good plant nutrient management.

With the prevailing farming systems, though, this average yield is bound to drop to even lower levels because of the irreversible mining effect of plant nutrients from the stressed soils. This critical, vicious cycle can no longer continue. That's why IITA has launched an integrated approach to overcome the food supply shortfall in West Africa. The research is funded by the Belgian government.

The BNMS for maize-based systems in the moist and humid

forest of West Africa builds on the achievements of previous SOM projects. However, it looks at the problem from a much wider angle, encompassing all major nutrients. Inorganic inputs may be appropriate. The farmer's perspective is considered, too.

The goal is to curb the various cycles of plant nutrient depletion in maize-based farming through integrated nutrient management geared to land-use practices which are economically viable, ecologically sound, and socially acceptable. The bottom line is to maintain improved soil nutrient balance by promoting utilization of locally available sources of plant nutrients, maximizing their nutrient-use efficiency and thus reducing the need for external, costly soluble fertilizers.

A research project to test this concept will increase production and quality of farm products, diversify land use, increase farm incomes, generate employment, and protect land resources and the natural environment.



Crop residues can help restore soil fertility and protect soil. The loss of soil nutrients and erosion are great threats to sustainable agricultural production in sub-Saharan Africa.

Alley farming's future brightens

IITA has developed and is promoting alley farming as a production alternative to slash-and-burn shifting cultivation

The system involves the cultivation of food crops between hedgerows (alleys) of multipurpose trees or shrubs, usually legumes. The legumes have deep roots that help in fixing nitrogen and nutrient cycling, their nitrogen-rich leaves can also be used as green manure and as fodder for livestock. By reducing the amount of land needed to be cleared for farming, alley farming saves natural vegetation.

With these acknowledged advantages, much enthusiasm was generated about the technology. However, the lack of immediate success, measured in terms of widespread farmers' adoption, led to skepticism about its relevance and adoptability. To assess the adoption of the technology and farmers' current management practices, IITA conducted surveys in Benin, Cameroon, and Nigeria, some 10 years after the introduction of this new technology.

Evidence from the studies shows that earlier skepticism about the adoption potential of the technology appears unjustified. Farmers are adopting the technology in villages characterized by (a) high land-use pressure, (b) soil fertility decline, (c) erosion problems, and (d) firewood and fodder scarcity. Farm-level adoption patterns vary across the three case-study countries.

In Nigeria, of the 223 farmers surveyed, 208 (93%) had heard about the technology, of which 66% had either initially adopted or experimented with it. Some of the initial adopters abandoned it, but 53% of these farmers continue to use it.

In Cameroon, of the 820 farmers surveyed, 256 (31%) had either initially experimented or adopted the technology. Of this group, 238 farmers (93%) continue to use it.

In Benin, of the 288 farmers surveyed, 225 (78%) had heard about the technology. Of the latter, only 72 farmers (32%) had either experimented or initially adopted it. Current level of use among this group is encouraging, as 93% continue to use the technology.

Solving the constraints to alley farming technology requires that researchers focus on technology modifications that will make the technology more flexible and adaptable to farmers' preferences

A farmer uses his machete to cut hedgerow legumes for use as green manure. The prunings when strewn on the soil also reduce runoff and erosion.





As farmers look for a better match of the technology with their resources and preferences, they have made important modifications to alley farming. The most significant is the use of fallow phases. For example, in Nigeria, of the sample farmers currently using the technology, 93% had made changes to the technology, of which 83% added a fallow phase.

Reasons for not adopting the technology now are traceable mainly to technology-related constraints. These include high labor demands, tree-crop competition, lack of knowledge about management, and hard-to-get leguminous trees and shrubs.


Land tenure, often believed to constrain adoption, wasn't a major problem in the three countries and appears to be overemphasized. It is thus highly unlikely that changes in land tenure to increase rights under private ownership of land will significantly change current adoption rates.

Also, if alley farming is not being adopted by some farmers, it is not due to lack of secure property rights alone. Farmers can be encouraged to take advantage of alley farming. If alley farming or other agroforestry technologies provide significant income, farmers, as rational people, will reallocate land under their ownership to take advantage of the economic benefits. But they must be convinced of the benefits.

Solving the constraints to alley farming technology requires that researchers focus on technology modifications that will make the technology more flexible and adaptable to farmers' preferences.

As researchers recognize and integrate these farmer modifications in future technology designs, adoption should increase. Such modified systems would need to be targeted to areas where incentives for land-use changes, i.e., land-use pressure, soil fertility decline, erosion, and fuelwood scarcity exist or are likely to occur in the near future.

In Africa, trees are being used more and more for staking material, firewood, and fodder. Here an enterprising farmer aloft gives his livestock assistance by cutting off branches (left). Cowpea bountifully growing between a nitrogen-fixing herbaceous legume is tended by a farmer. Alley farming is seen by more farmers in Africa as an alternative to slash and burn cultivation (above). Alley farming is an integrated system that sustains fertility of farmlands by imitating the self-renewing natural process of the forest. It also reduces the amount of land needed to be cleared for farming, thus conserving natural vegetation (below).



IITA and ILRI: collaborating for future generations

IITA is collaborating with the International Livestock Research Institute (ILRI), Ethiopia, to develop ways in which crop-livestock systems can function synergistically, without damaging the environment

Cattle grazing natural pastures supplemented with legumes gained up to 140 g a day during the dry season.

This joint-center effort has arisen because of increasing demands for crop and livestock products in sub-Saharan Africa. Human population there will top one billion within 20 years. Feeding so many will place great demands on agriculture.

Unfortunately, growing more crops, which requires more land, results in less fallow land. This uncropped land, though, allows the soil to recover its fertility and is commonly grazed by livestock. Such livestock vs. crop competition has the potential to severely degrade the natural resource base, which could be disastrous.

ILRI and IITA scientists believe they have found a partial solution to this situation: herbaceous legumes. The use of legumes is not new, of course. What is new is their multiple use, amidst changing conditions, in maintaining soil fertility, controlling weeds, pests, and erosion, and in providing feed for livestock.

For example, it's been found that species of *Stylosanthes*, *Centrosema*, and *Aeschynomene* can produce substantial quantities of nutritious fodder for livestock. This is especially so during the dry season, when the naturally available feed resources are of very



Crop legumes for better feed for livestock. The use of legumes is not new, of course. What is new is their multiple use, amidst changing conditions, in maintaining soil fertility, controlling weeds, pests, and erosion, and in providing feed for livestock.



Cowpea fodder is transported for storage. It will be ready for use during the dry season.

poor quality. At the same time, these legumes also enrich the soil to benefit future crops, and some species can contribute to weed control or *Stiga* reduction.

Legume options

In sub-Saharan Africa, appropriately managed forage legume pastures, as demonstrated in on-station trials, can support up to eight times as many cattle as native pasture and can more than double subsequent crop yields. Trials have shown that, with maize, crop yields following forage legumes were equivalent to those obtained by applying 90 kg of nitrogen fertilizer per hectare to natural fallow plots.

Also, cattle grazing natural pastures supplemented with legumes gained up to 140 g a day during the dry season. Those grazing only natural pasture lost 58 g a day. Such differences in weight gain, which can amount to as much as 30 kg in one dry season, have important, positive implications for the survival and productivity of the animals.

The researchers believe that by testing a number of herbaceous legume species, they can present options to farmers, so that they can choose what suits them best, for example, options for crop yield, weed

control, *Stiga* reduction, or livestock nutrition, etc.

Another major aspect of sustainable crop-livestock integration on which scientists from the two institutes are collaborating is concerned with the quantity and quality of residues from crops such as cowpea, maize, soybean, and cassava, which can be produced without detracting from grain or tuber yield.

After grain harvest, the remainder of crops (such as leaves, pods, and stems) can be used as a source of feed for livestock. Cowpea residue is already recognized by many farmers as a valuable feed, scientists from both institutes have identified dual-purpose varieties, thereby providing the maximum returns for farmers' investments.

Biotechnology is being considered as a tool to further enhance breeding and selection to improve crop residue quality.

These options provide opportunities to improve natural resource management through integrated crop-livestock systems. For example, animal manure contributes to soil fertility, an important attribute in view of increasing shortages of inorganic fertilizers.

Crop and livestock integration will impact on the welfare of future generations and the preservation of the environment. IITA and ILRI are pioneering ways to ensure that such integration has a positive role in sustainable agriculture.



Pueraria cover crop was simultaneously established with cassava intercropped with maize (photo taken six months after sowing)

Simultaneous cropping with *Pueraria*

After experimenting with many legume cover crop species for soil fertility improvement, IITA soil scientists have developed a soil fertility improvement technology without fertilizing simultaneous cropping with Pueraria phaseoloides

Over the years, IITA has increased productivity and enhanced sustainability of agroecosystems, partially through research on land and soil management. From the results of two decades of research on tropical soils and the ecological and social setting in which farmers cultivate their crops, IITA soil scientists have concluded that the essential principle in preventing or retarding soil degradation is to integrate appropriate fallow vegetation, such as leguminous cover crops, with food crops.

Pueraria seeds are planted at the same time as the maize/cassava intercrop. Later the *Pueraria* is cut down to a quarter of a meter in height. The second year involves an entire fallow of *Pueraria* regrowth. Before the next cropping season it is suggested that *Pueraria* be incorporated, though burning may be favored by the farmer's tradition.

Pueraria as a herbaceous legume cover crop improves the fertility of the soil through nitrogen fixation as well as nutrient cycling from the subsoil and stimulation of activity of soil biota

Pueraria as a herbaceous legume cover crop improves the fertility of the soil through nitrogen fixation as well as nutrient cycling from the subsoil and stimulation of activity of soil biota

The legume helps control soil erosion and it also reduces weeds. It is well suited to West and Central Africa's bimodal rainfall regions. It grows slowly initially, but withstands drought and thrives where the moisture-holding level of the soil is high. Because of the slow growth, *Pueraria* does not compete with associated food crops during the establishment period. And because of its ability to survive in the dry season, *Pueraria* can thrive at the onset of the rainy season, overgrow weeds, and accumulate a large amount of shoot and root biomass with no extra labor during the subsequent year.

Comparing systems

According to a long-term trial without the use of fertilizers and pesticides in Ibadan, Nigeria, the maize grain yield in the *Pueraria* simultaneous cropping was maintained at 2.5 t/ha after the third cycle, whereas in the sole maize/cassava continuous

cropping and cropping with natural regrowth there was a rapid decline to below 0.4 and 1.5 t/ha.

There was a higher cassava tuber yield with the *Pueraria* simultaneous cropping (14.2 t/ha) than in the natural regrowth cropping (10.1 t/ha) and sole continuous cropping (6.6 t/ha) after the third cycle.

Importantly, the *Pueraria* simultaneous cropping not only increased crop production, but also maintained soil fertility. With the *Pueraria* simultaneous cropping the soil organic carbon and mineral nitrogen stabilized very close to that of the secondary forest. The nutrient budget analysis indicated a positive balance for nitrogen in the *Pueraria* simultaneous cropping if the *Pueraria* residue is incorporated before cropping maize/cassava, assuming maize grain and cassava root yield will be sustained at about 2.5 and 15 t/ha, respectively.

The soil-available phosphorus was still not depleted after seven years of cropping because the phosphorus taken up from the subsoil by *Pueraria* can compensate for that removed by maize grain and cassava roots.

There was no decrease in soil pH under *Pueraria* simultaneous cropping. The number of earthworms with the *Pueraria* simultaneous cropping, an important soil-quality measure, conformed to that of the secondary forest.

This research implies that a self-sustainable food production is possible using the *Pueraria* simultaneous cover cropping. The technology is ready for off-station trials in West and Central Africa.

During the fallow stage *Pueraria* adds nitrogen to the soil thus restoring its fertility.



Farmers' perceptions of soil degradation


The nature of farming is such that farmers are highly aware of the environmental conditions in which growing crops and tending livestock occur. This makes sense, because food and income losses can occur from their not being aware of constraints and failing to take action when possible.

A study was conducted by IITA, in collaboration with Nigeria's Institute of Agricultural Research, to determine just what soil conservation issues in particular are continually assessed by farmers. It was done in the northern Guinea savanna of Nigeria. A total of 181 farmers in 15 villages, including village chiefs, were interviewed. The interviews lasted less than half an hour. The questions were open-ended and conversational in nature.

Chiefs were asked what the limitations were for crop production. In decreasing importance, the answers were fertilizer availability and cost (with cost less important), insect infestation, drought, lack of access to roads, nonavailability of machinery, and weed infestation.

More than a quarter said that unproductive fields exist in the village, but they did not attribute

Farmers were asked what they perceived as farming practices that made them better farmers than others. The most common responses were management of resources, good planning, and hard work.



In Katsina, Nigeria, soil conservation is a major cause of erosion and it is on the increase in the Sahelian Africa.

their existence to farming practices. Only half of them thought that sufficient good land would be available in the future. Slash-and-burn cultivation was practiced in the villages by 87% of the chiefs. The procedure used was to gather the trash or crop residue and set it on fire. This released nutrients and controlled weeds and insects.

Nearly 90% of the chiefs were aware of gullies, visible signs of erosion, and soil degradation. Only about 20% said they did anything about them.

Livestock and crop-livestock interactions were also assessed. Livestock was perceived mainly as a source of food and income by chiefs. They were also ranked high for their social value during traditional ceremonies, followed by their use for traction and as suppliers of manure. Rarely were livestock perceived as potentially harmful to soil, especially during the long, dry season when no cover is left to protect surface soil.

Need for fertilizer

For farmers, though, the causes for poor yield can be categorized as those that are most important at present, and those that are expected to increase or decrease with time. Currently, lack of fertilizer and *Striga* were perceived as the main causes for poor yields by 88% of farmers. Lack of fertilizer was the only cause for poor yield which farmers believe will grow worse.

Insect problems, other pests, and diseases were perceived as lessening. Except for *Striga* and climate conditions, most causes given for poor yield were related to soils with runoff, i.e., inherently poor soil fertility, erosion, waterlogging, and loss of soil fertility.

Farmers were asked what they would like to have in order to improve their farming. Fertilizer was number one. The use of tractors and the availability of bank loans were second and third on the farmers' wish list. Thus, other soil-related causes were viewed as minor problems, possibly due to the current lack of fertilizer. This corresponds to scientific evidence that no conservation practices can completely replace fertilizer for restoring fertility in the long run.

Moreover, crop rotation and adequate crop establishment were perceived as necessary for adequate yield. For instance, farmers consistently planted on ridges to facilitate crop establishment.

Farmers were asked what they perceived as farming practices that made them better farmers than others. The most common responses were management of resources, good planning, and hard work.



In Benin a village agricultural agent himself a farmer stands in his cotton field. He uses the teaching guide he holds to help his farmer neighbors solve crop production problems.

Mimicking the forest: multistrata systems for the humid tropics

IITA, in collaboration with farmers and national and international research institutes, is developing systems that can increase the productivity and income of smallholders

In the humid tropics, the development of multistrata systems is a priority. These systems, which integrate different annual and perennial crops, mimic the natural forest structure where vegetation forms several strata

In the humid tropics, the development of multistrata systems is a priority.

Although there is high demand for timber, fruits, nuts, and medicines, farmers have been reluctant to invest land and labor in new tree-based systems where the first harvest is often years away. Multistrata systems, therefore, are being developed which include annual food crops to provide income during the start-up years before the tree crops are harvested, they also prevent the bush from invading the young tree stand

In southern Cameroon, cocoa farmers are integrating upper canopy fruit trees such as *Dacryodes edulis*, the African prune, into their plantations and wish to add selected food crops as a lower stratum. Others are interested in growing oil palm and fruit trees with under-storey food crops. Forty such fields in five forest margin benchmark villages have been planted by farmers in partnership with researchers from IITA's Humid Forest Station and the Cameroonian Institute of Agricultural Research and Development. The effect and preference of frequency of annual cropping, introduction of nitrogen-fixing cover crops and hedgerows, and the use of chemical fertilizers are being addressed

Aware of the predicted timber shortage from Cameroon's permanent forests, IITA and Cameroon's National Forestry Institute are investigating the potential of integrating timber and food crops. This would allow farmers to plant timber trees and grow crops between them at establishment and once after each routine tree thinning, i.e., when the less vigorous trees are cut every five to ten years. The postthinning periods, with reduced competition and greater light, provide niches for cropping. This mimics traditional systems where crops like cocoyam and



...the farmer sets his plantains among the young, high-density timber trees, *Terminalia vorensis*. Plantain is a leading market food crop in southern Cameroon.

200



plantain are planted in forest gaps created by fallen trees

Prototype systems

At IITA's Humid Forest Station, prototype multistrata systems were established in six- and 17-year-old *Terminalia ivorensis* plantations in 1995. The effects of timber tree densities and low-input soil management regimes were assessed on the growth and yield of plantain, the most important marketed food crop in southern Cameroon.

Initial results are encouraging. Plantain bunches were heavier under tree stands with a 65% canopy cover compared to sole timber tree stands with about 15% cover. Bunches of up to 30 kg were harvested. Greater bunch yield was associated with a markedly lower degree of damage from black sigatoka disease. However, yields per hectare per year were lower, as greater shade prolonged the beginning of flowering. Shade in high tree density plots also reduced

the vigor of the aggressive arable weed *Chromolaena odorata*, allowing less aggressive forest species to establish, thus reducing weeding labor. Beneficial soil fauna, such as earthworms and soil-feeding termites, maintained their activity almost at the same level as in undisturbed forest, contributing to favorable soil properties.

Farmers and researchers have recognized the advantages of diversifying the production base in the humid tropics through the establishment of multistrata systems. A suitable combination of food crops adapted to the multistrata environment and early yielding fruit trees, oil palms, and timber trees is crucial to ensure a continuous flow of revenue and reduce the need to exploit natural forests.





East African Africa has great potential for sustainable agricultural production. Its various ecosystems are naturally balanced. Encouraging farmers to maintain the balance is crucial for the future.



Peter Neuenschwander Finds That Success Breeds Success

The Plant Health Management Division (PHMD) came into existence because of IITA's success in bringing the cassava mealybug under biological control in Africa. Its combining research and biological control, host plant resistance, and habitat management makes it unique in the Consultative Group on International Agricultural Research (CGIAR).

In 1991, in the reorganized structure of IITA, PHMD was given a big role. Since 1994, it has been headed by Peter Neuenschwander. "We're responsible for all integrated pest management projects, which assures collaboration across the different divisions. We work on all mandated crops in the humid and subhumid zones of Africa, some of which are extended into the dry savanna and even the Sahel," he explained.

Apart from the division's contribution to the continuous improvement of Integrated Pest Management (IPM) through research and implementation, recent successes brought major achievements. These involve biological control of the cassava green mite, mango mealybug, larger grain borer, *Striga* management, and the development of the first commercial entomopathogen for use against grasshoppers and locusts.

"We consider our accomplishments in the control of grasshoppers and locusts to be of immense scientific and socioeconomic importance," Neuenschwander noted. "The damage that these pests cause to crops, especially in Africa, is staggering. Since the use of chemical pesticides is undesirable worldwide and economically prohibitive in sub-Saharan Africa, use of environmentally friendly natural enemies to control them is in touch and in tune with the time."

IITA also promotes IPM activities across the entire CGIAR in the form of the Systemwide Program on Integrated Pest Management.

Locusts to be on

Wanted to see

W. H. H. H. H. H.

The year 1914

Integrated strategies to reverse plantain losses in Ghana

A Gesellschaft für Technische Zusammenarbeit (GTZ)-funded IITA project is responding to declining plantain yields in West and Central Africa. The aim of the project is to develop integrated management strategies, incorporating cultural and biological control, that are appropriate for smallholder plantain farmers, so as to reverse the decline in yields and plantation life.

Despite the importance of plantain as a preferred staple in Ghana, the area of production has decreased by up to 14% in recent years due to poor management, low soil fertility, short fallows, and an extensive pest and disease complex. Poor yields have led to abandonment of the crop in traditionally important plantain-growing areas.

The ultimate benefit of the project will be seen in increased plantain production and prolonged plantation life, which will contribute significantly to food security and sustainable agriculture in the region.

The analysis of the situation began with a participatory rural appraisal to determine farmers' perceptions of the constraints to plantain production in Ghana and Nigeria. The appraisals were followed by diagnostic surveys to determine the distribution, severity, and dynamics of pests and diseases, taking account of ecological factors and farming practices.

The preliminary evaluation highlighted the importance of nematodes, weevils, and black sigatoka as major constraints to plantain production. These results were confirmed by on-station trials, showing that infestation by a combination of nematodes and weevils can cause losses of 85%. The problems of nematodes and weevils are compounded by the repeated use of infested planting material.




*Tunnelling damage on a plantain pseudostem caused by the banana weevil *Cosmopolites sordidus*.*

Mycosphaerella fijiensis, a leaf fungus that reduces photosynthetic area, is the causal agent of black sigatoka and can cause up to 80% yield reductions

Radopholus similis, a nematode that causes necrotic lesions on the roots leading to uprooting of the plant, particularly after bunch emergence when the plant is already unstable

Cosmopolites sordidus, the banana weevil, causes significant losses by damaging the corm through burrowing. In severe cases, this causes breakage at the base of the stem



Multifaceted project

IITA, in collaboration with farmers and the Ghanaian Ministry of Food and Agriculture (GMOFA), is developing low-cost techniques to disinfect planting material, followed by rapid multiplication in village nurseries. Considerable success has been obtained with the use of clean planting material and improved management.

Another promising effort involves the use of a fungus to control the banana weevil. A highly virulent strain of the fungus *Beauveria bassiana* (from Uganda) has been identified and can be mass-produced in the laboratory. Preliminary field trials show weevil mortality ranging from 53 to 81%, compared with 7 to 8% mortality in the untreated control.


The project has other offerings. It includes graduate training for several Ghanaian scientists. Short-term training is provided for collaborating

GMOFA staff who assist farmers in improved plantain management. Moreover, the project is strengthening linkages between IITA and national programs in Ghana, while collaboration with a complementary project in East Africa enables information exchange between the two regions.

The ultimate benefit of the project will be seen in increased plantain production and prolonged plantation life, which will contribute significantly to food security and sustainable agriculture in the region.



A banana weevil infected by the entomopathogenic fungus *Beauveria bassiana*



IPM helps farmers increase cowpea production

Most cowpea farmers in sub-Saharan Africa are confronted with low yields, caused by insect pests and diseases. Over the past few years, however, this picture has been gradually changing due to the establishment of a regional pest management project

Known by its French title, Protection ecologiquement durable du niébe, or PEDUNE, it aims to increase cowpea production in the Sahel and African savannas by applying ecologically sustainable pest control. By its nature, integrated pest management is 'by farmers' and not 'for farmers,' so farmers' participation in this project is one of the key features.

During the last 15 years, IITA, in collaboration with the national agricultural research systems (NARS), has developed a number of efficient pest control technologies for cowpea, mainly based on resistant, higher-yielding varieties. Unfortunately, several surveys with the NARS revealed that until 1994 most of these technologies were yet to be adopted by farmers.

In order to increase adoption by farmers, IITA, with financial assistance from the Swiss Development Corporation (SDC), initiated the pilot phase of PEDUNE in 1994. An essential feature of the pilot phase, which started in Benin, Burkina Faso, Mozambique, Niger, and Nigeria, was to carry out diagnostic studies at the farmer level aimed at understanding the major socioeconomic circumstances which influence technology transfer.

Pilot project

A participatory evaluation of the pilot phase, which ended in 1996, revealed promising results and was thus the basis for continuing and

Among the most promising technologies are varieties resistant to Striga, aphids, and bruchids.

Intercropping of sorghum with cowpea traditionally (opposite page) and with an IITA improved cowpea variety (right). The improved variety increases soil fertility which greatly enhances the sorghum's growth.



Solarization is used prior to storage to disinfect cowpeas of bruchids, small beetles that bore into the seeds, damaging or rendering them inedible.

expanding the project to include four new countries: Cameroon, Ghana, Mali, and Senegal.

The second phase of the project started in 1997 and focuses on five areas: dissemination of proven technologies, training of farmers, extension agents, and scientists, farmer-managed on-farm trials to validate technologies, on-station applied research on priority areas, and monitoring and evaluation of project performance and rate of adoption.

Among the most promising technologies are varieties resistant to *Stigra*, aphids, and bruchids, improved storage techniques using solar drying, and the use of botanical pesticides in the field and in storage.

Strong linkages and collaboration among various disciplines and institutions are essential for the project's success, so emphasis is also placed on improving exchange of information and expertise, e.g., by encouraging exchange visits and short-term attachments for national project scientists.



IITA's insect pathology program: applying LUBILOSA's lessons

Using chemical pesticides for the control of insect pests is increasingly unacceptable because they pollute and adversely affect the health and habitat of humans and wildlife. Biopesticides are an alternative.

Until the mid-1980s, the principal means of controlling locusts, for example, was the use of persistent synthetic insecticides. However, because of concern about environmental damage and risks to human health and livestock, persistent chemical insecticides lost acceptability. Nonpersistent chemicals, with little residual activity, became widely used instead.


The Food and Agriculture Organization (FAO) of the United Nations accepted a dossier on the results, and the LUBILOSA mycopesticide was recommended by the pesticide referee group for locust control.

Because these chemicals kill largely by direct spray contact only, they are less effective and often require several applications to achieve control. This increases cost and environmental damage, and puts humans and livestock at risk.

For these reasons, especially during the 1986-89 locust plagues in Africa, a big donor investment was made in finding alternatives.

Almost a decade ago, IITA, in collaboration with the International Institute of Biological Control (IIBC) of CAB International, the Comité permanent inter-état de lutte contre la sécheresse dans le Sahel (CILSS), Gesellschaft für technische Zusammenarbeit (GTZ), and West African national partners, began work on biological control. Early on, the use of *Metarhizium flavoviride* was picked up by the LUBILOSA project, a French acronym for "biological control of locusts and grasshoppers."

Metarhizium anisopliae infecting the larger grain borer *Prostephanus truncatus*




Metarhizium is a fungus. It is applied in an oil mixture to avoid rapid drying of the spores. The fungus, which occurs naturally throughout Africa, has no adverse effects on mammals, including humans.

In the ongoing third phase, IITA leads the socioeconomic evaluation of *Metarhizium*, runs a comprehensive field trial program, studies *Metarhizium* impact on nontarget invertebrates, and produces fungal spores.

Safe fungus

The efficacy of *Metarhizium* was demonstrated to international standards against variegated grasshoppers (*Zoniocerus*



variegatus), rice grasshoppers (*Hieroglyphus daganensis*), the Sahelian grasshopper complex, and the Senegalese grasshopper (*Oedaleus senegalensis*)

Mortality started six days after applications and reached 70-95% after 14 days. There was no reinvasion of the treated plots. Desert locust (*Schistocerca gregaria*) and hopper bands were also killed.

Ecotoxicological laboratory and semifield tests, following the approved testing scheme of the US Environmental Protection Agency, revealed no harm by *Metarhizium* on several groups of beneficial and other ecologically important insects. Also, in large-scale operations in Niger, the nontarget fauna of the Sahel was shown not to be at risk. This also demonstrates the advantage, from the environmental point of view, of the biological product over chemical pesticides such as fenitrothion.

Other positive effects included secondary recycling of the pathogen after a single spraying. In other words, the fungus multiplies on the dead grasshoppers, increasing the killing action. This has significant consequences for the economics of biopesticide use.

The Food and Agriculture Organization (FAO) of the United Nations accepted a dossier on the results, and the LUBILOSА mycopesticide was recommended by the pesticide referee group for locust control. Transferring the technology to commercial partners in Europe and Africa has already begun, and it will lead to industrial mass production for application in Africa in a short time.

An economist is helping guide the complexities associated with commercial development of the technology.

Future uses

Because biological control poses less risk to humans, animals, and the environment, it is preferred over chemical insecticides. For this reason, IITA's insect pathology program is looking beyond LUBILOSА, to apply entomopathology to other pests. These include trials against Moroccan locust, brown locust, migratory and red locust, and implementation of the use of *Metarhizium* for *Zonocerus* control in humid-zone countries.

LUBILOSА has opened a door to the future of entomopathology in sub-Saharan Africa. This includes many fascinating research avenues, for example, production of mycopesticides in liquid fermentation and the use of protozoans, viruses, and bacteria.


For some of the pests, classical biological control using pathogens may be possible, for others, development of a biopesticide is preferred. Scientists are exploring using pathogens in biocontrol of the cassava green mite, against the larger grain borer, the banana weevil, and termites. A termite-active isolate of *Metarhizium anisopliae*, received from the International Centre for Insect Physiology and Ecology (ICIPE), was tested against four species of termites, both as a repellent and as a mortality agent. Each of these insects costs African farmers millions of dollars annually in crop losses.

BioNet International and IITA's insect taxonomy role

BioNet International supports conservation and the wise use of the environment. This global network is dedicated to achieving realistic self-reliance in taxonomy within developing country subregions. It pools expertise, experience, information, technologies, and infrastructures to support sustainable agricultural development.

Plans are on the drawing board to include South America, Meso-America, the Nile Basin, China, and the Middle East.

A drawer with Lepidoptera caught at one lighttrap site in Benin provides insights about moth diversity in West African environments.



BioNet is an initiative of CAB International, and it provides systematic backing for biodiversity. By strengthening taxonomic capabilities, it removes impediments to the development of integrated pest management systems and effective quarantine, which adversely affect diversified agricultural production in the developing world. Network hubs are focused on the Caribbean, southern Africa, the South Pacific, South Asia, Southeast Asia, and recently East and West Africa.


Plans are on the drawing board to include South America, Meso-America, the Nile Basin, China, and the Middle East. The network encourages South-South and North-South cooperation.

Biosystematics of arthropods, nematodes, and microorganisms (fungi, bacteria, and viruses) are the biological focus of BioNet. Recognizing the importance of sound taxonomic foundations for sustainable agricultural development, IITA has taken the lead to play a central function in establishing the West African node, to be known as WAFRINET or West African Net.

Collaborative and regional networking

IITA is thus playing a catalytic role in fostering regional expertise through active networking. In concert with scientists from 11 francophone and six anglophone West African countries, a two-year program has been outlined.

A cetonid beetle acquisition awaiting integration into IITA's reference collection at the Biological Control Center for Africa Cotonou Benin



emphasizing information and communication, training of biosystematists, rehabilitating collections and other sources, and developing and using new technologies, especially electronic aids to identification

IITA's unique institutional facility was established, in particular, with financial support from Austria. Linked with BioNet, it is providing an efficient information service, covering traditional taxonomy, molecular techniques, new records, and current pest distribution data. Information on natural enemies/biological control and bioremediation agents can ultimately be accessed through the network's databases. IITA will thereby keep participants up to date in its fields of expertise.

IITA's Plant Health Management Division trains science graduates in general and in applied taxonomy. This is


done through joint courses at overseas universities and institutions, and at local subregional academic and scientific centers. It leads to joint MSc and PhD degrees. Technician-level instruction is also offered through the network.

Information is vital to such an undertaking. Communication services provide key institutions with minilibraries and databases, computer equipment, and electronic communications systems, including E-mail capacity and Internet access.

The global network is supported by a consortium of developed-country expert centers, and it is managed by a coordinating committee and a technical secretariat in the CAB International headquarters in the UK.



The 200 square meter Insect Museum at IITA's Biological Control Center for Africa houses an insect reference collection of some 35 000 specimens and 2 100 identified species itemized in databases. With an acquisition of more than 5 000 specimens a year the museum contains one of the largest insect collections in West Africa.



Taking its nutrients directly from the water in which it floats the water hyacinth has become a nuisance in warm regions throughout the world



Partnership for food: the Technology Testing & Transfer Unit

IITA's Technology Testing & Transfer Unit operates on the principle that developing ecologically sustainable pest control technologies and transferring these technologies to farmers within various farming systems is a dynamic process

The Technology Testing & Transfer Unit (TT&TU), established in 1990 with funding initially from Germany and later from Austria and Switzerland, is guided by a philosophy as much as by objectives. As a guiding principle, collaboration between IITA and national scientists must be based on reciprocal trust and equal rights. TT&TU considers a project as being entirely owned by the national program.

Also, personal dialogue and interaction are more important than

letters, faxes, and mail.

Collaboration is based always on personal contacts and mainly determined by the interaction of individual personalities. This is the foundation of TT&TU's approach to strengthening national plant protection programs.

Based on an evaluation of its activities since its inception, the unit's objectives have been largely achieved. A solid informal network of information by various forms of contact has been established throughout sub-Saharan Africa.

The national agricultural research and extension systems will have to take the initiative themselves and introduce their requests for support according to their concrete work plans.

The next phase


The unit is already well into its next phase. TT&TU will continue to support national agricultural research and extension systems to help them test and implement biological control measures, adapt other integrated pest management (IPM) measures, and achieve sustainability through training and information sharing.

In keeping with the philosophy of equality, participants will not be selected to cooperate in the project. The national agricultural research and extension systems (NARES) will have to take the initiative themselves and introduce their requests for support according to their concrete work plans.


Also, the project will stress that all potential local resources for funding and equipment have to be exhausted before external support is given. This breaks any possible form of paternalism and conforms with the philosophy that help for self-help is paramount. Therefore, the unit trains national scientists and

technicians, characterizes ecosystems and the socioeconomic status of farming systems, conducts prerelease surveys and local rearing of beneficial agents for field release, does on-station and on-farm technology testing, and assesses the impact of technologies by assisting and improving the collaboration between national scientists and their counterparts in IITA's Plant Health Management Division.

Future collaborative research activities will continue to focus on technology testing and transfer with regard to IPM of cassava, cowpea, and maize. Special effort will be made on the biological control of cassava green mite, flower thrips, and pod borers on cowpea, maize stem borers, and the larger grain borer. New collaborative IPM projects are being developed with NARES on water hyacinth, mango mealybug, and *Stiga*.



The national agricultural program in Tanzania, supported by TT&TU, developed facilities for rearing agents to be used against major crop pests.



...anges in
Africa, the

...the human
...capability of
...al agricultural
...ch systems cannot
...emphasized





Michael Bassey Praises Training In Technology Transfer

Michael Bassey, director, IITA's International Cooperation Division, praises training as "the bedrock of IITA's support to national programs " Under his leadership, IITA has intensified its in-country training


Bassey believes that this capability can be maintained by using available training resources and preparing national programs to take lead roles in research-related training. Group training is the best way to accomplish this, he noted.


"Group training strengthens national programs so they can conduct the research essential for their own development. We encourage sustained advancement in research because it enhances collaboration."

In the past three years, IITA has trained more than 2000 technicians and scientists from sub-Saharan Africa, with an increasing proportion being women. Approximately half of these were trained at off-campus sites.

Research networks are a major conduit for training and dissemination of technology. Bassey stressed the networks' role. "Networks interconnect and channel technology, they also provide the organizational structure for training. IITA is among the few institutes which have invested significant core resources to bolster national research capacity."

In Bassey's view, training and network-strengthening go hand in hand. "Networks are a partnership, and the essence of a partnership is sharing responsibilities. As a good partner, one of IITA's major responsibilities is to train national collaborators in research, program management, and leadership."





In 1997 IITA established a home page on the World Wide Web. The initiative, along with the Institute's participation in AfricaLink, a US Agency for International Development project to link African researchers through E-mail with IITA and other IARC resources in Africa, reflects the International Cooperation Division's objective to improve the dissemination of information.

El Niño, NARS, and IITA News on the Web

El Niño's anticipated effects on weather and agriculture in eastern and southern Africa led IITA to prepare a plan to mitigate the disruption of root and tuber crop production.

Responding to a request from the US Agency for International Development, IITA focused its plan on cassava and sweetpotato. It was developed with the Southern Africa Center for Cooperation in Agricultural Research (SACCAR), the Association for Strengthening Agricultural Research in Eastern and Central Africa (ASARECA), and the International Potato Center (CIP).

The plan involved use of improved germplasm, rapid multiplication, and use of appropriate postharvest technologies. IITA is working through regional networks to implement production activities in selected countries.

National program collaboration

1997 was a milestone in IITA's work with the Conférence des responsables de la recherche agronomique africains (CORAF), which coordinates agricultural research in West and Central Africa. Both pledged to strengthen collaboration, which is expected to enhance the adoption of IITA technologies.

The primary objective of the International Cooperation Division is to develop long-term research strategies with national agricultural research systems in Africa.


CORAF, based in Dakar, Senegal, was established in 1977 to oversee research in the French-speaking countries of Africa. Its mandate, now covering anglophone and francophone countries in West and Central Africa, is to ensure food security. Strengthened CORAF-IITA collaboration is expected to lead to better management of research resources.

IITA was approached by the International Fund for Agricultural Development (IFAD) to provide training and technical assistance to Zanzibar's Food Crops and Cash Crops Commission. This involved smallholders who grow cassava and sweetpotato, and the operation of a soil and plant laboratory and tissue culture laboratory. Rapid multiplication and distribution of cassava and sweetpotato planting material were high on the agenda. The

project is supported by the World Bank. To further assist national programs, IITA's Training Unit produced a pocketbook entitled "Rapid Multiplication of Cassava."

In Ghana, another World Bank-financed, IFAD-executed project involving IITA was aimed at improved production, marketing, and processing of roots and tubers through adoption of new technology. As of the end of 1997, it had not yet been funded.

IITA, responding to the Democratic Republic of Congo's (DRC) need for seeds, delivered 12.4 metric tons of maize and top varieties of soybean seed to the country, in cooperation with World Vision International, within the framework of the US Agency for International Development-funded "Seeds of Change" program. The seeds are from plants bred by IITA for use in agroecological conditions in the major production areas of the DRC; they have good resistance to the prevailing diseases. They are being multiplied by the national agricultural sector and distributed to farmers.




Recognizing the wide need, IITA's Training Materials Unit produced Rapid Multiplication of Cassava. The handy pocketbook (US\$5) provides step by step instructions for increasing cassava planting material.

World Wide Web

1997 saw IITA establish a site on the World Wide Web. The initial concept and design of the Web site evolved through the efforts of computer and graphics specialists, writers, editors, scientists, and others in the Institute. IITA's Internet address is <http://www.cgiar.org/iita>.

The home page has six headings which are hypertext reference links, i.e., each heading is linked to another subheading that contains information about that particular heading. The links are research, information (IITA publications and library), partnerships, about IITA, news, and the CGIAR. The home page also contains information about IITA mandate, crops, and research divisions.

Being on the Internet helps meet the objective of improving the dissemination of IITA's research results. The primary objective of the International Cooperation Division is to develop long-term research strategies with national agricultural research systems in Africa.



**For
the
Record**

Research Highlights

IITA's work is organized around 16 multidisciplinary research projects and one project for the dissemination of results to national research systems. Some projects focus on production systems for specific crops or crop combinations, in some cases for a specific agroecological zone. Others are thematically oriented, cutting across commodities and agroecological zones.

IITA also serves as the convening organization for two international programs of the CGIAR: the Ecoregional Program for the Humid and Subhumid Tropics of Sub-Saharan Africa and the Systemwide Program for Integrated Pest Management.

The following section presents the goal and research highlights of each IITA project for 1997. The summaries are not exhaustive of the work begun or completed during the year; rather, they describe some key scientific results and are intended to give readers an idea of the breadth of research themes and problems being investigated by IITA scientists.

Project 1

Short Fallow Stabilization

Goal

Increase farm productivity and arrest resource degradation due to land use intensification through sustainable short fallow systems.

Highlights

- ✦ Herbaceous legume evaluation trials (single and mixed legume species) were established on low P soils with and without P application. The legumes included 12 accessions of *Mucuna pruriens*, 21 accessions of *Lablab purpureus*, 25 accessions of *Aeschynomene histrix*, and 54 accessions of *Centrosema pascuorum*. Preliminary results indicate large variation between species accessions for their adaptability to low P soils.
- ✦ The herbaceous legume seed collections of ILRI and IITA have been collated into a comprehensive list of some 388 seedlots. During 1997, 171 seedlots, totaling over 100 kg, were distributed to NARS and IARC scientists, mainly in West Africa.
- ✦ The center for cover crop seed and information exchange was established in Cotonou (with IDRC/IITA funding) to support cover

Research Highlights

crop seed and information exchange in West Africa. As a first task, several tonnes of *Mucuna* seed were produced for distribution to collaborators.

✦ Research was undertaken at Ibadan to measure benefits of 13 herbaceous legumes for subsequent maize. On average, subsequent maize grain yield was increased by 70%, compared to the natural grass fallow.

✦ Field studies were conducted to investigate the influence of three *Mucuna* varieties and two levels of fertilizer on the control of *I. cylindrica* during the year of cover crop planting and one year later. After one growing season, *M. pruriens* var *utilis*, *M. cochinchinensis* and *M. pruriens* var IRZ reduced *I. cylindrica* shoot density by 50, 76 and 68%, and shoot dry matter by 72, 92, and 79%, respectively. Maize grown one year after *Mucuna* required 50% less weeding than plots without *Mucuna*. Maize grain yield was higher in plots previously seeded to *Mucuna* than in plots without *Mucuna*.

✦ *Calliandra calothyrsus* hedgerow fallows and well-established *Pueraria phaseoloides* fallows can significantly suppress the growth of *Chromolaena odorata* and associated natural fallow species. At the end of 2 years of fallow, there was an 80% reduction in ground cover and a 90-95% reduction in biomass of these naturally occurring species compared to unplanted fallows.

✦ A legume based system close to the farmers' practices was evaluated over a period of 2 years. Maize and legumes were intercropped in the first year, and cassava/maize intercrop was rotated in the second year. Results from the first year data showed that the legume based systems are economically attractive.

✦ In 1997 farmers designed trials to integrate *Mucuna* into their maize-based systems in the northern Guinea savanna zone. A pre-maize *Mucuna* system was preferred, although many are also in favor of a maize/*Mucuna* relay system. Farmers recognized *Mucuna*'s ability to improve the soil, and to reduce weeds (including *Striga*).

✦ Using georeferenced databases on the importance of maize in the cropping systems and suitability of soils to support sustainable maize production, target domains for short fallow systems were proposed. They show that cereal based short-fallow systems are likely to have high, medium, and low impact on 386, 256 and 527 million hectares respectively in SSA. Potential adoption domains were also mapped. These will assist in selection of sites within the benchmark area for technology development and identify potential pilot sites for technology targeting.

Project 2

Agroecosystem Development Strategies

Goal

To guide ecoregional research and policies for poverty alleviation and sustainable development of agroecosystems in the subhumid and humid zones. Activities feed directly into the Ecoregional Program for the Humid and Subhumid Tropics of Sub-Saharan Africa (EPHTA).

Highlights

✦ Security over land tenure had a positive impact in Benin Republic on the adoption of *Mucuna*, though the technology is an annual crop probably because farmers perceived its positive long run effects on the improvement of natural resources.

- ✦ About 70% of the total labor input required in cassava farming is on postharvest activities, most of which is contributed by rural women and children. Significant factors that positively influence adoption of improved cassava processing technologies are the proportion of cassava processed, type of products, mechanization of land preparation, and the number of women with city life experience.
- ✦ A farmers' survey on the use and availability of organic and inorganic fertilizers was conducted in 5 villages in the derived savanna ecoregional benchmark and 2 villages in the NGS ecoregional benchmark area. The results showed that inorganic fertilizers are used by most farmers in both benchmark areas but that application rates vary widely between farmers' fields, and crops. Organic amendments are less commonly used and only on specific fields.
- ✦ A rapid appraisal of 2009 community-based organizations (CBOs) was completed in 475 villages. The survey shows that traditional credit-solidarity groups (tontines) (43%) and religious organizations (44%) form the overwhelming majority (87%) of village based organizations in southern Cameroon. The study shows a strong association between the presence of NGOs and the emergence of both farmers' federations and common initiatives groups at the local level. The information gathered from the surveys is now being used in the development of broad based partnerships with farmers' organizations and communities in the forest margin benchmark area of southern Cameroon.
- ✦ Representative farm models have been built for that area. Each model will be used to evaluate potential financial and economic benefits of the new crop and natural management technologies being developed and tested in the benchmark area.
- ✦ A household characterization survey was conducted in 15 villages among 225 households of the forest margins ecoregional benchmark area of southern Cameroon. Results show that, in general, households in areas with high land pressures have intensified their production systems to a much greater degree than in areas with low land-use pressure.

Project 3

Biological Control and Biodiversity

Goal

To enhance the livelihood of resource poor farmers and maintain sustainability of farming systems through biological control and preservation of biodiversity

Highlights

- ✦ A large-scale aerial application of *Metarhizium flavoviride* against the Senegalese grasshopper was carried out in collaboration with the Niger Plant Protection Service. Farmers preferred the mycopesticide to the chemical pesticide.
- ✦ Collaborative trials with a Malian NGO demonstrated good control of Sahelian grasshoppers with mycopesticides applied by village brigades.
- ✦ Ecotoxicological studies demonstrated the low negative environmental impact of the mycopesticide. The registration process for it has been initiated.

- ✦ The mass production unit was run at full capacity demonstrating the technical feasibility of the developed production process. Negotiations with commercial companies are well under way and commercial production is scheduled to start in 1998.
- ✦ Participatory field trials in southern Benin demonstrated control of the variegated grasshopper at very low doses in individual fields. Reduced crop loss is estimated to be higher than the cost of biocontrol, making individual purchase and application of the biopesticide attractive.
- ✦ Biosystematic capabilities were consistently enhanced, reflected in the extent of the identification service offered, the accumulation of voucher insect specimens and the availability of collection databases, complemented by a large reference bibliography.
- ✦ In 1997 it was documented in southern Benin that the two serendipitously introduced parasitoids, *Encarsia baitiensis* and *E. guadeloupae* had brought under control the spilling whitefly *Aleurodicus dispersus* a new exotic pest of cassava and ornamentals.

Project 4

Integrated Management of Legume Pests and Diseases

Goal

To reduce the risk of crop losses in farmer's fields in sub-Saharan Africa by means of integrated pest management technologies that increase cowpea and soybean productivity in a sustainable manner

Highlights

- ✦ Higher levels of resistance identified for flower thrips (initial progeny testing from crosses conducted under intense natural population) and PSBs in cultivated cowpea.
- ✦ Multiple pest resistance confirmed in elite cowpea breeding lines, and resistant varieties to CoBB identified.
- ✦ Preliminary results from feeding bioassays on transgenic cowpea plants using MPB neonates indicated some level of successful transformation with *Bt* protoxin *Cry* IA genes.
- ✦ Resistance to MPB confirmed in new gene products and secondary metabolites from African yam bean obtained locally.
- ✦ A drastic disruption of the utilization of fat in the process of oogenesis in MPB in *Vigna oblongifolia* (e.g. TVNu 42) explains biochemical basis of resistance.
- ✦ MPB resistance of TVNu 72 (*Vigna vexillata*) mainly biophysical (trichomes and/or other podwall characteristics). *C. tomentosicollis* resistance due both to biophysical (trichomes) and biochemical factors.
- ✦ Survey data from over 200 cowpea fields in Benin indicate very low population levels for the local strain of the thrips parasitoid *Ceranisus menes* (Hym, Eulophidae).
- ✦ Overall mortality of *M. utrata* larvae due to pathogens 13%, higher percentages on plants with prostrate growth habit.
- ✦ Semiselective medium and polyclonal antibodies for detection of CoBB bacteria developed and tested.

- ✦ Pathogen diversity of *Xanthomonas campestris* pv *vignicola* determined pathologically, biochemically and genetically, host ∞ pathogen interactions studied, no races identified
- ✦ Genetic diversity of *M phaseolina* strains from various origins characterized
- ✦ Existence of symptomless virus infection demonstrated consequences for breeding programs and virus indexing assessed
- ✦ Over 120 cowpea lines screened for resistance to viruses, including cowpea aphid-borne mosaic, blackeye cowpea mosaic, cowpea mottle, cucumber mosaic, and southern bean mosaic viruses and the results sent to the breeder for pyramiding of resistance
- ✦ Heat treatment disinfects contaminated seeds (CoBB and *M phaseolina*)
- ✦ The combined use of solar drying and improved storage for cowpea demonstrated at PEDUNE pilot sites
- ✦ PEDUNE countries have conducted trials with different formulations of neem for both field and storage applications

Project 5

Integrated Management of Maize Pests and Diseases

Goal

To reduce pre and postharvest losses of maize caused by insects, diseases, and fungal grain contaminants IITA scientists work with NARES partners to diagnose constraints, test resistant germplasm, and explore options in host plant resistance, habitat management, and biological control. Combinations of control options are assembled in an integrated pest management (IPM) package and tested with the participation of scientists, farmers, and socioeconomists

Highlights

- ✦ First discovery of oospores of the maize strain of *Peronosclerospora sorghi*, indicating that removal of crazy top at the early stage of development will preclude oospore development
- ✦ A deployment strategy to saturate an area with DM-resistant maize variety was initiated in 9 villages across 2 local government areas of Ogbomoso in Oyo State, Nigeria. About 25 ha of land was planted and 6 tonnes of seeds generated from selected farms
- ✦ Following establishment of an East African strain of the larval parasitoid *Cotesia sesamiae* on *Sesamia calamistis* in southern Benin, first releases have been made in eastern Nigeria
- ✦ New candidates for redistribution BC against *E saccharina* and *B fusca* were identified in West and South Africa, respectively
- ✦ Data from the trapping network has confirmed the biological control of *P truncatus* by *T nigrescens* in southern Benin and has revealed serious *P truncatus* infestations in northern Benin
- ✦ Weather driven population models for *P truncatus* and *S zeamais* have been completed, and these will be linked via grain damage rates to a grain loss model for stored maize
- ✦ Atoxicogenic isolates of *Aspergillus flavus* were identified from each of four agroecologies of the Benin Republic and characterized as to

Project 6

Integrated Management of Cassava Pests and Diseases

their vegetative compatibility. Initial trials indicated good biocompetitive fitness of two of the isolates against toxic strains of *Aspergillus* spp.

✎ The causal agent of a maize disease new to the region was identified as the basidiomycete *Marasmiellus* spp. The symptom of the disease, a horizontal banded leaf blight, was seen in Ghana, Nigeria, and Cameroon and was reproduced in Koch's postulates by spraying a mycelial suspension on 12 day old maize plants.

Goal

Increase cassava productivity in sub-Saharan Africa

Highlights

✎ Exotic predatory mites, released for the biological control of the cassava green mite (CGM) showed excellent establishment and spread in West Africa. In field trials the predators reduced CGM populations by an average of two thirds and increased cassava yields by a third, resulting in farmer benefit estimated at US\$70 per ha. Based on these results an Africa wide biological control of the pest was initiated. The virulence of exotic isolates of the pathogenic fungus, *Neozyglites floridana*, was evaluated and the indigenous pathogenic fungus, *Hirsutella thompsonii*, isolated for experimental releases.

✎ Cassava root yield losses to the cassava mosaic disease (CMD) pandemic in western Kenya were estimated at 42%.

Characterization of epidemiology and diagnoses of the whitefly vector were initiated in diverse agroecologies in West and Central Africa (Benin, Cameroon, Ghana, Nigeria), EARRNET countries (Kenya, Rwanda, and Uganda) and SARRNET countries (Malawi, Mozambique, Tanzania and Zambia), largely within the framework of the system wide IPM initiative.

✎ Yield loss of 20-30% due to cassava bacterial blight (CBB) was recorded in resistant TMS 30572 and susceptible varieties comparing 2 infection levels. CBB epidemiology trial results stressed the importance of soils and weeds as inoculum sources of the bacteria and of variegated grasshopper in the transmission of the disease.

✎ Yield loss of 30% was recorded for cassava brown streak disease in Tanzania and Malawi, with the variety TMS 30040 and five locally selected clones showing resistance to the disease.

✎ Two new diseases of cassava, *Curtulania* leaf and stem blight on field and stored cassava stems and *Nattrassia* root and stem rot, were reported and their prevalence determined in Benin and Nigeria.

✎ ELISA-based CMD diagnostic technology was decentralized to NARS, and PCR-based technology was established to detect new cassava mosaic virus variants.

✎ Twelve women scientists from West and Central Africa completed postgraduate training in cassava IPM and extension methods. Eleven NARS technicians from 5 SADC countries were

trained in cassava IPM, and farmer field school training was conducted at 25 sites in West and Central Africa

✦ Cassava information resources were produced. These included geo-referenced distribution maps of plant protection constraints in Ghana, Benin, Nigeria, and Cameroon, GIS maps of the distribution and characteristics of CMD in Uganda and the wider Lake Victoria basin, cassava IPM extension materials, and directories of personnel, institutions, cassava projects, databases of gray literature, bibliographies, and cassava references.

Project 7

Improving Plantain- and Banana-based Systems

Goal

Development and dissemination of improved technologies for sustainable Musa production in different ecologies of sub-Saharan Africa

Highlights

- ✦ Geo-referenced databases. A diagnostic survey of pests and diseases in Uganda has been completed.
- ✦ Knowledge of pests and diseases enhanced. Identification of banana streak virus in Benin, Ghana, Kenya, Malawi, Nigeria, Tanzania, and Uganda confirmed its widespread occurrence. Also, a new virus, tentatively named banana die-back virus, was identified in Nigeria.
- ✦ Distribution patterns for nematode species were determined in Ghana, Nigeria, Rwanda, and Uganda.
- ✦ Yield losses from pests and diseases determined. Yield loss in plantain (AAB) from nematodes and weevils in Ghana was established. More than 75% of plants infested with both nematode and weevil failed to reach maturity, and there was an overall yield loss of 85% from both pests.
- ✦ Yield losses in East African highland bananas (EAHB) (AAA) from weevil increased with crop cycle, reaching 50% in the 4th year, while lesion nematodes caused a 30–40% yield loss in EAHB, but no detectable yield loss in cooking/beer banana (ABB).
- ✦ IPM strategies available. Pseudostem traps reduced weevil numbers in on-farm trials in Uganda, and endophytic fungi isolated from banana corms caused egg mortality under laboratory conditions and reduced weevil damage in nursery trials.
- ✦ Utilization of germplasm enhanced. Breeding capability was fully established in East Africa following transfer of breeding technology from IITA Onne. A first diploid hybrid, derived from highland banana and with black sigatoka resistance, was selected for use in EAHB breeding.
- ✦ Characterization of host plant tolerance and resistance to nematodes, including the identification of 4 resistant and tolerant hybrids, was achieved.
- ✦ Improved genotypes and populations available. Eight new triploid and tetraploid plantain and banana hybrids were selected, based on high bunch weight, large number of hands per bunch and

large fruits, black sigatoka resistance and good growth habit
Selection for BSV resistance/tolerance has identified 4 promising banana and plantain hybrids

✎ Virus indexing techniques were developed to ensure that IITA's plantain and banana hybrids can again be distributed to other African countries

✎ NARS capability enhanced Twenty eight national scientists were trained in *Musa* breeding, nematology and/or entomology, and the use of BSV diagnostics

Project 8

Integrated Management of Striga and Other Parasitic Plants

Goal

To reduce infestations of parasitic plants with a focus on Striga spp. The project is implemented in collaboration with NARES. Through integrated management practices, emphasizing cereal rotations with selected nitrogen fixing cultivars of legumes, crop yield losses due to parasite infestations are reduced while soil conditions are improved

Highlights

✎ A very positive external review recommended that a world center for parasitic plant research be developed at IITA

✎ Demonstration of an integrated crop and land management program for *Striga hermonthica* control was conducted in the moist savanna of Nigeria. Crops planted in this season included cultivars of soybean, cowpea, and cotton selected for their ability to stimulate *S. hermonthica* seed germination. Improved tillage and weed control practices were also implemented. A successful field day was conducted for farmers, NARES, and NGOs to observe the demonstration and learn about integrated *Striga hermonthica* control

✎ Significant *Striga hermonthica* seedbank reduction was again demonstrated in farmers' fields using rotations with selected soybean cultivars

✎ Incorporation of acetolactate synthase (ALS) inhibitor resistance into open-pollinated IITA maize lines and inbreds was completed

✎ Ethylene-producing bacteria were identified and successfully tested for their ability to stimulate *S. aspera*, *S. gesnerioides* and *S. hermonthica* seed germination. Results showed that 2 isolates of *Pseudomonas* sp. were more effective than the definitive synthetic germination stimulant (GR24) in stimulating parasite seed germination

✎ Over 1600 progeny from 5 maize populations were screened under artificial infestation with *Striga hermonthica* in replicated trials. Partially inbred lines from TZL Comp 1 C3 with very low *S. hermonthica* emergence in the field were selected. Six STR populations were multiplied for on-farm trials with funding from the Korean government, to be organized by PASCON in 1998. A tolerant early maturing cultivar, EVDT 97STR, as well as EV IWDSTR, and a resistant inbred (1102) will be available for regional testing in 1998

Seventy five S2 lines were extracted from a backcross population derived from *Zea diploperennis* and initial crosses were made for molecular mapping of resistance in a Rockefeller funded project in collaboration with CIMMYT and KARI. These lines were also sent to CIMMYT and KARI Kenya for evaluation under natural *S. bermonthica* infestation.

A survey on farmers' harvest practices related to *S. bermonthica* seed contamination of crop seeds was completed on about 80 farmers' fields cropped to cereals. A large proportion of cereal seed samples was found contaminated before and after harvesting. The results confirm the role of the farmer's harvesting method in the contamination of crop seeds and the spread of *S. bermonthica* with crop seeds.

A total of 25 improved cowpea breeding lines were developed with resistance to *Striga gesnerioides*. A project on cowpea resistance to *S. gesnerioides* has been funded in collaboration with the University of Virginia, USA.

Hybridization and cytological studies with *S. bermonthica* and *S. aspera* showed that the 2 species differ by only one haploid chromosome, they hybridize readily, and that hybrids can be found in fields where the 2 species are sympatric.

Project 9

Improving Postharvest Systems

Goal

To increase the income generating capability and improve the nutritional status of farmers, processors, and consumers in both the rural and urban communities of Africa.

Highlights

Results from monitoring surveys in the third year of the Soybean Utilization Project in Nigeria, Ghana, and Cote d'Ivoire showed increasing interest in the use of soybean at the household level.

A socioeconomic analysis of improved postharvest processing showed that the majority of equipment significantly improved income generation. The most remunerative equipment had the capacity for multi-crop applications.

Adoption of the high quality cassava flour technology is increasing in Nigeria, and there is evidence that similar activities have started in Tanzania. Developed at IITA, the technology is being disseminated through NGOs, extension agents, and from farmer/processor to farmer/processor.

Surveys in Nigeria revealed that cassava is being used as the raw material for several commercial-scale processed products including biscuit manufacturing, starch extraction, noodle production, and industrial distillation into alcohol.

Results from a study on yam chips processing was completed in Benin, Nigeria, and Togo. This investigation showed that yam processing was highly profitable when developed with specific varieties and in association with short fallow systems.

Tests on the efficiency of cassava processing equipment, in Uganda and Kenya, led to the development of a number of modified designs in processing equipment for root crops.

Project 10

Farming Systems Diversification

✦ Cassava processing plants were established in three districts of Uganda. The projects were designed to develop stronger linkages between IARCs, national programs, NGOs and community-based organizations (CBOs), and this approach proved to be highly effective with an internal rate of return of 30% in the first year.

✦ Four training courses were held on manufacturing and servicing of improved agro processing equipment for 32 technicians in Benin, Ethiopia, Ghana, Guinea, and Togo.

✦ A workshop was organized in Nigeria which brought together industrialists, cassava producers, NGOs, extension agents, extension services and government agencies for product quality.

Goal

NARES and international system scientists develop, with farmers, new, diversified, and complementary income generating enterprises in West and Central Africa

Highlights

✦ **Multistrata systems** Focused group interviews with 79 villages in the forest margins benchmark area and the East Province of Cameroon were conducted to examine the reaction of cocoa farmers to recent relative price declines in the cocoa market. Farmers were observed to have diversified their holdings in three ways: diversification from tree crop cultivation into food crops (most common); diversification within the plantation through an increased inclusion of fruit trees, and diversification into other tree crops.

✦ **Specialty crop systems** A survey of over 500 market women in the humid forest zone of Cameroon indicated that average daily earnings selling indigenous vegetables were significantly lower than those for exotic vegetables, such as tomatoes and cabbage. However, indigenous vegetables offer a particularly important employment option for the lower economic rungs of urban and rural women in Cameroon.

✦ Cassava leaves are an important leafy green vegetable in Central Africa. Six cassava clones with different plant architectures were selected to investigate the possibility of increasing edible leaf production. Preliminary results indicate big differences in the production of edible leaves. Although pruning facilitates leaf harvest by women, monthly harvests can lead to a significant drop in leaf yield. Pruning also decreases subsequent tuber yields.

✦ **Inland valley systems** Cowpea variety IT90K-284-2 was shown to be a good crop for diversification of crop production in inland valleys of the northern Guinea savanna benchmark area in Nigeria. Over 2 years of testing on farmers' fields during the dry season, it produced more than 1 t/ha of grain, even without insecticide application and a net revenue of approximately US\$500/ha.

✦ **Mixed farming systems** The incipient market for crop residues is expanding in the Guinea savannas of West Africa. An

economic analysis showed that the future of the market for crop residues is constrained by the transportation cost which was found to represent 47–58% of their total cost

✎ A survey was conducted with 150 farmers in northern Nigeria to assess the impact of a cattle fattening program on the socioeconomic well-being of the participating farmers. It showed a net benefit of US\$113 per cattle fattened or about US\$450 per farmer per year. The findings suggest that income generation could be increased if the cost of feeding could be reduced and the supply of both veterinary services and medication could be improved through a credit facility program in favor of cattle fatteners.

✎ Reciprocal benefits from crops and livestock in a mixed farming system were investigated from a survey of 150 agropastoralists in two areas at different gradients of resource-use intensification in the northern Guinea savanna of Nigeria. The productivity of the system was threefold higher in a market-driven area than in a population-driven area. Crop and livestock enterprises yielded the highest returns to land at higher cultivation intensity while the returns to labor were the highest at low cultivation intensity for areas with poor and good market opportunities. In areas where land becomes scarce and labor abundant, increased intensification in resource use is driving farming systems towards greater crop-livestock integration and productivity.

Project 11

Cowpea-Cereals Systems Improvement in the Dry Savanna

Goal

To develop adoptable crop varieties and agronomic practices for the Sudan savanna and Sahel, which will increase the total productivity of the dominant farming systems. The project integrates research by scientists from IITA, ILRI, and ICRISAT who are working on grain legumes, cereals, and livestock in the dry savannas of West Africa. Research institutes in Nigeria (IAR/ABU) and Niger (INRAN) are also active members.

Highlights

✎ **Constraints analysis** Survey of cowpea fields in the Sudan-Sahelian region in 1996 and 1997 indicated a decreasing gradient of *Maruca* pod borer pressure from high rainfall areas to low rainfall areas but an increasing pressure of aphid, thrips *Striga gesnerioides* and drought.

✎ **Improvement of local landraces** From 150 local landraces of cowpea screened in 1996, 9 were crossed with donor parents for incorporation of genes for resistance to aphid, thrips, bruchid, *Striga Alectra* and viruses by the backcrossing method.

✎ **Improved grain-type cowpea varieties** A total of 746 new cowpea breeding lines were evaluated for grain and fodder yields in different cropping systems. A number of lines showed considerable resistance to several biotic constraints and yielded between 1.5 and 2 t/ha.

✎ **Improved cowpea varieties for intercropping** Several new improved breeding lines yielded 50 to 200% higher than local varieties under intercropping with millet and sorghum without insecticide spray.

✎ **Screening method for drought tolerance** A simple method (using a shallow box for seedling screening) has been developed, which discriminates between drought-tolerant and susceptible lines of cowpea. Using this method 2 types of drought-tolerant lines have been identified and their genetics studied.

✎ **Early maize and soybean varieties** The variety trials in 1996 and 1997 have shown that improved extra-early-maturing maize varieties can yield over 3 t/ha in areas with about 600 mm rainfall and early-maturing soybean varieties produce 1.5-2.0 t/ha grain yield without insecticide, rhizobium inoculum, or fertilizer.

✎ **Farmer-to-farmer diffusion of improved cowpea seeds** Preliminary results of IITA/GTZ farmers' participatory seed multiplication and diffusion program have shown great promise for rapid distribution of improved cowpea seeds. Three kg seed of one improved cowpea variety was given on credit to each of 36 selected farmers in June 97. By Dec 97, a total of 6672 kg seed had been produced by these farmers. Similarly, farmer-to-farmer diffusion of an improved cowpea variety has increased cowpea cultivation in the dry season from 1 farmer in 1993 to over 2000 farmers in 1997 in northern Nigeria. Socioeconomic studies revealed that the main driving forces of dry-season cowpea were membership in farmers' association, high grain yield, farm size, and contact with extension workers.

Project 12

Improvement of Maize-Grain Legume Systems in the Moist Savanna of West and Central Africa

Goal

To enhance the productivity of maize grain legume systems in the moist savanna by improving crop varieties and the management of cropping systems

Highlights

✎ The maize breeding team of scientists based in Cote d'Ivoire and Nigeria made significant advances in the development of both open-pollinated varieties and hybrids with higher levels of N-use efficiency (NUE) and drought tolerance. The 10 best families selected from the low N-tolerant pool had 60% higher grain yield under N stress and 19% higher yield under high N than the commercial hybrid. These results confirmed the progress that was reported for this trait in 1996.

✎ The improvement of maize for NUE is expected to have a major impact on both the productivity and sustainability of maize grain legume systems in the moist savanna, with the majority of producers not able to obtain adequate quantities of N fertilizers.

✎ About 200 soybean breeding lines were evaluated for total nitrogen produced through fixation (from the atmosphere). This has enabled selection of breeding lines that produce high grain yield and also contribute nitrogen to the production systems even when the grain is harvested. Several breeding lines have been identified with grain yield equal to the best check and producing 50% to 75% higher fodder yields. These new breeding lines have the potential to considerably improve the productivity and sustainability of the farming system.

✎ Strip cropping maize with improved varieties of cowpea gave encouraging results for both maize and cowpea grain yields with no insecticide spray. Both maize and cowpea were planted at the beginning of the rainy season, as this helps the cowpea to escape the time when the most damaging insects are present in high numbers.

✎ Response of key maize and soybean genotypes could be predicted with simulation models in diverse cropping systems in on station studies. Using models, we can suggest ways for maximizing nutrient use efficiency in various agroecologies. Initial on-farm experimentation and modeling suggest weaknesses in current understanding of physiological processes as well as interactions among N and P nutrients. Regional analysis, combining simulation and geographic information system (GIS), shows large areas of potential impact for soybean-maize rotational systems in West Africa. Models and GIS tools are being combined to develop a decision-support system that can be used by NARS to help identify the appropriate maize-grain legume systems for testing and dissemination.

Project 13

Improvement of Yam-based Systems

Goal

To ensure that farmers achieve a sustainable increase in the productivity of yam based production systems through adoption of improved technologies. The project develops relevant technologies targeted at enhanced productivity of such systems, in partnership with NARES

Highlights

✎ Gender implications for the development of resource management technologies for yam production were investigated in a survey of more than 600 yam growers in the southern Guinea savanna of Nigeria. Women were observed to be heavily involved in yam production, contrary to conventional belief. For both sexes, yam was mainly grown in newly cleared land from long fallow and for a continuous period of less than 3 years. The major constraints in yam production for both sexes were pests and diseases in the field and during storage. Other constraints were weeds, declining soil fertility, lack of staking materials, and labor. It was concluded that the practice of growing yam first on newly cleared land and avoiding continuous yam cultivation may have more to do with the control of pest and pathogen buildup than with declining soil fertility.

✎ In studies at IITA, Ibadan, *Scutellonema bradys* (yam nematode) caused 60% loss of marketable tubers under low nematode pressure, and 100% loss under high pressure. Both situations led to complete loss of planting material.

✎ A survey of yam viruses in Nigeria revealed the presence of yam mosaic potyvirus (previously reported from Nigeria), *Dioscorea alata* potyvirus and cucumber mosaic cucumovirus (previously reported in yams in West Africa), *Dioscorea alata* badnavirus and *Dioscorea dumetorum* potyvirus (not previously reported from Africa). Three new viruses, tentatively named *Dioscorea* mottle, *Dioscorea* mild chlorotic and *Dioscorea* necrosis viruses, have been isolated from their natural *Dioscorea* spp. hosts. Diagnostic techniques have been developed for all yam viruses found to date in Nigeria.

✦ A protocol to ensure the safe international movement of *D alata* germplasm was developed. Consequently, two genotypes of *D alata* were certified for international distribution for the first time. Eleven new genotypes of *D rotundata* were also certified for international distribution.

✦ Over 6,300 virus-tested plantlets representing 22 genotypes of *D rotundata* were delivered to NARS in 7 countries while 6 767 minitubers, produced from virus-tested plantlets of the same species were distributed to NARS in 8 countries.

✦ The development of yams in Uganda got a major boost during the year. More than 140 farmers (including 104 women) from 3 districts were trained in techniques for rapid field/nursery multiplication of yam planting materials. Twenty six varieties of introduced *D rotundata* and 7 local varieties of *D alata* and one of *D cayenensis* are under rapid field multiplication. Over 84% of 3,000 virus tested plantlets of *D rotundata* were successfully established, following a fresh introduction of 22 genotypes from IITA headquarters.

✦ Prerelease varietal trials in Nigeria conducted in partnership with the National Root Crops Research Institute confirmed the superiority of 3 IITA derived hybrid clones of *D rotundata* over popular local varieties used as checks.

Project 14

Cassava Productivity in Lowland and Mid- Altitude Agroecologies of Sub-Saharan Africa

Goal

To develop, evaluate, and promote improved and adapted cassava germplasm for the lowland and mid altitude agroecological zones of sub-Saharan Africa, and to develop agronomic and other practices to ensure sustainable cassava production and utilization. Interaction with national programs is fostered through long established links in West and Central Africa, two root crop research networks (EARRNET and SARRNET) covering East and Southern Africa, and cassava research at the East and Southern Africa Regional Center (ESARC) in Uganda.

Highlights

✦ Efforts to broaden the genetic base of cassava in SSA continued, with the distribution of 497 000 seeds of broad-based improved populations (2,764 families) with multiple pest resistance and special traits to national programs in 21 countries. IITA also received 30,000 seeds derived from landraces of 8 SSA countries and 70,000 seeds from CIAT with adaptation to the semiarid mid-altitude, and subtropical ecologies, respectively.

✦ Improved genotypes incorporating resistant genes from African landraces, combining acceptable agronomic and end-user characteristics with very high levels of multiple resistance to cassava mosaic disease (CMD), bacterial blight (CBB), anthracnose (CAD), and green mite (CGM) have been developed. Forty one of these genotypes were virus tested and certified, and a total of 344 genotypes are now available at IITA for international distribution. Of these materials, IITA distributed 21 040 in vitro plantlets to 9 collaborating countries. In addition, the production and delivery was completed of 18 090 certified cassava in vitro

plantlets to the Angola Seed of Freedom Project through World Vision International

✦ Sourcing seeds from broad based germplasm at IITA Ibadan the EARRNET/ESARC regional germplasm program in Serere, Uganda distributed a total of 1 222 promising clones to the national programs of Rwanda (593 clones), Kenya (555 clones), and Uganda (74 clones) for further evaluation under local conditions

✦ Collaborative work with the International Livestock Research Institute for smallholder crop livestock farming systems showed large variations in root and foliage yields and nutritive value

✦ Early-bulking varieties continue to be identified from IITA germplasm In Nigeria, genotypes adapted to dry-season cropping in the inland valley ecosystem could give average yields of 25 tonnes/ha and dry-matter percentage of 36% in a 6 month period

✦ Nineteen participants were trained on rapid multiplication of cassava and sweetpotato in Rwanda, in an effort to resuscitate research and development of these commodities

Project 15

Molecular and Cellular Biotechnology for Crop Improvement

Goal

To advance the efficacy of genetic improvement and germplasm dissemination beyond the norms associated with the application of conventional breeding and diagnostic techniques The project makes new molecular and cellular tools and products available to collaborating scientists working on IITA's mandated crops

Highlights

✦ Roots and plantlets were obtained from 3-week and 7-week-old immature seeds, respectively, in *D. alata*

✦ Transgenic cowpea lines were produced with constructs containing *Gus* reporter genes, *Bacillus thuringiensis* genes coding for the *Cry* IAb and *Cry* IAc insecticidal crystal endotoxin The transgenics are being characterized and evaluated for insect resistance

✦ Putative genomic and cDNA fragments corresponding to lectins, thionin phenylalanine ammonia lyase (PAL), and chalcone synthase (CHS) genes associated with pest resistance have been isolated from African yam bean, wild cowpea and/or cultivated cowpea Most of the fragments correspond to the predicted size

✦ Affinity chromatographic procedures have been used to purify lectins from African yam bean and *V. vexillata*

✦ Six mapping populations from controlled hybridizations between selected yam genotypes of *D. rotundata* and *D. alata* were developed for use in developing linkage maps The parental lines used for generating the mapping populations were selected for their contrasting phenotypic expressions for virus and nematode resistance in *D. rotundata* and for anthracnose in *D. alata*

✦ Recombinant inbred (RI) lines developed for the genome mapping of cowpea are now available Some DNA markers have been generated and placed on the map which presently spans 665 cM

indicating an average map distance of 7.2 cM. Additional DNA markers are being generated for placement on the map.

✦ Some markers associated with quantitative trait loci (QTL) in cowpea such as days to flowering, 100 seed weight, and days to pod maturity were identified, among others.

✦ Candidate microsatellite markers for parthenocarp, earliness, and regulated suckering in plantain have been identified and will be tested further.

✦ IITA, in collaboration with John Innes Centre and Horticulture Research Institute (UK), developed reliable detection systems for *Dioscorea alata* badnavirus, facilitating safe movement of improved *D. alata* germplasm.

✦ Immunocapture reverse transcriptase polymerase chain reaction (IC-RT-PCR), a very sensitive protocol, was developed to detect yam mosaic potyvirus.

✦ Scottish Crops Research Institute (Dundee, UK) donated to IITA six hybridoma cell lines raised against cassava mosaic geminiviruses. Ascetic and culture fluids were produced at IITA and distributed to 10 laboratories in Africa, thereby enhancing the capability of national programs to diagnose geminiviruses in cassava.

Project 16

Conservation and Genetic Enhancement of Plant Biodiversity

Goal

To improve the conservation and utilization of plant biodiversity to promote sustainable food production in sub-Saharan Africa.

Highlights

✦ Field collections were made of 330 local cassava accessions from Togo, 97 local yam cultivars from Cote d'Ivoire, and 11 wild yam and 25 wild *Vigna* spp. from Nigeria. A total of 331 accessions of yam and 44 accessions of cassava was transferred from the field genebank to in vitro cultures. A large number of indigenous yam cultivars (557) was collected in Benin Republic. A total of 388 seedlots of herbaceous legumes were identified and cataloged, by combining former ILRI and IITA collections, 171 seedlots were distributed for experiments in IITA/ILRI (42) and to NARS (74) and NGOs (55).

✦ User-friendly database management systems for over 37,000 germplasm accessions of cowpea, wild *Vigna* spp., Bambara groundnut, soybean, rice, yams, and cassava were developed.

Genetic diversity and heterotic groups in Guinea yams (*D. rotundata* and *D. cayenensis*) and their wild relative *D. praehensilis* were assessed, using amplified fragment length polymorphisms (AFLPs). Molecular markers were used for the first time to characterize landrace collections of plantains.

✦ Many improved breeding lines (431) and African landraces (561) of cassava are being evaluated and characterized for agrobotanical traits, and resistance to cassava mosaic disease (ACMD), cassava bacterial blight (CBB), and cassava anthracnose disease (CAD).

Project 17

Improving the Dissemination of IITA Research Results

✍ From a study of combining ability of *Musa* hybrids, 18 promising diploid and 6 triploid banana hybrids were identified with large bunch size, large number of hands and fruits, big fruit size, improved ratooning, earliness, and resistance to black sigatoka. A promising diploid hybrid with resistance to nematodes was identified.

✍ Maize varieties were extracted from TZL COMP4 and TZE COMP3 and were higher yielding than varieties extracted from previous cycles of selection. Two reciprocal populations formed for the mid altitudes were screened for resistance to maize streak virus *Eusserohilum turcicum* and ear rot.

✍ Four maize lines from TZPB-SR prol. C1 had over 90% prolific (2 eared) plants as well as high yield and disease resistance in the forest zone of Nigeria. Across the forest zone and savanna sites, the best line had an average yield of 6.5 t/ha, with 70% prolific plants. S₂ and S₄ testcrosses from two white maize populations (IWD and IWF) and their STR versions produced over 50% higher yield than the commercial hybrid Oba Super 1.

✍ Some inbred lines were identified (e.g., 1102, 1309) with levels of resistance to *S. hermonthica* comparable to or better than that of inbred 1368. Hybrids generated from these lines were high yielding and resistant to abiotic stresses.

✍ Some 25-30 S₁ families derived from early-maturing maize populations (DR-W Pool and DR-Y Pool) with drought tolerance and resistance to maize streak virus were selected for recombination.

Goal

To strengthen the effectiveness of NARS in the generation and utilization of appropriate research results. Major activities are human resource development, project development and monitoring, technology transfer among NARS, information dissemination, and public awareness generation.

Highlights

✍ Group training courses were held both at IITA and in several countries on matters such as rapid multiplication of cassava and sweetpotato, banana and plantain research, biometric applications, soil and plant analysis.

✍ IITA played a lead role in intercenter collaboration on training. A meeting involving NARS and IARCs was held at IITA in November 1997. The intercenter training group is now called the IARCs and NARS Training Group (INTG).

✍ The Technology Testing & Transfer Unit initiated research support in various countries in biological control of *Striga hermonthica*, mango mealybug, and water hyacinth.

✍ As for training materials, ten IITA Research Guides (IRGs) were produced during 1997. Fifteen IRGs were distributed to 125 addresses in several countries. Collaboration on the production of training materials took place between IITA and Ghana, Cote d'Ivoire, Zambia, CIMMYT, and ICRISAT.

✎ Information was generated and disseminated for research purposes and for public awareness generation. There has been a substantial increase in the volume of publications available in French.

✎ The Library continued to provide assistance to a wide range of scientists and students.

✎ Public awareness is an important aspect of the corporate activities of IITA. IITA's research results and their impact have been presented in various fora. Information has been routinely disseminated through news releases, weekly articles in newspapers, CGIAR publications, international and local radio.

✎ EARRNET, SARRNET, and WECAMAN, the EU OFAR, TT&TU, and the GTZ-supported Seed Project continued to provide a strong link between IITA and NARS in technology transfer.

✎ IITA-NARS interaction was further strengthened by IITA's lead role in preparing a USAID/OFDA supported strategic plan to mitigate the effects of the El Niño weather disturbances using cassava and sweetpotato, for countries in East and Southern Africa.

✎ IITA and CORAF held a meeting to discuss working relations and the harmonization of research within the West/Central African subregion. It resulted in an agreement to conduct regional research under the umbrella of CORAF, and to harmonize research networks on maize and on roots and tuber crops.

Project 18

Ecoregional Program for the Humid and Subhumid Tropics of Africa

Goal

To assist smallholder and medium scale farmers to improve their well being and alleviate poverty through the use of sustainable production technologies and postharvest systems that increase productivity and food security and minimize natural resources degradation.

Highlights

✎ The 1997 EPHTA work plan which focused on benchmark area development and pilot studies was endorsed by the Program Committee (PC) at its first meeting held at IITA, Ibadan, 24-26 Feb 1997.

✎ Benchmark area development. All 5 countries hosting the 6 benchmark areas appointed Benchmark Area Coordinators. Several meetings were held in all benchmark areas to sensitize stakeholders and potential EPHTA partners.

✎ Characterization surveys. Surveys have been completed in the forest margins benchmark area in Cameroon and in the northern Guinea savanna and the degraded forest benchmark areas in Nigeria.

✎ The forest margins benchmark area in Cameroon was officially launched on 26 May 1997 in Yaounde, as was the degraded forest benchmark area in southeastern Nigeria on 6 Oct 1997 at Owerri.

✦ Nominations have been received from countries for membership of the 7 working groups indicated in the program document. The working groups will be formed and become operational in 1998.

✦ From 12 to 16 May 1997 a workshop on ecoregional research methodology was held in Cameroon. It was attended by 3 participants from each of the 6 benchmark areas, including the Benchmark Area Coordinators and each of the Pilot Site Coordinators. The workshop was also attended by a 5-man delegation from CIRAD and representatives from ILRI, WARDA, and IITA, the organizing center.

✦ From 15 to 19 Sep 1997, EPHTA, in collaboration with the Systemwide Programs for Integrated Pest Management (SP-IPM) and Alternatives to Slash-and-Burn (ASB) sponsored the Integrated Weed Management Workshop, which was held in Yaounde, Cameroon. It was attended by participants from EPHTA member countries, representatives from WARDA, CIRAD, NRI, and IIBC (Britain), and IITA.

Project 19

Systemwide Program on Integrated Pest Management

Goal

To ensure, by encouraging better communication, coordination, and the adoption of more effective approaches, that CG research on integrated pest management (IPM) is more responsive to the needs of IPM practitioners, gains wider recognition, and thereby contributes fully to sustainable agricultural development.

Highlights

✦ Researchers from many national programs and several international centers participating in the intercenter project on whiteflies and geminiviruses have begun field work at numerous sites in Africa and Latin America with close support from specialist laboratories in Europe and North America. This technically and organizationally complex project is the first to be launched under the auspices of the Systemwide Program on IPM. Its successful establishment, under the leadership of CIAT, will serve as a model for future project development.

✦ In this first, diagnostic phase of the project, researchers in the different regions will use a common set of methodologies to assess the importance of losses associated with different combinations of crops, viruses, and their whitefly vectors. The knowledge gained will provide a sound basis for prioritizing future research efforts.

✦ In Africa, recognizing the urgency of the situation resulting from a highly destructive outbreak of African cassava mosaic disease, researchers involved in the project are moving rapidly to carry out analytical research and implement appropriate counter-measures. In this regard, the network of collaborators and sound knowledge base already established by ESCaPP and IITA's other root crop research and development efforts have been an important factor in facilitating the timely launch of project activities.

✦ A workshop on management of weeds in the farming system, convened at IITA's Humid Forest Station in September, pioneered a new mode of collaboration between the Institute and various inter-institutional initiatives. The workshop cosponsored by the SP-IPM,

Research Highlights

EPHTA (both of which are led by IITA) and ASB (the inter-center initiative on Alternatives to Slash-and-Burn led by ICRAF) was co-convened by IITA and WARDA. Weed scientists from national programs across West and Central Africa came together with disciplinary specialists from the involved centers and from various institutions in Europe to analyze weed problems encountered in the region, examine options for tackling them, and outline plans to carry out the necessary research. Several detailed research proposals are now being prepared within the framework of EPHTA, with plans for pilot studies at various benchmark sites in the different agroecological zones.

✈ Other meetings of intercenter task forces were held, focusing on specific topics. When available CGIAR information is brought together into an authoritative document during 1998, it will provide a more reliable basis for researching and implementing sustainable pest management solutions.




Graduate Research Completed at IITA in 1997

Degree	Country	Division	University	Sponsor(s) Network	Name	M/F	Research topic
Crop Improvement							
PhD	Nigeria	CID	University of Ibadan, Nigeria	IITA, Self	Daniel I	M	Storage of seeds and pollen grains of yams (<i>Dioscorea</i> spp.) for germplasm conservation and enhancement in crop improvement
PhD	Nigeria	CID	University of Ibadan Nigeria	Self	Akparobi S O	M	Response of cassava (<i>Maniot esculenta</i> Crantz) clones to temperature regimes
PhD	Sierra Leone	CID	Njala University Sierra Leone	IITA	Lahri T M	M	Comparative studies on yield physiology and quality aspects of cassava (<i>Maniot esculenta</i> Crantz) grown in the lowland and upland
PhD	Nigeria	CID	University of Jos Nigeria	Self	Abimiku S E	M	Genetics of resistance to <i>Ascochyta</i> blight disease of cowpea
PhD	Nigeria	CID	University of Jos Nigeria	Self	Yulwa, V	F	Genetics of plant type and branching habit in cowpea
PhD	Cameroon	CID	Obafemi Awolowo University, Ile Ife Nigeria	Self	Hankoua, B B	M	Micropropagation and diosgenin production of <i>Dioscorea dumetorum</i>
PhD	Nigeria	CID	University of Ibadan Nigeria	Italian grant	Ikea J	M	Investigating the transformability of cowpea by particle bombardment
MSc	Belgium	CID	Katholieke Universiteit Leuven	SP	De Beule Hilde	F	Evaluation of the damage by nematodes to the size of the root system of plants at flowering stage
MSc	Uganda	CID	Makerere University Uganda	IITA/ESARC	Bukenya C	M	Utilization of cassava in a rural economy with specific reference to processing and marketing of cassava based products
MSc	Benin	CID	Ahmadu Bello University Zaria, Nigeria	IITA/FF	Adjadi O	F	Genetic variability in cowpea, <i>Vigna unguiculata</i> Walp under various crop management systems
MSc	Nigeria	CID	University of Ibadan Nigeria	Self	Egesi Ngozi C	M	Taxonomy of the African wild <i>Vigna</i> gene pool <i>Vigna luteola</i> Benth and <i>Vigna marina</i> Merrill
MSc	Nigeria	CID	University of Ibadan Nigeria	Self	Ombokun A	F	Control of bacterial contaminations in yam cultures use of antibiotics
Plant Health Management							
PhD	Uganda	PHMD	University of Reading UK	RF	Tusheremeirwe W	M	Assessment of banana pathogens in Uganda with emphasis on factors favoring development of black sigatoka
PhD	Nigeria	PHMD	University of Reading UK	IITA	Olatunwo, R.	M	Incidence of <i>Stenocarpella macrospora</i> on maize and evaluation of potential management strategies
PhD	Entrea	PHMD	Georg August University Göttingen Germany	BMZ	Fessehaie A	M	Development of diagnostic methods for cassava bacterial blight
PhD	Gambia	PHMD	University of Reading UK	IIBC/DFPV	Sidi S	M	Synergistic effect of the tungus <i>Metarhizium flavoviride</i> with pyrethroid and insect growth regulator insecticides for the control of grasshoppers and locusts
PhD	Germany	PHMD	Humboldt University Berlin, Germany	BMZ/IITA	Lorek, C	M	<i>Prostephanus</i> fungal antagonists Honduras and Benin


Degree	Country	Division Network	University	Sponsor(s)	Name	M/ F	Research topic
MSc	Senegal	PHMD	University of Ghana, Legon	IITA/FF	Diop K.	F	The biology of meles <i>Ceranisus</i> (Walker) (Hym. Ghana Eulophidae), a parasitoid of the bean flower thrips <i>Megalurothrips sjostedti</i> Trybom (Thys., Thripidae)
MSc	Nigeria	PHMD	University of Ibadan Nigeria	Self	Odu B	M	Characterization of a virus potyvirus in <i>Dioscorea alata</i>
MSc	Cameroon	PHMD	University of Nigeria Nsukka	IITA	Abang, M M	M	Characterization of populations of <i>Colletotrichum gloeosporioides</i> (Penz.) from yam based cropping systems in Nigeria
MSc	Uganda	PHMD	Makerere University Uganda	RF	Abera A M	F	Timing of attack by the banana weevil and damage distribution in the host plant
Resource and Crop Management							
PhD	Nigeria	RCMD	University of Nigeria Nsukka	GTZ/Self	Asawalam O D	M	The role of earthworms in sustaining soil fertility and crop production under subhumid and humid tropical conditions
PhD	Nigeria	RCMD	University of Ibadan Nigeria	IITA/Self	Onvia, N	F	Movement and persistence of imazapyr in soils of the moist savanna zone in relation to control of <i>Imperata cylindrica</i> (speargrass)
PhD	Cameroon	RCMD	University of Georgia, USA	IITA	Elad R. L	F	The implicit valuation of resources within the African agricultural household
PhD	Nigeria	RCMD	University of Hohenheim Germany	IITA	Ndubuisi A	M	Economic analysis of crop livestock integration among the agropastoralists in the northern Guinea savanna of Nigeria
PhD	Ghana	RCMD	University of Saskatchewan, Saskatoon, Canada	IITA	Gana, B K	F	Evaluating the effects of land use changes on soil quality in northeastern Ghana
PhD	Nigeria	RCMD	University of Ibadan, Nigeria	Self	Modupe V O	M	Soybeans phosphorus requirements and production efficiency in the humid rain forest region of Nigeria using simulation modelling
PhD	Nigeria	RCMD	Federal University of Technology Owerri Nigeria	Self	Asumugha G N	M	Efficiency in the marketing system for ginger (a root crop) compared to cassava in Nigeria
PhD	Nigeria	RCMD	Katholieke Universiteit Leuven	KUL/SOM	Ibewro B E	M	Nitrogen availability in relation to SOM dynamics under <i>Mucuna</i> and <i>Lablab</i> cover cropping compared to <i>Imperata</i> systems
PhD	Nigeria	RCMD	Federal University of Technology, Akure, Nigeria	Self	Epebun Cofe F F	F	Evaluation of the soil surface change of characteristics in the humid tropical zone of Nigeria
PhD	Nigeria	RCMD	University of Ibadan Nigeria	Self	Awoyemi T T	M	Gender role and productivity differential in cassava business
MSc	Belgium	RCMD	Katholieke Universiteit Leuven	BNMSProject	Delaure S	M	Using soil information for the characterization and selection of research domains and representative farmer fields in Zana Kaduna benchmark fields in northern Nigeria

Financial Statement

Core costs by operating segment, 1997 and 1996



Research expenditure by CGIAR program, 1997 and 1996



Note: The core budget is used to fund those research-related activities essential in meeting the CGIAR objectives for developing countries

**Statement of Financial
Position**

at 31 December
Expressed in US\$ thousands

	1997	1996
ASSETS		
Current assets		
Cash and cash equivalents	20,090	19,292
Accounts receivable		
Donors	3,287	3,488
Others	861	804
Inventories	1,249	1,444
Prepaid expenses	490	685
Other assets	<u>189</u>	<u>178</u>
Total current assets	<u>26,166</u>	<u>25,891</u>
Fixed assets		
Property plant and equipment	74,037	71,489
Less accumulated depreciation	<u>(41,290)</u>	<u>(39,926)</u>
Total fixed assets—net	<u>32,747</u>	<u>31,563</u>
Total assets	<u>58,913</u>	<u>57,454</u>
LIABILITIES AND FUND BALANCES		
Current liabilities		
Accounts payable and other liabilities	4,656	4,089
Accrued salaries and benefits	4,959	5,338
Payments in advance—donors	<u>8,272</u>	<u>7,304</u>
Total current liabilities	<u>17,887</u>	<u>16,731</u>
Net assets		
Capital invested in fixed assets	32,747	31,563
Capital fund	2,758	3,722
Operating fund	<u>5,521</u>	<u>5,438</u>
Total net assets	<u>41,026</u>	<u>40,723</u>
Total liabilities and net assets	<u>58,913</u>	<u>57,454</u>

Statement of Activity
at 31 December
Expressed in US\$ thousands

	1997	1996
REVENUE		
Grants	29,170	31,180
Investment income	<u>1,474</u>	<u>1,004</u>
Total revenue	<u>30,644</u>	<u>32,184</u>
EXPENSES		
Research programs	20,713	20,801
Conferences and training	2,403	2,639
Information services	574	793
General administration	1,633	2,529
General operations	2,873	3,477
Depreciation	<u>3,424</u>	<u>3,714</u>
Total expenses	<u>31,620</u>	<u>33,953</u>
Excess of expenses over revenue	<u>(976)</u>	<u>(1,769)</u>

	1997	1996	Statement of Cash Flow at 31 December Expressed in US\$ thousands
Cash flows from operating activities			
Excess of expenses over revenue	(976)	(1,769)	
Adjustments to reconcile net cash Provided by operating activities			
Depreciation	3,424	3 714	
Gain on disposal of assets	267	474	
Decrease (Increase) in assets			
Accounts receivable—donors	201	(555)	
Accounts receivable—others	(57)	331	
Inventories	195	(116)	
Prepaid expenses	195	(442)	
Other assets	(11)	(47)	
Increase (Decrease) in liabilities			
Accounts payable and other liabilities	567	(236)	
Accrued salaries and benefits	(379)	498	
Payments in advance—donors	968	(386)	
Total adjustments	<u>5,370</u>	<u>3,235</u>	
Net cash provided by operating activities	4,394	1 466	
Cash flow used in investment activities			
Acquisition of fixed assets	(3,596)	(2,914)	
Net increase (decrease) in cash and cash equivalents	<u>798</u>	<u>(1,448)</u>	
Cash and cash equivalents End of year	20,090	19 292	
Beginning of year	19,292	20,740	
Increase (Decrease) in the year	<u>798</u>	<u>(1,448)</u>	

Donors	1997		1996		Donors Expressed in US\$ thousands
	Core Funding	Complementary Funding	Core Funding	Complementary Funding	
Austria	401	84	150	1 501	
Belgium	1,166	-	1 253	297	
BMZ Germany	1,412	86	1 776	93	
Canada	847	-	911	-	
Commission of the European Communities	766	-	202	130	
Common Fund for Commodities	81	-	-	-	
Denmark	1,370	57	1 668	37	
Food and Agriculture Organization	-	1	-	2	
Ford Foundation	-	4	-	83	
France	227	-	310	334	
Gatsby Charitable Foundation	34	200	-	106	
International Centre for Research in Agroforestry	-	69	-	97	
International Development Research Centre	61	207	-	251	
International Fund for Agricultural Development	258	186	136	110	
International Institute of Biological Control	-	700	-	505	
Italy	400	-	300	-	
Japan	3,400	-	4 281	-	
Korea Republic of	50	-	50	-	
Netherlands	1,232	-	1 171	-	
Nigeria	47	12	6	-	
Norway	1,147	-	665	-	
Overseas Development Administration (UK)	-	94	-	65	
Rockefeller Foundation	281	129	349	94	
Sasakawa Africa Association	123	-	-	128	
Sweden	310	-	313	-	
Switzerland	749	546	558	577	
Thailand	50	-	-	-	
United Kingdom	556	-	723	-	
United Nations Development Programme	767	48	128	985	
United States Agency for International Development	4,977	1,347	3 326	3 025	
United States Department of Agriculture	-	118	-	-	
University of Hohenheim	-	108	-	114	
World Bank	3,447	-	3 800	90	
Multiple/Other Donors	884	125	302	161	
Closed Projects	-	6	-	17	
TOTAL	<u>25,043</u>	<u>4,127</u>	<u>22,378</u>	<u>8,802</u>	Note The National Fertilizer Company of Nigeria (NAFCON) provides fertilizer from time to time as their stock allows



Contributions by IITA staff to scientific literature that became available during 1997 including journal articles, book chapters, papers in monographs or conference proceedings, training materials, published abstracts and research notes or disease reports Also included are (1) publications based on work done by IITA staff prior to their joining IITA especially where the work reported is of interest to IITA (2) publications by staff who have left IITA which are based on work done while they were at the Institute

Journal Articles

Ahiabu, R K A , K E Danso, G Y P Klu, P Bieler, and W Msikita 1997

In vitro cassava research capability at the Biotechnology and Nuclear Agricultural Institute (BNARI)—Ghana Potentials and constraints African Journal of Root and Tuber Crops 2 209–212

Akano, A O , G I Atiri, S Y C Ng, and R Asiedu 1997

Effect of African cassava mosaic disease on growth and yield components of virus tested cassava genotypes derived from meristem culture in early and late planting periods in three agroecologies of Nigeria African Journal of Root and Tuber Crops 2 44–48

Asadu, C L A , J Diels, and B Vanlauwe 1997

A comparison of the contributions of clay silt and organic matter to the effective CEC of soils of sub Saharan Africa Soil Science 162 785–794

Asanzi, M C , N A Bosque Perez, and L R Nault 1995 (publ in 1997)

Movement of *Cicadulina storeyi* [Homoptera Cicadellidae] in maize fields and its behavior in relation to maize growth stage Insect Science and its Application 16 39–44

Beare, M H , M Vikram Reddy, G Tian, and S C Srivastava 1997

Agricultural intensification soil biodiversity and agroecosystem function in the tropics the role of decomposer biota Applied Soil Ecology 6 87–108

Berner, D K , F O Ikie, and J M Green 1997

ALS inhibiting herbicide seed treatments control *Striga hermonthica* in ALS modified corn (*Zea mays*) Weed Technology 11 704–707

Borgemeister, C , W G Merkle, C Adda, P Degbey, and R H Markham 1997

Seasonal and meteorological factors influencing the annual flight cycle of *Prostephanus truncatus* (Coleoptera Bostrichidae) and its predator *Teretriosoma nigrescens* (Coleoptera Histeridae) in Benin Bulletin of Entomological Research 87 239–246

Bridge, P D , C Prior, J Sagbohan, C J Lomer, M Carey, and A Buddie 1997

Molecular characterisation of isolates of *Metarhizium* from locusts and grasshoppers Biodiversity and Conservation 6 177–189

Buresh, R J , and G Tian 1997

Soil improvement in sub Saharan Africa Agroforestry Systems 38 51–76

Cardwell, K F , and T Wehrly 1997

A nonparametric significance test for combating crop disease Biometrics 53 207–218


Cardwell, K , F Schulthess, R Ndemah, and Z Ngoko 1997

A systems approach to assess crop health and maize yield losses due to pests and diseases in Cameroon Agriculture Ecosystems and Environment 65 33–47

Cardwell, K F , J Kling, and C Bock 1997

Comparison of field inoculation methods for screening maize against downy mildew (*Peronosclerospora sorghi*) Plant Breeding 116 221–226

- Carsky, R J , R Abaidoo, K Dashiell, and N Sanginga 1997**
Effect of soybean on subsequent maize grain yield in the Guinea savanna zone of West Africa African Crop Science Journal 5 31–38
- Chabi Olaye, A , F Schulthess, T G Shanower, and N A Bosque Perez 1997**
Factors influencing the developmental rates and reproductive potentials of *T busseolae* (Gahan) (Hymn Scelionidae) an egg parasitoid of *S calamistis* Hampson (Lep Noctuidae) Biological Control 8 15–21
- Chukwumah, Y , S Y C Ng, and M Bokanga 1997**
Genotypic and tissue differences in biosynthetic activity for linamarin production in cassava African Journal of Root and Tuber Crops 2 73–76
- Craenen, K , and R Ortiz 1997**
Effect of the bs1 gene in plantain–banana hybrids on response to black sigatoka Theoretical and Applied Genetics 95 497–505
- Dansi, A , J Zoundjhekon, H D Mignouna, and F M Quin 1997**
Collecte des ignames cultivees du complexe *Dioscorea cayenensis–rotundata* au Benin Plant Genetic Resources Newsletter 112 81–85
- Defoer, T , A Kamara, and H De Groote 1997**
Gender and variety selection Farmers assessment of local corn varieties in southern Mali African Crop Science Journal 5 65–76
- Dixon, A G O , and E N Nukenine 1997**
Statistical analysis of cassava yield trials with the additive main effects and multiplicative interaction (AMMI) model African Journal of Root and Tuber Crops 3 46–50
- Douro Kpindou, O K , P A Shah, J Langewald, C J Lomer, H van der Pau, A Sidibe, and C O Daffe 1997**
Essais sur l'utilisation d un biopesticide (*Metarhizium flavoviride*) pour le controle des sauteriaux au Mali de 1992 a 1994 Journal of Applied Entomology 121 285–291
- Fargues, J , M S Goettel, N Smits, A Ouedraogo, C Vidal, L A Lacey, C J Lomer, and M Rougier 1996**
Variability in susceptibility to simulated sunlight of conidia among isolates of entomopathogenic hyphomycetes Mycopathologia 135 171–181
- Fargues, J , A Ouedraogo, M S Goettel, and C J Lomer 1997**
Effect of temperature humidity and inoculation method on susceptibility of *Schistocerca gregaria* to *Metarhizium flavoviride* Biocontrol Science and Technology 7 345–356
- Ferris, R S B , R Ortiz, U Chukwu, Y O Akalumhe, S Akele, A Ubi, and D Vuylsteke 1997**
The introduction and market potential of exotic black sigatoka resistant cooking banana cultivars in West Africa Quarterly Journal of International Agriculture 36 141–152
- Gold, C S , and M I Bagabe 1997**
Banana weevil *Cosmopolites sordidus* Germar (Coleoptera Curculionidae) infestation of cooking and beer bananas in adjacent stands in Uganda African Entomologist 5 103–108
- Hailemichael, Y , F Schulthess, J W Smith Jr , and W A Overholt 1997**
Suitability of West African gramineous stemborers for the development of *Cotesia* species (Hymenoptera Brachonidae) Insect Science and its Application 17 89–95
- Hall, A E , B B Singh, and J D Ehlers 1997**
Cowpea breeding Plant Breeding Reviews 15 215–274
- Hayashi, Y , and R J Carsky 1997**
Effect of use of savanna tree leaves as mulch materials on germination and growth of selected annual crop seedlings Japanese Journal of Tropical Agriculture 41 14–21
- Inaizumi, H 1997**
Determinants of the factors of the expansion of improved cassava processing technologies in West African countries Journal of Agricultural Development Studies (Japan) 8 40–51
- Inaizumi, H 1997**
Importance of cassava production in Africa and application of COSCA database Journal of Rural Community Studies (Japan) 85 93–99 (in Japanese)
- Kadiata, B D , K Mulongoy, and N O Isirimah 1997**
Influence of pruning frequency of *Albizia lebbek* *Gliricidia sepium* and *Leucaena leucocephala* on nodulation and potential nitrogen fixation Biology and Fertility of Soils 24 255–260



- Kang, B T , F K Salako, I O Akobundu, J L Pleysier, and J N Chianu 1997**
Amelioration of a degraded Oxic Paleustalf by leguminous and natural fallows Soil Use and Management 13 130–136
- Kim, S K 1996**
Horizontal resistance core to a research breakthrough to combat *Striga* in Africa Integrated Pest Management Reviews 1 229–249
- Kim, S K , and V O Adetimirin 1997**
Responses of tolerant and susceptible maize varieties to timing and rate of nitrogen under *Striga hermonthica* infestation Agronomy Journal 89 38–44
- Kim, S K , and V O Adetimirin 1997**
Striga hermonthica seed inoculum rate effects on maize hybrid tolerance and susceptibility expression Crop Science 37 1066–1071
- Kim, S K , V O Adetimirin, and A Y Akintunde 1997**
Nitrogen effects on *Striga hermonthica* infestation grain yield and agronomic traits of tolerant and susceptible maize hybrids Crop Science 37 711–716
- Kim, S K , S T O Lagoke, and C The 1997**
Observations on field infection by witchweed (*Striga* species) on maize in West and Central Africa International Journal of Pest Management 43 113–121
- Kolawole, G O , and B T Kang 1997**
Effect of seed size and phosphorus fertilization on growth of selected legumes Communications in Soil Science and Plant Analysis 28 1223–1235
- Kooyman, C , and I Godonou 1997**
Infection of *Schistocerca gregaria* (Orthoptera Acrididae) hoppers by *Metarhizium flavoviride* (Deuteromycotina Hyphomycetes) conidia in an oil formulation applied under desert conditions Bulletin of Entomological Research 87 105–107
- Kooyman, C , R P Bateman, J Langewald, C J Lomer, Z Ouambama, and M B Thomas 1997**
Operational scale application of entomopathogenic fungi for control of Sahelian grasshoppers Proceedings of the Royal Society B 264 541–546
- Langewald, J , C Kooyman, O K Douro Kpindou, C J Lomer, A Dahmoud, and H Mohamed 1997**
Field treatment of desert locust (*Schistocerca gregaria* Forskal) hoppers in the field in Mauritania with an oil formulation of the entomopathogenic fungus *Metarhizium flavoviride* Biocontrol Science and Technology 7 603–611
- Langewald, J , M B Thomas, C J Lomer, and O K Douro Kpindou 1997**
Use of *Metarhizium flavoviride* for control of *Zonocerus variegatus* A model relating mortality in caged field samples with disease development in the field Entomologia Experimentalis et Applicata 82 1–8
- Lomer, C J , C Prior, and C Kooyman 1997**
Development of *Metarhizium* spp for the control of locusts and grasshoppers Memoirs of the Entomological Society of Canada 171 265–286
- Lomer, C J , M B Thomas, I Godonou, P A Shah, O K Douro Kpindou, and J Langewald 1997**
Control of grasshoppers particularly *Hieroglyphus daganensis* in northern Benin using *Metarhizium flavoviride* Memoirs of the Entomological Society of Canada 171 301–311
- Manyong, V M , and B Oyewole 1997**
Spatial patterns of biological constraints to cassava and yam production in West and Central Africa Implications for technology development and transfer African Journal of Root and Tuber Crops 3 15–21
- Mba, R E C , and A G O Dixon 1997**
Heritability estimates for ACMD resistance for some newly developed cassava clones in Nigeria African Journal of Root and Tuber Crops 2 56–58

- Meikle, W G , A P Gutierrez, and H R Herren 1996**
Decision making by smallholder farmers in an oil palm based economy in southern Benin West Africa Tropical Agriculture (Trinidad) 73 217–225
- Mignouna, H D , and A G O Dixon 1997**
Genetic relationships among cassava clones with varying levels of resistance to African mosaic disease using RAPD markers African Journal of Root and Tuber Crops 2 28–32
- Mignouna, H D , F Camara, and S Kourouma 1997**
Collecting germplasm of wild and cultivated yam (*Dioscorea* spp) in Guinea Plant Genetic Resources Newsletter 109 15–16
- Mobambo, K N , C Pasberg Gauhl, F Gauhl, and K Zuofa 1997**
Host response to black sigatoka in *Musa* germplasm of different ages under natural inoculation conditions Crop Protection 16 359–363
- Moore, D , J Langewald, and F Obognon 1997**
Effects of rehydration on the conidial viability of *Metarhizium flavoviride* mycopesticide formulations Biocontrol Science and Technology 7 87–94
- Msikita, W , J S Yaninek, M Ahounou, H Baimey, and R Fagbemissi 1997**
A system to screen and select for resistance to *Fusarium solani* African Journal of Root and Tuber Crops 2 59–61
- Muller, R , C Pasberg Gauhl, F Gauhl, J Ramser, and G Kahl 1997**
Oligonucleotide fingerprinting detects genetic variability at different levels in Nigerian *Mycosphaerella fijiensis* Journal of Phytopathology 145 25–30
- Ng, S Y C , and I J Ekanayake 1997**
Responses of cassava to polyethylene glycol mediated osmotic stress in vitro African Journal of Root and Tuber Crops 2 228–231
- Ng, S Y C , and N Q Ng 1997**
Cassava in vitro germplasm management at the International Institute of Tropical Agriculture African Journal of Root and Tuber Crops 2 232–234
- Oduor, G I , J S Yaninek, G J de Moraes, and L P S van der Geest 1997**
The effect of pathogen dosage on the pathogenicity of *Neozygites floridana* (Zygomycetes Entomophthorales) to *Mononychellus tanajoa* (Acari Tetranychidae) Journal of Invertebrate Pathology 70 127–130
- Oduor, G I , M W Sabelis, R Lingeman, G J de Moraes, and J S Yaninek 1997**
Modelling fungal (*Neozygites* cf *floridana*) in local populations of cassava green mites (*Mononychellus tanajoa*) Experimental and Applied Acarology 21 485–506
- Ogbe, F O , G Thottappilly, and F M Quin 1997**
Implementation in Africa of serological diagnostic test for cassava mosaic geminiviruses African Journal of Root and Tuber Crops 2 33–36
- Ortiz, R 1997**
Genetic and phenotypic correlations in plantain–banana euploid hybrids Plant Breeding 116 487–491
- Ortiz, R 1997**
Morphological variation in *Musa* germplasm Genetic Resources and Crop Evolution 44 393–404
- Ortiz, R 1997**
Occurrence and inheritance of 2n pollen in *Musa* Annals of Botany 79 449–453
- Ortiz, R 1997**
Secondary polyploids heterosis and evolutionary crop breeding for further improvement of the plantain and banana genome Theoretical and Applied Genetics 94 1113–1120
- Ortiz, R , and J H Crouch 1997**
The efficiency of natural and artificial pollinators in plantain (*Musa* spp AAB group) hybridisation and seed production Annals of Botany 80 693–695
- Ortiz, R , and H Langie 1997**
Path analysis and ideotypes for plantain breeding Agronomy Journal 89 988–994
- Ortiz, R , and D Vuylsteke 1997**
Improved polyploid *Musa* germplasm developed through ploidy manipulations African Crop Science Journal 5 107–117
- Ortiz, R , P D Austin, and D Vuylsteke 1997**
IITA High Rainfall Station Twenty years of research for sustainable agriculture in the West African humid forest HortScience 32 969–972

- Ortiz, R , D Vuylsteke, R S B Ferris, J U Okoro, A N Guessan, O B Hemeng, D K Yeboah, K Afreh Nuamah, E K S Ahiekpor, E Foure, B A Adelaja, M Ayodele, O B Arene, F E O Ikiedrugwu, A N Agbor, A N Nwogu, E Okoro, G Kayode, I K Ipimoye, S Akele, and A Lawrence 1997**
Developing new plantain varieties for Africa *Plant Varieties and Seeds* 10 39–57
- Ortiz, R , K Craenen, and D Vuylsteke 1997**
Ploidy manipulations and genetic markers as tools for analysis of quantitative trait variation in progeny derived from triploid plantains *Hereditas* 126 255–259
- Osuji, J O , B E Okoli, and R Ortiz 1997**
Histochemical localization of calcium oxalate crystals in fruits of plantain and banana cultivars *Fruits* 52 5–10
- Osuji J O , G Harrison, J H Crouch, and J S Heslop Harrison 1997**
Identification of the genomic constitution of *Musa* L lines (bananas plantains and hybrids) using molecular cytogenetics *Annals of Botany* 89 787–793
- Osuji, J O , B E Okoli, D Vuylsteke, and R Ortiz 1997**
Multivariate pattern of quantitative trait variation in triploid banana and plantain cultivars *Scientia Horticulturae* 71 197–202
- Paustian, K , O Andren, H H Janzen, R Lal, P Smith, G Tian, H Tiessen, M Van Noordwijk, and P L Woome 1997**
Agricultural soils as a sink to mitigate CO₂ emissions *Soil Use and Management* 13 230–244
- Pellegrineschi, A 1997**
In vitro plant regeneration via organogenesis of cowpea [*Vigna unguiculata* (L.) Walp.] *Plant Cell Reports* 17 89–95
- Pellegrineschi, A , C A Fatokun, G Thottappilly, and A A Adepoju 1997**
Cowpea embryo rescue 1 Influence of culture media composition on plant recovery from isolated immature embryos *Plant Cell Reports* 17 133–138
- Peters, M , S A Tarawali, and J Alkamper 1997**
Dry season growth performance of tropical pasture legumes in subhumid Nigeria *Tropical Grasslands* 31 201–231
- Risasi, E L , G Tian, E E Opuwaribo, and B T Kang 1997**
Effect of root size on root nutrient concentrations of four woody trees and maize *Forest Farm and Community Tree Research Report* 2 13–17
- Sanginga, N , K Dashiell, J A Okogun, and G Thottappilly 1997**
Nitrogen fixation and N contribution by promiscuous nodulating soybeans in the southern Guinea savanna of Nigeria *Plant and Soil* 195 257–266
- Scholz, D , C Borgemeister, W G Meikle, R H Markham, and H M Poehling 1997**
Infestation of maize by *Prostephanus truncatus* initiated by male produced pheromone *Entomologia Experimentalis et Applicata* 83 53–61
- Schulthess, F , N A Bosque Perez, A Chabi Olaye, S Gounou, R Ndemah, and G Goergen 1997**
Exchanging natural enemies species of lepidopterous cereal stemborers between African regions *Insect Science and its Application* 17 97–108
- Schulthess, F , P Neuenschwander, and S Gounou 1997**
Multi trophic interactions in cassava *Manihot esculenta* cropping systems in the subhumid tropics of West Africa *Agriculture Ecosystems and Environment* 66 211–222
- Setamou, M , K F Cardwell, F Schulthess, and K Hell 1997**
Aspergillus flavus infection and aflatoxin contamination of preharvest maize in the Republic of Benin *Plant Disease* 81 1323–1327
- Shoyinka, S A , G Thottappilly, G G Adebayo, and F O Anno Nyako 1997**
Survey on cowpea virus incidence and distribution in Nigeria *International Journal of Pest Management* 43 127–132

- Smith, J W , A Naazie, A Larbi, K Agyemang, and S A Tarawali 1997**
Integrated crop–livestock systems in sub Saharan Africa an option or imperative? Outlook on Agriculture 26 237–246
- Staubli Dreyer B , J Baumgartner, P Neuenschwander, and S Dorn 1997**
The functional responses of two *Hyperaspis notata* strains to their prey the cassava mealybug *Phenacoccus manihoti* Bulletin de la Societe Entomologique Suisse 70 21–28
- Staubli Dreyer, B , P Neuenschwander, J Baumgartner, and S Dorn 1997**
Trophic influences on survival development and reproduction of *Hyperaspis notata* (Col Coccinellidae) Journal of Applied Entomology 121 249–256
- Stonehouse, J M , C Gbongboui , A de Groot, C J Lomer, S Ly, I Marga, and Tijani 1997**
Grasshopper control in the Sahel Farmer perceptions and participation Crop Protection 16 733–741
- Thomas, M B , J Langewald, and S N Wood 1996**
Evaluating the effects of a biopesticide on populations of the variegated grasshopper *Zonocerus variegatus* Journal of Applied Ecology 33 1509–1516
- Thomas, M B , S N Wood, J Langewald, and C J Lomer 1997**
Persistence of biopesticides and consequences for biological control of grasshoppers and locusts Pesticide Science 49 47–55
- Thomas, M B , S Blanford, and C J Lomer 1997**
Reduction of feeding by the variegated grasshopper *Zonocerus variegatus* following infection by the fungal pathogen *Metarhizium flavoviride* Biocontrol Science and Technology 7 327–334
- Thottappilly, G , and H W Rossel 1997**
Identification and characterization of viruses infecting bambara groundnut (*Vigna subterranea*) in Nigeria International Journal of Pest Management 43 177–185
- Tian, G , and L Brussaard 1997**
Mulching effect of plant residues of chemically contrasting compositions on soil organic matter content and cation exchange capacity Communications in Soil Science and Plant Analysis 28 1603–1611
- Tian, G , B T Kang, and L Brussaard 1997**
Effect of mulch quality on earthworm activity and nutrient supply in the humid tropics Soil Biology and Biochemistry 29 369–373
- Tonukari, N J , G Thottappilly, N Q Ng, and H D Mignouna 1997**
Genetic polymorphism of cassava (*Manihot esculenta* Crantz) germplasm within the Republic of Benin detected with RAPD markers African Journal of Crop Science 5 219–228
- Umoren, U E , O O Tewe, M Bokanga, and L E N Jackai 1997**
Protein quality of raw and autoclaved cowpeas Comparison between some insect resistant and susceptible varieties Plant Foods for Human Nutrition 50 301–315
- Vanlauwe, B , N Sanginga, and R Merckx 1997**
Decomposition of four *Leucaena* and *Senna* prunings in alley cropping systems under subhumid tropical conditions The process and its modifiers Soil Biology and Biochemistry 29 131–137
- Vuyisteke, D , R Ortiz, R S B Ferris, and J H Crouch 1997**
Plantain improvement Plant Breeding Reviews 14 267–320

Book Chapters/Conference Papers

- Abera, A C S Gold, and S Kyamanywa 1997**
Banana weevil oviposition and damage in Uganda Pages 1199–1205 in Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda
- Adejuyigbe, C O , G Tian, G O Adeoye, and B T Kang 1997**
Impact of land use practices on soil microarthropod populations Pages 499–505 in Proceedings 3rd African Soil Science Society Conference 19–26 Aug 1995 Ibadan Nigeria OAU/STRC Lagos Nigeria
- Asiedu, R , N M Wanyera, S Y C Ng, and N Q Ng 1997**
Yams Pages 57–66 in Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK



- Badejo, M A , G Tian, and L Brussaard 1996**
Response of cryptostigmatid mites (Oribatida) to plant residues of different chemical compositions in a maize cropping system Pages 595–597 *in* Proceedings Acarology IX Vol 1 edited by R Mitchell D J Horn G R Needham and W C Welbourn The Ohio Biological Survey Columbus OH USA
- Bigirwa, G , E Adipala, P Esele, and K F Cardwell 1997**
Disease progress of *Peronosclerospora sorghi* on some Ugandan maize genotypes Pages 229–232 *in* Maize productivity gains through research and technology dissemination edited by J K Ransom A F E Palmer B T Zambezi Z O Mduruma S R Waddington K V Pixley and D C Jewell Proceedings 5th Eastern and Southern Africa Regional Maize Conference 3–7 Jun 1996 Arusha Tanzania CIMMYT Addis Ababa Ethiopia
- Blade, S F , S V R Shetty, T Terao, and B B Singh 1997**
Recent developments in cowpea cropping systems research Pages 114–128 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Blomme, G , and R Ortiz 1997**
Preliminary evaluation of variability in *Musa* root system development Pages 51–52 *in* Biology of root formation and development edited by A Altman and Y Waisel Plenum Publishing Corporation New York USA
- Bonierbale, M , C Guevara, A G O Dixon, N Q Ng, R Asiedu, and S Y C Ng 1997**
Cassava Pages 1–20 *in* Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK
- Bosque Perez, N A , J G Kling, and S I Odubiyi 1997**
Recent advances in the development of sources of resistance to pink stalk borer and African sugarcane borer Pages 234–240 *in* Insect resistant maize Recent advances and utilization edited by J A Mihm Proceedings 2nd International Symposium on Insect Resistant Maize 27 Nov to 2 Dec 1994 El Batan Mexico International Wheat and Maize Improvement Center (CIMMYT) Mexico DF Mexico
- Bottenberg, H , M Tamo, D Arodokoun, L E N Jackai, B B Singh, and O Youm 1997**
Population dynamics and migration of cowpea pests in northern Nigeria Implications for integrated pest management Pages 271–284 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Chikoye, D , F Ekeleme, and I O Akobundu 1997**
Weed composition and population dynamics in intensified smallholder farms in West Africa Pages 161–166 *in* The 1997 Brighton Crop Protection Conference—Weeds Proceedings of an International Conference 17–20 Nov 1997 Brighton UK British Crop Protection Council Farnham Surrey UK
- Dashiell, K , and C Fatokun 1997**
Soyabean Pages 181–190 *in* Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK
- De Groot, H 1997**
Women s income versus family income as a determinant for food security an example from southern Mali Pages 12–22 *in* Issues in agricultural competitiveness Markets and policies Proceedings 22nd Conference of Agricultural Economists Harare Zimbabwe 22–29 Aug 1994 edited by R Rose C Tanner and M Bellamy International Association of Agricultural Economists Dartmouth Aldershot UK

- De Groote, H , and N Coulibaly 1997**
Acces aux ressources productives selon genre et generation Pages 253–264 *in* Institutions and technologies for rural development in West Africa Proceedings International Symposium 16–22 Feb 1996 Cotonou Benin edited by T Bierschenk Y Le Meur and M von Oppen Margraf Verlag Weikersheim Germany
- Emechebe, A M , and D A Florini 1997**
Shoot and pod diseases of cowpea induced by fungi and bacteria Pages 176–192 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Emechebe, A M , and B B Singh 1997**
Technology options and research challenges to increase cowpea production under *Striga* and drought stress in semi arid Africa Pages 207–217 *in* Technology options for sustainable agricultural production in sub Saharan Africa edited by T Bezuneh A M Emechebe J Sedgo and M Ouedraogo OAU/STRC—SAFGRAD Ouagadougou Burkina Faso
- Fatokun, C A , N D Young, and G O Myers 1997**
Molecular markers and genome mapping in cowpea Pages 352–360 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Fatokun, C A , P Perrino, and N Q Ng 1997**
Wide crossing in African *Vigna* species Pages 50–57 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Ferris, R S B , J B A Whyte, B Khizzah, and J Legg 1997**
Opportunities for commercializing cassava in East and Southern Africa Pages 1427–1433 *in* Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda
- Fery, R L , and B B Singh 1997**
Cowpea genetics A review of the recent literature Pages 13–29 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Florini, D A 1997**
Nematodes and other soilborne pathogens of cowpea Pages 193–206 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Gold, C S , S H Okech, E B Karamura, and R Ssendege 1997**
Banana weevil population densities and related damage in Ntungamo and Mbarara districts Uganda Pages 1207–1219 *in* Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda
- Gold, C S , A Kiggundu, and A M Abera 1997**
Farmer banana cultivar selection criteria in Uganda Pages 139–158 *in* Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda
- Goli, A E , F Begemann, and N Q Ng 1997**
Characterization and evaluation of IITA s bambara groundnut collection Pages 101–108 *in* Bambara groundnut *Vigna subterranea* (L) Verdc edited by J Heller F Begemann and J Mushonga Promoting the conservation and use of underutilized and neglected crops 9 Proceedings of the Workshop on Conservation and Improvement of Bambara Groundnut (*Vigna subterranea* (L) Verdc) 14–16 Nov 1995 Harare Zimbabwe Institute of Plant Genetics and Crop Plant Research Gatersleben Germany/Department of Research & Specialist Services Harare Zimbabwe/International Plant Genetic Resources Institute Rome Italy



- Hampton, R O , G Thottappilly, and H W Rossel 1997**
 Viral diseases of cowpea and their control by resistance conferring genes Pages 159–175 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Hauser, S , B Vanlauwe, D O Asawalam, and L Norgrove 1997**
 Role of earthworms in traditional and improved low input agricultural systems in West Africa Pages 113–136 *in* Soil ecology in sustainable agricultural systems edited by L Brussaard and R Ferrera Cerrato CRC Press/Lewis Publishers Boca Raton/New York USA
- Horry, J P , R Ortiz, E Arnaud, J H Crouch, R S B Ferris, D R Jones, N Mateo, C Picq, and D Vuylsteke 1997**
 Banana and plantain Pages 67–81 *in* Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK
- Inaizumi, H , and O Brodie Mends 1997**
 Determinants of adoption of improved cassava processing technologies in West African countries Pages 376–381 *in* Journal of Rural Economics (Special Issue) Proceedings Annual Conference of the Agricultural Economics Society of Japan 1997 Tokyo Japan
- Inaizumi, H , A Enete, O Brodie Mends, and E Oyetunji 1997**
 Determinants of mechanized cassava processing technology adoption in West Africa Pages 51–62 *in* Ecosystem and sustainable development edited by J L Uso C A Brebbia and H Power Computational Mechanics Publications Southampton UK
- Jackai, L E N , and C B Adalla 1997**
 Pest management practices in cowpea A review Pages 240–258 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Jackson, M T , G C Loresto, S Appa Rao, M Jones, E P Guimaraes, and N Q Ng 1997**
 Rice Pages 273–291 *in* Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK
- Jeon, Y W , and L Halos Kim 1997**
 Improving postharvest technology development in Africa Pages 133–141 *in* Women agricultural intensification and household food security edited by S A Breth Proceedings Workshop 25–28 Jun 1996 University of Cape Coast Ghana Sasakawa Africa Association Mexico City Mexico
- Kamara, A Y , S C Jutzi, I O Akobundu, and D Chikoye 1997**
 The effect of mulch from three multipurpose trees on weed composition and biomass in maize Pages 653–654 *in* The 1997 Brighton Crop Protection Conference—Weeds Proceedings of an International Conference 17–20 Nov 1997 Brighton UK British Crop Protection Council Farnham Surrey UK
- Khizzah, B W , J B A Whyte, J Legg, S Ferris, and H Ojulong 1997**
 Cassava germplasm enhancement for food security and industrial utilization in East and Southern Africa Pages 1289–1295 *in* Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda
- Lagoke, S T O , J Y Shebayan, I Magani, P Olorunju, O O Olufajo, K A Elemo, I Uvah, A A Adeoti, P S Chindo, I Kureh, S Jatau, A M Emechebe, W B Ndahi, S K Kim, G Weber, B B Singh, C Odion, and A Avan 1997**
Striga problem and development of appropriate control strategies in various crops in Nigeria Pages 89–100 *in* Integrated management of *Striga* for the African farmers edited by S T O Lagoke I E van der Straten and S S Mboob Proceedings 3rd General Workshop of the Pan African *Striga* Control Network (PASCON) 18–20 Oct 1993 Harare Zimbabwe FAO Accra Ghana

- Lattanzio, V , A Cardinali, V Linsalata, P Perrino, and N Q Ng 1997**
 Flavonoid HPLC fingerprints of wild *Vigna* species Pages 66–74 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Manyong, V M , and B Oyewole 1997**
 Geographic definition of risk area for maize production in West Africa A socio economic perspective Pages 130–134 *in* Maize productivity gains through research and technology dissemination edited by J K Ransom A F E Palmer B T Zambezi Z O Mduruma S R Waddington K V Pixley and D C Jewell Proceedings 5th Eastern and Southern Africa Regional Maize Conference 3–7 Jun 1996 Arusha Tanzania CIMMYT Addis Ababa Ethiopia
- Minde I J , J M Teri, V W Saka, K Rockman, and I R M Benesi 1997**
 Accelerated multiplication and distribution of cassava and sweetpotato planting material in Malawi Pages 162–167 *in* Alternative strategies for smallholder seed supply Proceedings of an International Conference on Options for Strengthening National and Regional Seed Systems in Africa and West Asia 10–14 Mar 1997 Harare Zimbabwe edited by D D Rohrbach Z Bishaw and A J G van Gastel International Crops Research Institute for the Semi Arid Tropics Patancheru 502 324 Andhra Pradesh India
- Modder, W W D 1997**
 Attraction and diurnal behaviour of the African pest grasshopper *Zonocerus variegatus* (L) at oviposition sites Pages 283–292 *in* New strategies in locust control edited by S Krall R Peveling and D Ba Diallo Birkhauser Verlag Basel Switzerland
- Monti, L M , L L Murdock, and G Thottappilly 1997**
 Opportunities for biotechnology in cowpea Pages 341–351 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Mortimore, M J , B B Singh, F Harris, and S F Blade 1997**
 Cowpea in traditional cropping systems Pages 99–113 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Msikita, W , R M Skirvin, and S Y Chen 1997**
 Micropropagation of *Brassica oleracea* (Cole crops) Pages 30–47 *in* Biotechnology in agriculture and forestry Vol 39 High tech and micropropagation V edited by Y P S Bajaj Springer Verlag Berlin Germany
- Muhr, L , S A Tarawali, M Peters, R Schultze Kraft, and D Berner 1997**
 Multiple uses of tropical forage legumes for sustainable farming in the moist savannas of Africa Pages 19 55–19 56 *in* Proceedings XVIII International Grassland Congress 8–19 Jun 1997 Winnipeg and Saskatoon Canada International Grassland Congress Calgary Alberta Canada
- Ng, N Q , and B B Singh 1997**
 Cowpea Pages 82–99 *in* Biodiversity in trust Conservation and use of plant genetic resources in CGIAR Centres edited by D Fuccillo L Sears and P Stapleton Cambridge University Press Cambridge UK
- Ng S Y C , and N Q Ng 1997**
 Germplasm conservation in food yams (*Dioscorea* spp) Constraints application and future prospects Pages 257–286 *in* Conservation of plant genetic resources *in vitro* Vol 1 General aspects edited by M K Razdan and E C Cocking Science Publishers Enfield NH USA
- Oikeh, S O , J G Kling, W J Horst, and V O Chude 1997**
 Yield and N use efficiency of five tropical maize genotypes under different N levels in the moist savanna of Nigeria Pages 163–167 *in* Maize productivity gains through research and technology dissemination edited by J K Ransom A F E Palmer B T Zambezi Z O Mduruma S R Waddington K V Pixley and D C Jewell Proceedings 5th Eastern and Southern Africa Regional Maize Conference 3–7 Jun 1996 Arusha Tanzania CIMMYT Addis Ababa Ethiopia



Okech, S H O , A Karugaba, C S Gold, A Nyakuni, H Ssali, and E B Karamura 1997

Influence of soil conservation bunds compost manure coffee and bean intercropping on weevil incidence banana vigour and bunch weight in Bagambaa sub county a hilly environment in Mbarara southwestern Uganda Pages 1221–1228 in Vol 3 Proceedings African Crop Science Conference 13–17 Jan 1997 Pretoria South Africa edited by E Adipala J S Tenywa and M W Ogenga Latigo African Crop Science Society Kampala Uganda

Padulosi, S , and N Q Ng 1997

Origin taxonomy and morphology of *Vigna unguiculata* (L) Walp Pages 1–12 in Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria

Peters, M , S A Tarawali, R Schultze Kraft, and A Musa 1997

Legume–legume complementarity for sustainable pasture development in the tropics Pages 22 123–22 124 in Proceedings XVIII International Grassland Congress 8–19 Jun 1997 Winnipeg and Saskatoon Canada International Grassland Congress Calgary Alberta Canada

Singh, B B , and S F Blade 1997

Potential of dry season cowpea in West Africa Pages 227–233 in Technology options for sustainable agricultural production in sub Saharan Africa edited by T Bezuneh A M Emechebe J Sedgo and M Ouedraogo OAU/STRC—SAFGRAD Ouagadougou Burkina Faso

Singh, B B , and A M Emechebe 1997

Advances in research on cowpea *Striga* and *Alectra* Pages 215–224 in Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria

Singh, B B , and S A Tarawali 1997

Cowpea An integral component of sustainable mixed crop/livestock farming systems in West Africa and strategies to improve its productivity Pages 79–100 in Crop residues in sustainable mixed crop–livestock farming systems edited by C Renard CAB International Wallingford Oxon UK in association with the International Crops Research Institute for the Semi Arid Tropics (ICRISAT) and the International Livestock Research Institute (ILRI)

Singh, B B , I D K Atokple, and A M Emechebe 1997

Controlling *Striga* and *Alectra* on cowpea by host plant resistance Pages 32–35 in Integrated management of *Striga* for the African farmers edited by S T O Lagoke I E van der Straten and S S Mboob Proceedings 3rd General Workshop of the Pan African *Striga* Control Network (PASCON) 18–20 Oct 1993 Harare Zimbabwe FAO Accra Ghana

Singh, B B , O L Chambliss, and B Sharma 1997

Recent advances in cowpea breeding Pages 30–49 in Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria

Singh, B N , A T Maji, N Q Ng, C Paul, C Williams, and M N Ukwungwu 1997

Utilization of *Oryza glaberrima* genetic resources for lowland rice improvement Pages 177–187 in Interspecific hybridization Progress and prospects edited by M P Jones M Dingkuhn D E Johnson and S O Fagade WARDA/ADRAO Bouake Cote d'Ivoire

- Smith, J, G Weber, V M Manyong, and M A B Fakorede 1997**
Fostering sustainable increases in maize productivity in Nigeria Pages 107–124 *in* Africa's emerging maize revolution edited by D Byerlee and C K Eicher Lynne Rienner Publishers Boulder CO USA
- Sonnante, G, A R Piergiovanni, N Q Ng, and P Perrino 1997**
Isozyme markers and taxonomic relationships among *Vigna* species Pages 58–65 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Tamo, M, H Bottenberg, D Arodokoun, and R Adeoti 1997**
The feasibility of classical biological control of two major cowpea insect pests Pages 259–270 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Tarawali, S A, B B Singh, M Peters, and S F Blade 1997**
Cowpea haulms as fodder Pages 313–325 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Tarawali, S A, B B Singh, S Fernandez Rivera, M Peters, J W Smith, R Schultze Kraft, and H A Ajeigbe 1997**
Optimizing the contribution of cowpea to food and fodder production in crop–livestock systems in West Africa Pages 19 53–19 54 *in* Proceedings XVIII International Grassland Congress 8–19 Jun 1997 Winnipeg and Saskatoon Canada International Grassland Congress Calgary Alberta Canada
- Terao, T, I Watanabe, R Matsunaga, S Hakoyama, and B B Singh 1997**
Agro physiological constraints in intercropped cowpea An analysis Pages 129–140 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria
- Tian, G, M A Badejo, and C O Adejuyigbe 1997**
Soil faunal activity as an important component of soil fertility through influencing plant decomposition Pages 579–590 *in* Proceedings 3rd African Soil Science Society Conference 19–26 Aug 1995 Ibadan Nigeria OAU/STRC Lagos Nigeria
- Tian, G, L Brussaard, B T Kang, and M J Swift 1997**
Soil fauna mediated decomposition of plant residues under constrained environmental and residue quality conditions Pages 125–134 *in* Driven by nature Plant litter quality and decomposition edited by G Cadisch and K E Giller CAB International Wallingford Oxon UK
- Vanlauwe, B, J Diels, N Sanginga, and R Merckx 1997**
Residue quality and decomposition An unsteady relationship? Pages 157–166 *in* Driven by nature Plant litter quality and decomposition edited by G Cadisch and K E Giller CAB International Wallingford Oxon UK
- Watanabe I, S Hakoyama, T Terao, and B B Singh 1997**
Evaluation methods for drought tolerance of cowpea Pages 141–146 *in* Advances in cowpea research edited by B B Singh D R Mohan Raj K E Dashiell and L E N Jackai Copublication of the International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS) IITA Ibadan Nigeria

Training Materials

- Berner, D K, M D Winslow, A E Awad, K F Cardwell, D R Mohan Raj, and S K Kim (editors) 1997**
Striga research methods A manual (2nd edition) International Institute of Tropical Agriculture Ibadan Nigeria 82 pp



- Buckland, L , and J Haleegoah 1996**
Gender analysis in agricultural production IITA Research Guide 58 Training Program, IITA, Ibadan Nigeria 24 pp
- Ekanayake, I J , D S O Osiru, and M C M Porto 1997**
Agronomy of cassava IITA Research Guide 60 Training Program IITA Ibadan Nigeria 30 pp
- Ekanayake, I J , D S O Osiru, and M C M Porto 1997**
Morphology of cassava IITA Research Guide 61 Training Program IITA Ibadan Nigeria 30 pp
- Ferris, R S B 1997**
Improving storage life of plantain and banana IITA Research Guide 62 Training Program IITA Ibadan Nigeria 22 pp
- Swennen, R , and R Ortiz 1997**
Morphology and growth of plantain and banana IITA Research Guide 66 Training Program IITA Ibadan Nigeria 32 pp
- Thottappilly, G , and H W Rossel 1996**
Virus diseases of cowpea in Africa IITA Research Guide 53 Training Program IITA Ibadan Nigeria 28 pp

Abstracts

- Afouda, L , and K Wydra 1997**
Pathological characterization of root and stem rot pathogens of cassava and evaluation of antagonists for their biological control DPG Working Group Plant Protection in the Tropics and Subtropics Berlin July 1997 Phytomedizin 27(4) 43–44
- Akano, A O , S Y C Ng, A G O Dixon, and G Thottappilly 1997**
Application of embryo culture in establishing African cassava mosaic disease resistant gene mapping population Page 61 in Book of abstracts 11th Symposium of the International Society for Tropical Root Crops (ISTRC) 19–24 Oct 1997 Trinidad and Tobago
- Akano, A O , R Asiedu, and S Y C Ng 1997**
Effect of African cassava mosaic disease on the yield of cassava derived from cuttings from tissue culture plants in two agroecological zones of Nigeria African Journal of Root and Tuber Crops 2 48
- Anegebeh, P O , T Simons, B Duguma, I Dawson, H Jaenicke, and P D Austin 1997**
Household prospects for fruit production from *Irvingia gabonensis* (Aubry Lecomte ex O Rorke) Baill Pages 24–25 in Abstracts 15th Annual Conference Horticultural Society of Nigeria (HORTSON) 8–11 Apr 1997 National Horticultural Research Institute (NIHORT) Ibadan Nigeria
- Dahal, G , J d'A Hughes, G Thottappilly, and B E L Lockhart 1997**
Effect of temperature on symptom expression and detection of banana streak badnavirus in plantain and bananas Phytopathology 87 S 22
- Dahal, G, J d A Hughes, F Gauhl, C Pasberg Gauhl, G Thottappilly, A Tenkouano, D Vuylsteke, and B E L Lockhart 1997**
Evaluation of micropropagated plantain and banana hybrids for banana streak badnavirus incidence in Nigeria Phytopathology 87 S 22
- Dahal, G , C Pasberg Gauhl, F Gauhl, G Thottappilly, and J d'A Hughes 1997** Intraplant variation in symptom expression and distribution of banana streak badnavirus in naturally infected plantain and banana (*Musa spp*) Phytopathology 87 S 22
- Dochez, C , P Spejler, D De Waele, and R Ortiz 1997**
Host plant response to nematodes in *Musa* landraces and hybrids Pages 21–22 in Abstracts 13th Symposium of the Nematological Society of Southern Africa 9–13 Mar 1997 San Lamber Kwazulu Natal Republic of South Africa

- Fessehaie, A , K Wydra, and K Rudolph 1997**
 Detection of *Xanthomonas campestris* pv *manihotis* specific proteins by electrophoresis and immunoblotting for development of monospecific polyclonal antibodies DPG Working Group Plant Protection in the Tropics and Subtropics Berlin July 1997 Phytomedizin 27(4) 45
- Fessehaie, A , K Wydra, and K Rudolph 1997**
 Immuno elektrophoretische Untersuchungen mit Proteinen von *Xanthomonas campestris* pv *manihotis* für die Herstellung hochspezifischer Antikörper Phytomedizin 27(1) 54
- Hughes, J d'A , L N Dongo, and S Y C Ng 1997**
 Diagnosis of yam viruses Page 55 in Book of abstracts 11th Symposium of the International Society for Tropical Root Crops (ISTRC) 19–24 Oct 1997 Trinidad and Tobago
- Hughes, J d A , L N Dongo, and G I Atiri 1997**
 Viruses infecting cultivated yams (*Dioscorea alata* and *D rotundata*) in Nigeria Phytopathology 87 S 45
- Jackai, L E N , G Thottappilly, and T I Ofuya 1997**
 Variable demography in the Nigeria population of *Callosobruchus maculatus* Fab on cowpea *Vigna unguiculata* (L) Walp and another food legume *Sphenostylis sternocarpa* Harms Pages 97–98 in Joint Congress of the Entomological Society of Southern Africa (11th Congress) and the African Association of Insect Scientists (12th Congress) 30 Jun–4 Jul 1997 Stellenbosch South Africa
- Kling, J G , D K Berner, and O A Ibikunle 1997**
 Sources of resistance to *Striga* in tropical maize and teosinte Page 71 in Agronomy Abstracts American Society of Agronomy Madison WI USA
- Ng, S Y C , and S H Mantell 1997**
 Influence of carbon source on in vitro tuberization and growth of white yam (*Dioscorea rotundata* Poir) Page 26 in Book of abstracts 11th Symposium of the International Society for Tropical Root Crops (ISTRC) 19–24 Oct 1997 Trinidad and Tobago
- Ortiz, R , and D Vuylsteke 1997**
 Improved polyploid *Musa* germplasm developed by IITA through ploidy manipulations Page 99 in Abstracts First All Africa Crop Science Congress 13–17 Jan 1997 University of Pretoria Pretoria Republic of South Africa
- Singh, B B 1997**
 Performance of cowpea breeding lines developed from pedigree selection under sole and intercrop in different cropping systems Page 67 in Agronomy Abstracts American Society of Agronomy Madison WI USA
- Singh, B B , R Tabo, S C Gupta, and H Ajeigbe 1997**
 Increasing productivity of sorghum–cowpea and millet–cowpea intercrops through improved varieties Page 41 in Agronomy Abstracts American Society of Agronomy Madison WI USA
- Thottappilly, G , G Dahal, G Harper, R Hull, and B E L Lockhart 1997**
 Banana streak badnavirus Development of diagnostics by ELISA and PCR Phytopathology 87 S 97
- Wydra, K , A Fanou, R Sikirou, I Adamou, V Zinsou, and A Avocan 1997**
 Expression of field resistance and tolerance in cassava and cowpea to cassava bacterial blight and cowpea bacterial blight respectively DPG Working Group Plant Protection in the Tropics and Subtropics Berlin July 1997 Phytomedizin 27(4) 45–46

Other Reports/ Miscellaneous

- Crouch, J H , H K Crouch, R Ortiz, and R L Jarret 1997**
 Microsatellites for molecular breeding of *Musa* InfoMusa 6(1) 5–6
- Czerwenka Wenkstetten, I M , D K Berner, and A Schilder 1997**
 First report and pathogenicity of *Myrothecium roridum* *Curvularia eragrostidis* and *C lunata* on seeds of *Striga hermonthica* in Nigeria Plant Disease 81 832
- Dixon, A G O 1997**
 Cassava mosaic disease in sub Saharan Africa potential of African landraces as sources of resistance International Society for Tropical Root Crops (ISTRC) Newsletter 24 5–6



Dixon, A G O 1997

Promoting broader diversity in crops African farmers winning the fight against cassava disease *Geneflow* 1997 (IPGRI) 19

Gauhi, F , C Pasberg Gauhi, and J d'A Hughes 1997

First report of banana streak badnavirus in plantain landraces in southern Cameroon Central Africa *Plant Disease* 81 1335

IPGRI/IITA 1997

Descriptors for yam (*Dioscorea* spp) International Institute of Tropical Agriculture Ibadan Nigeria/International Plant Genetic Resources Institute Rome Italy 64 pp

Msikita, W , J S Yaninek, M Ahounou, H Baimey, and R Fagbemissi 1997

First report of *Curvularia lunata* associated with stem disease of cassava *Plant Disease* 81 112

Msikita, W , J S Yaninek, M Ahounou, H Baimey, and C O Fagbemissi 1997

First report of *Nattrassia mangiferae* root and stem rot of cassava in West Africa *Plant Disease* 81 1332

Muimba Kankolongo, A , A Chalwe, P Sisupo, and N C Kanga 1997

Distribution prevalence and outlook for control of cassava mosaic disease in Zambia *Roots Newsletter* 4(1) 2-7

Ogbe, F O , J Legg, M D Raya, A Muimba Kankolongo, M P Theu, N A Phiri, G Kaitisha, and A Chalwe 1997


Diagnostic survey of African cassava mosaic viruses in Tanzania Malawi and Zambia *Roots Newsletter* 4(2) 12-15

Ortiz, R 1997

A delivery system of improved banana and plantain propagules *InfoMusa* 6(2) 14-15

Schill, P , K Afreh Nuamah, C Gold, F Ulzen Apiah, E Paa Kwesi, S A Peprah, and J K Twumasi 1997

Farmers perception of constraints in plantain production in Ghana *Plant Health Management Division Monograph* No 5 IITA Ibadan Nigeria 54 pp



Pierre Dubreuil, Chair*

c/o CIRAD Paris France

Enrico Porceddu, Chair

Professor Dipartimento di Agrobiologia e Agrochimica
Universita Degli Studi Della Tuscia Viterbo Italy

Adamu Aliyu**

Director of Agricultural Sciences and Head of NARP
Federal Ministry of Agriculture and Natural Resources
Abuja Nigeria

Lukas Brader**

Director General IITA
Ibadan Nigeria

Lmamoulave Cisse

Head Agronomy Division
Institut mondial du phosphate
Casablanca Morocco

Martin Clucas

Martin Clucas & Associates
West Mersea Essex UK

Volker Hoffmann

Professor Universitat Hohenheim
Stuttgart Germany

Jørgen Jakobsen

Head Department of Plant Pathology and Pest Management
Danish Institute of Plant and Soil Science
Lyngby Denmark

Anastasios Leventis

Leventis Overseas
London UK

Constance McCorkle

Senior Research Scientist and Consultant
Washington D C USA

Joseph Mukubi

Director General
National Agricultural Research Organization
Entebbe Uganda

Mary Okelo

Kaboke House
Nairobi Kenya

Kyoko Saito

Tokyo Metropolitan Food and Technology Research Center
Tokyo Japan

John Stewart*

Dean of Agriculture University of Saskatchewan
Saskatoon Canada

Michel Sedogo

Directeur General Centre national de la recherche
scientifique et technologique
Ouagadougou Burkina Faso

Eric Tollens

Professor Faculty of Agricultural Sciences
Katholieke Universiteit Leuven
Hervelee Belgium

Marc van Montagu*

Director Laboratorium voor Genetica
Gent Belgium

Permanent Secretary**

Federal Ministry of Agriculture and Natural Resources
Abuja Nigeria

Vice Chancellor**

By rotation among Nigerian agricultural universities

*Left during the year

**Ex officio

Professional Staff

The listings presented here indicate the staff member's name, designation, and work location (in bold type). Staff from other CGIAR centers and institutions working in collaboration with IITA are also listed.

Director General

L Brader PhD

Office of the Director General

R H Booth PhD deputy director general **Ibadan**
S A Adetunji PhD special assistant to director general **Ibadan***
J Cramer BA executive assistant to the director general **Ibadan**
C F McDonald MSc assistant to the deputy director general **Ibadan**

F O Oke records/archival officer **Ibadan**

Crop Improvement Division

F M Quin PhD agronomist director **Ibadan**

J N Agba BSc station administrative manager **Onne**
M B Alti, MBA station administrative manager **Kano**

Grain Legume Improvement Program

K E Dashiell PhD breeder/geneticist program leader **Ibadan**
C A Fatokun PhD breeder/geneticist **Ibadan**
B B Singh, PhD breeder/geneticist and officer in charge **Kano**

Visiting scientist

O Nakayama PhD food technologist **Ibadan**

Maize Improvement Program

F M Quin, PhD agronomist program leader **Ibadan**
J G Kling PhD breeder/geneticist **Ibadan**
A Menkir PhD breeder/geneticist **Ibadan**

Visiting scientists

O S Ajala PhD breeder (WVI) **Ibadan***
B Maziya Dixon PhD food technologist **Ibadan***

Plantain and Banana Improvement Program

D Vuylsteke Ir agronomist/breeder program leader & ESARC team leader **Namulonge**
J H Crouch PhD germplasm enhancer/biotechnologist **Onne***
M Pillay PhD cytogeneticist **Onne**
A Tenkouano PhD breeder/geneticist **Onne**

Postdoctoral fellow

M Tshionza PhD socioeconomist **Onne**

Associate experts

G Blomme Ir agronomist **Onne**
C Dochez **Onne***

Root and Tuber Improvement Program

R Asiedu PhD breeder/geneticist program leader **Ibadan**
M Bokanga PhD biochemist and food technologist **Ibadan**
A G O Dixon PhD breeder/geneticist **Ibadan**
I Ekanayake PhD physiologist **Ibadan***
S Y C Ng MSc tissue culture specialist **Ibadan**
P Venier PhD yam specialist (CIRAD) **Cotonou**

Visiting scientist

O Girardin PhD yam scientist **Ibadan**

Postdoctoral fellow

N Wanyera PhD breeder/agronomist **Ibadan***

Biotechnology Research Unit

G Thottappilly PhD virologist unit head **Ibadan**
D H Mignouna PhD molecular biologist **Ibadan**

Postdoctoral fellows

J S Machuka PhD biotechnologist **Ibadan**
A Pellegrineschi PhD, biotechnologist **Ibadan**

Visiting scientist

O G Omitogun PhD molecular biologist **Ibadan***

Genetic Resources Unit

N Q Ng PhD germplasm scientist unit head **Ibadan**

Research Farms Unit

P Austin BSc unit head and officer in charge **Onne**
G O Olayode MSc farm management officer **Ibadan***

Plant Health Management Division

P Neuenschwander PhD entomologist director **Cotonou**

Administration and Support Services

J N Quaye MA chief administrator **Cotonou***
J B Akinwumi MSc engineer **Cotonou**
J B Adjovi chief accountant **Cotonou**
S Nyampong executive assistant to the director **Cotonou**
A Ratie maintenance services officer **Cotonou**

Biological Control Program

C J Lomer PhD insect pathologist program leader **Cotonou**
C Borgemeister PhD entomologist **Cotonou***
A Cherry PhD entomologist (NRI) **Cotonou**

Left during the year

H De Groot PhD socioeconomist **Cotonou**
G Goergen PhD entomologist **Cotonou**
B D James PhD coordinator ESCaPP project **Cotonou**
J Langewald PhD entomopathologist **Cotonou**
P Le Gall PhD entomologist (ORSTOM) **Cotonou**
R H Markham PhD coordinator systemwide project on IPM **Ibadan**
B Megevand MSc mite rearing specialist **Cotonou***
W Meikle, PhD entomologist **Cotonou**
K Wydra PhD plant pathologist **Cotonou**
J S Yaninek PhD acarologist, **Cotonou**

Postdoctoral fellows

S Dara PhD entomopathologist **Cotonou**
W Msikita PhD plant pathologist, **Cotonou**

Visiting scientist

M Toko, PhD entomologist **Cotonou**

Associate experts

D Muller MSc PG Dip , **Cotonou**
C Nansen MSc **Cotonou**

Habitat Management Program

M Tamo PhD insect ecologist program leader **Cotonou**
K F Cardwell PhD plant pathologist **Cotonou/Ibadan**
K Green PhD, coordinator IITA/GTZ West African plantain project, **Accra**
W N O Hammond PhD coordinator PEDUNE **Cotonou**
P Schill PhD entomologist **Accra***
F Schulthess PhD ecologist **Cotonou**

Visiting scientist

V Adenle PhD plant pathologist **Ibadan**

Host Plant Resistance Program

D Berner PhD parasitic weeds biologist program leader **Ibadan**
G Dahal PhD virologist **Ibadan***
F Gauhl PhD plant pathologist **Yaounde***
J d A Hughes PhD virologist, **Ibadan**
L E N Jackai PhD entomologist **Ibadan**
C Pasberg Gauhl PhD plant pathologist **Yaounde***

Postdoctoral fellows

S K Asante PhD entomologist **Kano**
O Coulibaly PhD agricultural economist **Yaounde**
K Green, PhD plant pathologist **Ibadan***
J Legg PhD virologist **Namulonge**
A M Schilder PhD plant pathologist **Ibadan**

Associate experts

J C Meerman MSc **Ibadan**
M Zweigert Dipl Ing unit head (GTZ) **Cotonou**
A A Sy PhD regional coordinator **Cotonou**

Resource and Crop Management Division

D Baker PhD agricultural economist director **Yaounde***
H Grimme PhD soil chemist director **Ibadan**

Administration and Support Services

S Claassen MSc farm manager and station administrator **Yaounde***
Y Aboubakar station administration supervisor **Yaounde**

M Atangana MSc administrative officer **Yaounde**
E Ndingjock MBPA finance officer **Yaounde**
C Yumga station assistant **Yaounde**

Moist Savanna Program

R Carsky PhD agronomist program leader **Cotonou**
J Diels PhD modeler **Ibadan**
S Jagtap PhD crop modeler and GIS specialist **Ibadan**
G Kirchhof PhD soil physicist **Ibadan***
O Lyasse Ir agronomic engineer (VVOB) **Ibadan**
A M Manyong PhD agricultural economist **Ibadan**
N Sanginga PhD soil microbiologist **Ibadan**
S Tarawali PhD agronomist (joint with ILRI) **Ibadan**
G Tian PhD soil fertility specialist **Ibadan**
B P Vanlauwe PhD soil microbiologist **Ibadan**

Postdoctoral fellows

D Chikoye PhD weed scientist **Ibadan**
L S Koutika PhD soil scientist **Yaounde**

Visiting scientist

H Inaizumi PhD agricultural economist **Ibadan**

Humid Forest Program

S Weise PhD weed scientist and officer in charge Humid Forest Station **Yaounde**
A Adesina PhD agricultural economist and social science coordinator, **Yaounde**
S Hauser PhD soil physicist **Yaounde**
C Nolte PhD soil fertility specialist **Yaounde**
J Wendt PhD soil chemist **Yaounde**

Postdoctoral fellows

O Coulibaly PhD agricultural economist **Yaounde***
I Riviere PhD crop ecologist **Yaounde***

Visiting scientists

C Diaw PhD anthropologist (Rockefeller/ASB) **Yaounde**
J Gockowski PhD agricultural economist (Rockefeller/ASB) **Yaounde**
S E Johnson PhD agroecologist **Yaounde**

Analytical Services Unit

J Uponi MSc laboratory manager **Ibadan**

Biometrics Unit

S Nokoe, PhD biometrician unit head **Ibadan**

Postharvest Technology

Y W Jeon PhD agricultural engineer unit head **Ibadan**
L S Halos Kim MSc food and agricultural engineer **Ibadan**

Collaborative Study of Cassava in Africa (COSCA)

F I Nweke PhD agricultural economist/coordinator **Ibadan***

Visiting scientist

C L A Asadu PhD soil agronomist **Ibadan**

East and Southern Africa Regional Center (ESARC)

D R Vuylsteke Ir agronomist/breeder research team leader ESARC (CID) **Namulonge**
S Ferris PhD postharvest technologist ESARC (CID) **Namulonge**
C Gold PhD entomologist ESARC (PHMD) **Namulonge**

Professional Staff

S H O Okech PhD regional research fellow Africa Highlands
Namulonge

P Spejjer PhD nematologist ESARC (PHMD) **Namulonge**

Postdoctoral fellows

J Hartman PhD breeder/geneticist **Namulonge**

J Legg, PhD virologist **Namulonge**

Visiting scientist

N Wanyera PhD root crops agronomist **Namulonge**

Associate expert

E Boonen Ir VVOB **Namulonge** *

International Cooperation Division

M W Basse, PhD, mechanical engineer director **Ibadan**

Information Services

J Reeves JD head information services **Ibadan**

D R Mohan Raj, PhD head editorial unit **Ibadan**

R Zachmann PhD head multimedia unit **Ibadan**

Y Adedigba MA head library & documentation **Ibadan**

J I Adeyomoye MLS principal librarian **Ibadan**

O Ajayi BEd audiovisual specialist **Ibadan**

T Babaleye MCA, public affairs manager **Ibadan**

E O Ezomo MLS, principal librarian **Ibadan**

B O Ojuronbe BSc production manager **Ibadan***

Y Olatunbosun BSc, editor **Ibadan**

T T Owoeye MLS editor **Ibadan**

A Oyetunde MA editor **Ibadan**

R W Yussuf MA, production manager **Ibadan**

Training

J L Gulley PhD, program leader **Ibadan***

R Zachmann PhD head training materials **Ibadan**

A A Adekunle MSc research training specialist **Ibadan**

M Ajayi MSc research training specialist **Ibadan**

R Obubo MSc research training specialist **Ibadan**

C Okafor MBA, administrative manager **Ibadan**

O A Osinubi MSc, research training specialist **Ibadan**

Collaboration and Networks Program

A P Unyo PhD soil scientist, program leader, **Ibadan**

E A Atayi, PhD, agricultural economist, coordinator,

ecoregional program, **Ibadan**

J Fajemisin, PhD, breeder research liaison scientist **Bouake**

J Suh, PhD entomologist, research liaison scientist **Kumasi**

Interpretation and Translation Unit

B F Sall, MA, unit head **Ibadan**

C H Dia MA interpreter/translator **Ibadan**

O B Hounvou, MA, interpreter/translator, **Ibadan**

C Lord MA interpreter/translator, **Yaounde***

H Songre, MA interpreter/translator **Ibadan**

Monitoring and Evaluation Unit

O M Ogunyinka MSc coordinator **Ibadan**

Cooperative Special Projects

East Africa Root Crops Research Network (EARRNET)

J B A Whyte PhD breeder coordinator **Namulonge**

B W Khizzah PhD agronomist **Namulonge**

Southern Africa Root Crops Research Network (SARRNET)

J Teri PhD, pathologist coordinator, **Lilongwe**

M I Andrade PhD agronomist **Maputo**

N M Mahungu PhD breeder **Dar es Salaam**

T Malila, administrative officer, **Lilongwe**

A Mumba Kankolongo PhD pathologist **Mutanda**

I N Kasele PhD multiplication expert **Marondera**

West and Central Africa Collaborative Maize Research Network (WECAMAN)

B Badu Apraku PhD breeder coordinator **Bouake**

Resident scientists of IARCs/AROs

H A Mendoza PhD, breeder CIP **Ibadan**

O Ndoye PhD economist CIFOR **Yaounde**

M Bernard soil scientist coordinator Hohenheim students (UNIHO) **Cotonou**

B Duguma PhD agroforester ICRAF **Yaounde**

K Agyemang PhD animal scientist ILRI **Ibadan**

A Larbi PhD agronomist ILRI **Ibadan**

M Peters PhD agronomist ILRI **Ibadan***

S Tarawali PhD agronomist ILRI **Ibadan**

J W Smith PhD animal scientist ILRI **Ibadan**

B N Singh PhD breeder WARDA **Ibadan***

C T Williams PhD entomologist WARDA/CABI **Ibadan***

Corporate Services Division

W Powell BSc director corporate services division **Ibadan**

Human Resources

O I Osotimehin BSc DDA personnel manager **Ibadan**

J B Adenuga, AMIPN employee services manager **Ibadan**

A Ohanwusi MBA planning/training manager **Ibadan**

Medical Unit

F O A Ajose MD unit head **Ibadan**

I I Anetor MSc medical laboratory services officer **Ibadan**

E O Lawani MD physician **Ibadan**

F O Dwyer BSc nurse/practitioner **Ibadan**

Budget and Finance

G McIntosh CMA chief financial officer **Ibadan**

B A Adeola, FCIS accountant **Ibadan**

L O Afemikhe MSc senior technical analyst (computer)

Ibadan

J E Bolarinwa MBA payroll accountant **Ibadan**

N N Eguozie BSc financial information systems manager

Ibadan

R Obikudu, MBA materials manager **Ibadan**

O Oduberu ICAN FIS senior analyst **Ibadan**

S Ogunade treasury manager **Ibadan**

K O Olatifede ACA special projects accountant **Ibadan**

O Sholola ACA corporate accounting manager **Ibadan**

S J Udoh AMNIM chief accountant **Ibadan**

Left during the year

Internal Audit

R S Lafond CMA head **Ibadan**

A Ajakaiye ACA intermediate auditor **Ibadan**

R A Fagbenro ACA senior internal auditor, **Ibadan**

Physical Plant Services

A Bhatnagar BSc, head **Ibadan**

A C Butler, building and site services officer **Ibadan***

P G Gualinetti, construction site engineering services officer

Ibadan*

A C Micheletti construction site engineer/services officer

Ibadan

S W Quader BSc electronic services officer **Ibadan**

E O Akintokun research vehicle services officer **Ibadan**

F K Alude HND heavy equipment & fabrication services

manager **Ibadan**

P T Lamuren BSc telecommunication services officer **Ibadan**

M A Oyediji electrical services officer **Ibadan**

O O Taiwo HND assistant services officer **Ibadan**

Computer Services

M C A Simmonds, MSc manager **Ibadan**

S A Folayan MSc computer systems manager **Ibadan***

Auxiliary Services

A Beard manager hotel and catering services **Ibadan**

C Eykyn, BSc specialist science teacher, **Ibadan**

C Inniss Palmer MA specialist english teacher **Ibadan**

A Jackson deputy head international school of IITA **Ibadan**

N Jackson head international school of IITA **Ibadan**

A R Middleton BA manager hotel and catering services

Ibadan*

D J Sewell manager aircraft operations **Ibadan**

W Ekpo ACIS security manager **Ibadan**

C A Enahoro HND head, travel services **Lagos**

Abbreviations and acronyms used in the text

ASB	Global Initiative on the Alternatives to Slash-and Burn
BC	Biological control
CABI	Commonwealth Agriculture Bureau International, UK
CBO	Community based Organization
CCER	Center Commissioned External Review
CGIAR	Consultative Group on International Agricultural Research
CIAT	Centro Internacional de Agricultura Tropical (Colombia)
CIMMYT	Centro Internacional de Mejoramiento de Maiz y Trigo (Mexico)
CIRAD	Centre de cooperation internationale en recherche agronomique pour le developpement
CORAF	Conference des responsables de la recherche agronomique africains
DM	Downy mildew
EARRNET	East African Root and Tubers Research Network
EPHTA	Ecoregional Program for the Humid and Subhumid Tropics of Sub-Saharan Africa
FAO	Food and Agriculture Organization of the United Nations
GIS	geographic information system
GTZ	German Agency for Technical Cooperation
IARC	international agricultural research center
ICIPE	International Centre for Insect Physiology and Ecology
ICRISAT	International Crops Research Institute for the Semi-Arid Tropics
IDRC	International Development Research Centre (Canada)
IIBC	International Institute of Biological Control (UK)
ILRI	International Livestock Research Institute
INTG	IARC/NARS training group
IPM	integrated pest management
KARI	Kenya Agricultural Research Institute
LUBILOSA	Lutte biologique contre les locustes et sauteriaux
MPB	Maruca pod borer
NARES	national agricultural research and extension systems
NARS	national agricultural research system
NGO	nongovernmental organization
NRI	Natural Resources Institute (UK)
PEDUNE	Protection ecologiquement durable du niebe
PSB	Pod-sucking bug
SADC	Southern Africa Development Community
SARRNET	Southern Africa Root Crops Research Network
SP-IPM	Systemwide Program on Integrated Pest Management
USAID	United States Agency for International Development
WARDA	West Africa Rice Development Association
WECAMAN	West and Central Africa Collaborative Maize Research Network

IITA Annual Report 1997

ISSN 0331-4340

EDITOR-IN-CHIEF AND WRITER

Jack Reeves

MANAGING EDITOR

Taiwo Owoeye

DESIGN AND LAYOUT

Remi Yussuf

EDITING

Taiwo Owoeye
D R Mohan Raj
Jack Reeves

PUBLISHING CONSULTANT

Alan Mark Fletcher

PRINTING

Finger Lakes Press
Auburn, New York

COVER DRAWING

'Farming Mothers'
Ronke Lawal-Campbell

International Institute of Tropical Agriculture
PMB 5320
Ibadan Nigeria

Telephone (234 2) 241 2626
Fax (234 2) 241 2221
E mail iita@cgnnet.com

International Mailing Address

c/o L W Lambourn & Co
Carolyn House 26 Dingwall Road
Croydon CR9 3EE England

97