

**Indonesia
Demographic
and Health
Survey
2007**

**Preliminary
Report**

**Statistics Indonesia
Jakarta, Indonesia**

**National Family Planning Coordinating Board
Jakarta, Indonesia**

**Ministry of Health
Jakarta, Indonesia**

**MEASURE DHS
Macro International
Calverton, Maryland USA**

This report summarizes the findings of the 2007 Indonesia Demographic and Health Survey (IDHS) carried out by Statistics Indonesia (Badan Pusat Statistik—BPS). The IDHS is part of the worldwide Demographic and Health Surveys program, which is designed to collect data on fertility, family planning, and maternal and child health.

The Government of Indonesia supported the local costs of the survey. The United Nations Population Fund (UNFPA) provided funds for questionnaires printing and shipment. Macro International Inc. (Macro) provided limited technical assistance under the auspices of the Demographic and Health Surveys (MEASURE DHS) program, which is supported by the U.S. Agency for International Development (USAID). The Ford Foundation provided funds for the expansion of the sample in 15 districts in Java, to allow estimates at the district level. UNICEF also provided funds to allow estimates at the district level in Nanggroe Aceh Darussalam Province and two districts Nias and South Nias in North Sumatera Province.

Additional information about the survey may be obtained from the Directorate for Population and Labor Force Statistics, BPS, Jalan Dr. Sutomo No. 6-8, Jakarta 10710, Indonesia (Telephone/fax 345-6285, e-mail: Kependudukan@mailhost.bps.go.id), or the National Family Planning Coordinating Board, BKKBN, Jalan Permata 1, Halim Perdanakusumah, Jakarta 13650, Indonesia (Telephone/fax 800-8535), or the Institute for Research and Development, Ministry of Health, Jalan Percetakan Negara 29, Jakarta 10560, Indonesia.

Additional information about the DHS program may be obtained by contacting: MEASURE DHS, Macro International Inc., 11785 Beltsville Drive, Suite 300, Calverton, MD 20705, USA (Telephone 301-572-0200; Fax 301-572-0999; E-mail: reports@macrointernational.com; Internet: www.measuredhs.com).

Indonesia Demographic and Health Survey 2007

Preliminary Report

**Statistics Indonesia
Jakarta, Indonesia**

**National Family Planning Coordinating Board
Jakarta, Indonesia**

**Ministry of Health
Jakarta, Indonesia**

**MEASURE DHS
Macro International Inc.
Calverton, Maryland USA**

July 2008



CONTENTS

Page

TABLES	v
FIGURES	vii
I. INTRODUCTION	1
II. SURVEY IMPLEMENTATION	3
A. Questionnaires	3
B. Sample Design	3
C. Training and Fieldwork	4
D. Data Processing.....	4
III. RESULTS	5
A. Response Rates	5
B. Characteristics of the Respondents	6
C. Fertility	7
D. Family Planning	8
E. Unmet Need for Family Planning Services	11
F. Fertility Preferences	12
G. Maternity Care	13
H. Vaccination of Children	15
I. Childhood Illnesses	17
J. Infant Feeding Practices.....	18
K. Childhood Mortality	19
L. Knowledge of HIV/AIDS	21
REFERENCES	25
APPENDIX	27

TABLES

		Page
Table 1	Results of the household and individual interviews.....	5
Table 2	Background characteristics of respondents.....	6
Table 3	Current Fertility	7
Table 4	Knowledge of contraceptive methods by background characteristics	9
Table 5	Current use of contraception.....	10
Table 6	Need and demand for family planning among currently married women	12
Table 7	Fertility preferences by number of living children	13
Table 8	Maternal care indicators.....	14
Table 9	Vaccinations by background characteristics	16
Table 10	Treatment for acute respiratory infection, fever, and diarrhea.....	18
Table 11	Breastfeeding status by age.....	19
Table 12	Early childhood mortality rates.....	20
Table 13	Knowledge of AIDS	21
Table 14	Knowledge of ways to reduce the risk of getting the AIDS virus	22
Table 15.1	Knowledge about voluntary counseling and testing (VCT): Women.....	23
Table 15.2	Knowledge about voluntary counseling and testing (VCT): Men	24
Table A.1	Background characteristics of respondents by province.....	27
Table A.2	Fertility by province.....	28
Table A.3	Knowledge of contraceptive methods by province.....	29
Table A.4	Current use of contraception by province	30
Table A.5	Unmet need and the demand for family planning by province.....	31
Table A.6	Maternal care indicators by province.....	32
Table A.7	Vaccinations by province.....	33
Table A.8	Treatment of acute respiratory infection, fever, and diarrhea by province	34
Table A.9	Early childhood mortality rates by province.....	35
Table A.10	Knowledge of ways to reduce the risk of getting the AIDS virus	36
Table A.11	Knowledge of HIV prevention methods by province	37
Table A.12	Knowledge about voluntary counseling and testing (VCT).....	38

FIGURES

	Page
Figure 1	Trends in total fertility rates, Indonesia 1991-2007 8
Figure 2	Trends in contraceptive use among currently married women, Indonesia 1991-2007 11
Figure 3	Trends in maternal health care indicators, Indonesia 2002-03 and 2007 15
Figure 4	Trends in vaccination coverage among children age 12-23 months, Indonesia 2002-03 and 2007 17
Figure 5	Trends in early childhood mortality rates, Indonesia 1991-2007 20

I. INTRODUCTION

The 2007 Indonesia Demographic and Health Survey (IDHS) was implemented by Statistics Indonesia (Badan Pusat Statistik—BPS). The Government of Indonesia supported the local costs of the survey. The United Nations Population Fund (UNFPA) provided funds for questionnaires printing and shipment. Macro International Inc. (Macro) provided limited technical assistance under the auspices of the Demographic and Health Surveys (MEASURE DHS) program, which is supported by the U.S. Agency for International Development (USAID). The Ford Foundation provided funds for the expansion of the sample in 15 districts in Java, to allow estimates at the individual district level. UNICEF also provided funds to allow estimates at the district level in Nanggroe Aceh Darussalam Province and two districts Nias and South Nias in North Sumatera Province.

The 2007 IDHS is the sixth survey conducted in Indonesia under the auspices of the DHS program. Most of the data collected in the 2007 IDHS provide updated estimates of basic demographic and health indicators covered in previous IDHS surveys.

The 2007 IDHS is designed to provide information on population, family planning, and health. A scientifically selected sample of ever-married women age 15 to 49 years and currently married men age 15-54 were interviewed. Women were asked questions about their background, the children they had given birth to, their knowledge and use of family planning methods, the health of their children, reproductive health, and other information that will be helpful to policymakers and administrators in the health and family planning fields. The questionnaire for men was shorter than that for women because it excluded detailed questions on individual children and children's health. However, men were asked about their knowledge, attitudes and practices regarding health care for their wife and children.

This report presents a first look at selected findings of the 2007 IDHS.¹ A comprehensive analysis of the data will be published later this year. While considered provisional, the results presented here are not expected to differ significantly from those presented in the final report.

¹ The 2007 Indonesia Young Adult Reproductive Health Survey (IYARHS), a survey on the reproductive attitudes and behavior of the never-married population age 15-24, was conducted at the same time as the IDHS in a subsample of the households. A special report on the IYARHS findings will be prepared and issued at the same time as the IDHS main report.

II. SURVEY IMPLEMENTATION

A. Questionnaires

The 2007 IDHS used three questionnaires: the Household Questionnaire, the Women's Questionnaire and the Men's Questionnaire. In consultation with the National Family Planning Coordinating Board (BKKBN) and the Ministry of Health (MOH), BPS made a decision to largely base the 2007 IDHS survey instruments on the questionnaires used in the 2002-03 IDHS in order to facilitate trend analysis. Input was solicited from other potential data users, and several modifications were made to optimize the draft 2007 IDHS instruments to collect the needs for population and health data. The draft IDHS questionnaires were also compared to the most recent version of the standard questionnaires used in the DHS program and minor modifications incorporated to facilitate international comparison.

The Household Questionnaire was used to list all the usual members and visitors in the selected households. Basic information collected on each person listed includes: age, sex, education, and relationship to the head of the household. The main purpose of the Household Questionnaire was to identify women and men who were eligible for the individual interview. Information on characteristics of the household's dwelling unit, such as the source of water, type of toilet facilities, construction materials used for the floor and outer walls of the house, and ownership of various durable goods were also recorded in the Household Questionnaire. These items reflect the household's socioeconomic status.

The Women's Questionnaire was used to collect information from all ever-married women age 15-49. These women were asked questions on the following topics:

- Background characteristics (marital status, education, media exposure, etc.)
- Knowledge and use of family planning methods
- Reproductive history and fertility preferences
- Antenatal, delivery and postnatal care
- Breastfeeding and infant feeding practices
- Vaccinations and childhood illnesses
- Marriage and sexual activity
- Woman's work and husband's background characteristics
- Infant's and children's feeding practices
- Childhood mortality
- Awareness and behavior regarding AIDS and other sexually transmitted infections (STIs)
- Sibling mortality, including maternal mortality.

The Men's Questionnaire was administered to all currently married men age 15-54 living in every third household in the IDHS sample. The Men's Questionnaire collected much of the same information included in the Women's Questionnaire, but was shorter because it did not contain questions on reproductive history, maternal and child health, nutrition and maternal mortality. Instead, men were asked about their knowledge and participation in the health-care-seeking practices for their children.

B. Sample Design

The sampling frame for this survey is the list of census blocks (CBs) developed for the 2007 National Labor Force Survey (Sakernas). A total of 1,694 CBs, 676 in urban areas and 1,018 in rural areas were selected from the list of CBs. The number of CBs selected in each district was not allocated proportional to their total population. For the Sakernas, in each CB, a complete household listing and

mapping was conducted in July 2007, which formed the basis for the second-stage sampling. An average of 25 households were selected systematically from each CB.

The 2007 IDHS sample is aimed at providing reliable estimates of key characteristics for ever-married women 15-49 and married men 15-54 in Indonesia as a whole, in urban and rural areas, and in each of the 33 provinces included in the survey. The sample was targeted to yield a total of 33,850 completed interviews with women and 12,200 interviews with men.

C. Training and Fieldwork

A total of 832 persons, 468 women and of 364 men, participated in the main survey training for interviewers. Training took place in June 2007 in seven training centers between (Medan, Padang, Banten, D.I. Yogyakarta, Denpasar, Banjarmasin, and Makasar), and in July 2007 in two training centers (Jayapura and Manokwari). The training included class presentations, mock interviews, and tests. All of the participants were trained using the Women's Questionnaire. Once the materials for the women's interview were completed, the male participants were trained in conducting an interview using the Men's Questionnaire. The training included practice interviews in Bahasa Indonesia and the participant's local language.

The 2007 IDHS employed 104 interviewing teams to collect the data. Fieldwork principally took place from 25 June to 31 December, 2007. However, fieldwork had to be extended in several provinces including Riau Islands, Papua and West Papua because of flooding and other problems. Fieldwork was completed in all areas in February 2008.

D. Data Processing

All completed questionnaires for the IDHS, accompanied by their control forms, were returned to the BPS central office in Jakarta for data processing. This consisted of office editing, coding of open-ended questions, data entry, verification, and editing computer-identified errors. A team of 42 data entry clerks, data editors and data entry supervisors processed the data. Data entry and editing was carried using a computer package program called CSPro, which was specifically designed to process DHS-type survey data. During the preparation of the data entry programs, a BPS staff spent several weeks at Macro International offices in Calverton, Maryland. Data entry and editing activities, which began in September 2007 were completed in March 2008.

III. RESULTS

This section of the report focuses on key findings from the 2007 IDHS in the areas of fertility and family planning, maternal and child health, and HIV/AIDS awareness. The results in the main body of the report focus on the findings for Indonesia as a whole. Additional tables showing findings by province are included in the appendix of this report.

A. Response Rates

Table 1 shows response rates for the 2007 IDHS. A total of 42,341 households were selected in the sample, of which 41,131 were occupied. Of these households, 40,701 were successfully interviewed, yielding a household response rate of 99 percent.

In the interviewed households, 34,227 women were identified for individual interview and, of these, completed interviews were conducted with 32,895 women, yielding a response rate of 96 percent. In half of the households, 9,716 eligible men were identified, of which 8,758 were successfully interviewed, yielding a response rate of 90 percent. The lower response rate for men was due to the more frequent and longer absence of men from the household.

Table 1 Results of the household and individual interviews			
Number of households, number of interviews, and response rates, according to residence (unweighted), Indonesia 2007			
Result	Residence		Total
	Urban	Rural	
Household interviews			
Households selected	16,920	25,421	42,341
Households occupied	16,429	24,702	41,131
Households interviewed	16,224	24,477	40,701
Household response rate ¹	98.8	99.1	99.0
Individual interviews: women			
Number of eligible women	13,608	20,619	34,227
Number of eligible women interviewed	13,087	19,808	32,895
Eligible women response rate ²	96.2	96.1	96.1
Individual interviews: men			
Number of eligible men	3,927	5,789	9,716
Number of eligible men interviewed	3,510	5,248	8,758
Eligible men response rate ²	89.4	90.7	90.1
¹ Households interviewed / households occupied			
² Respondents interviewed / eligible respondents			

B. Characteristics of the Respondents

Table 2 shows the distribution of ever-married women age 15-49 and currently married men age 15-54 in the 2007 IDHS sample by selected background characteristics. Approximately one in three women are under 30 years of age, 94 percent are currently married, and 42 percent live in urban areas. Over time, women are becoming better educated, but the change is diminishing in recent years. The percentage of women with no education declined from 13 percent in 1997 to 8 percent in 2002-2003 and 7 percent in 2007. Over the same time period, the percentage of women with some secondary education increased from 38 percent to 46 percent.

Among the currently married men who were interviewed in the survey, 17 percent are under 30 years, 36 percent are in their thirties, and 47 percent are age 40-54. In general, men are better educated than women. The percentage of men with no education is lower than that of women, while the percentage of men with secondary or higher education is higher than that of women.

Table 2 Background characteristics of respondents						
Percent distribution of ever-married women age 15-49 and currently married men age 15-54 by selected background characteristics, Indonesia 2007						
Background characteristic	Women			Men		
	Weighted percent	Weighted number	Unweighted number	Weighted percent	Weighted number	Unweighted number
Age						
15-19	2.6	845	914	0.3	29	19
20-24	12.4	4,094	4,156	4.9	432	442
25-29	17.5	5,771	6,170	12.7	1,116	1,190
30-34	18.3	6,020	6,317	16.2	1,418	1,587
35-39	18.3	6,004	5,898	19.2	1,679	1,708
40-44	16.3	5,365	5,034	17.9	1,570	1,500
45-49	14.6	4,795	4,406	15.5	1,359	1,268
50-54	na			13.2	1,155	1,044
Marital status						
Married	94.0	30,931	30,869	100.0	8,758	8,758
Divorced	3.1	1,012	1,035	na	na	na
Widowed	2.9	952	991	na	na	na
Residence						
Urban	41.8	13,745	13,087	42.6	3,728	3,510
Rural	58.2	19,150	19,808	57.4	5,030	5,248
Education						
No education	6.9	2,270	2,234	4.1	361	344
Some primary	16.9	5,572	5,503	18.3	1,605	1,444
Completed primary	30.6	10,077	8,834	26.7	2,339	2,084
Some secondary	20.6	6,781	7,048	19.7	1,721	1,868
Secondary +	24.9	8,193	9,273	31.1	2,727	3,016
Religion						
Islam	88.5	29,104	26,185	88.2	7,724	6,881
Protestant	6.0	1,989	3,598	6.1	531	990
Catholic	2.9	958	1,406	3.0	263	368
Hindu	1.8	592	1,286	2.0	174	408
Buddha	0.4	139	253	0.4	33	62
Confucian	0.2	74	120	0.1	7	12
Missing	0.1	38	47	0.1	8	11
Total	100.0	32,895	32,895	100.0	8,758	8,758

Note: Education categories refer to the highest level of education attended, whether or not that level was completed.
na = Not applicable

C. Fertility

All women who were interviewed in the 2007 IDHS were asked to report the total number of sons and daughters to whom they had ever given birth in their lifetime. To promote complete reporting of children, women were asked separately about children still living at home, those living elsewhere, and children who had died. A complete birth history was then obtained, including information on the sex, date of birth and survival status of each child and the age at death for dead children.

Age-specific fertility rates for the three-year period before the 2007 IDHS are shown in Table 3. Age-specific and total fertility rates were calculated directly from the birth history data. The sum of the age-specific fertility rates (known as the total fertility rate, or TFR) is a summary measure of the level of fertility. It represents the number of children a woman would have by the end of her childbearing years if she were to pass through those years bearing children at the currently observed age-specific rates. If fertility were to remain constant at current levels, a woman in Indonesia would bear an average of 2.6 children in her lifetime. Table 3 also shows that, overall, urban women have 0.6 fewer children than women in rural areas. This is true for age-specific fertility rates in all age groups but 25-29.


Table 3 also shows the general fertility rate (GFR) and crude birth rate (CBR). As in the case of the TFR, fertility levels in urban areas are lower than those in rural areas.

Table 3 Current fertility			
Age-specific and total fertility rate, the general fertility rate, and the crude birth rate for the three years preceding the survey, by residence, Indonesia 2007			
Age group	Residence		Total
	Urban	Rural	
15-19	26	74	51
20-24	116	153	135
25-29	138	131	134
30-34	104	110	108
35-39	59	70	65
40-44	17	21	19
45-49	4	7	6
TFR	2.3	2.8	2.6
GFR	80.0	97.0	89.0
CBR	20.2	21.5	20.9
TFR ¹	2.3	2.8	2.6
GFR ¹	80.0	97.0	89.0
CBR ¹	20.3	21.5	21.0

TFR: Total fertility rate for ages 15-49, expressed per woman
 GFR: General fertility rate (births divided by the number of women age 15-44), expressed per 1,000 women
 CBR: Crude birth rate, expressed per 1,000 population
¹ Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua). Rates for age group 45-49 may be slightly biased due to truncation.

Figure 1 shows total fertility rates from the 1991, 1994, 1997, 2002-03, and 2007 IDHS surveys. There is a steady decline from 3 children per woman in 1991 to 2.6 children at the time of the 2002-2003 IDHS. Fertility appears to have stabilized at this level, with no change observed during the roughly five-year period between 2002-2003 and 2007.

**Figure 1. Trends in Total Fertility Rates
Indonesia 1991-2007**



Note: The 2002-03 IDHS did not include Naggroe Aceh Darussalam, Maluku North Maluku, and Papua provinces. Previous surveys included East Timor.

D. Family Planning

Knowledge of Contraception

Acquiring knowledge about fertility control is an important step towards gaining access to and then using a suitable contraceptive method in a timely and effective manner. The 2007 IDHS collected information on knowledge and use of contraception. To obtain these data, respondents were first asked to name all of the contraceptive methods that they had heard about. For methods not mentioned spontaneously, a description of the method was read, and the respondents were asked if they had heard of the method. For each method named or recognized, respondents were asked if they had ever used the method. Finally, women were asked if they (or their partners) were currently using a method. For analytical purposes, contraceptive methods were grouped into two types: modern and traditional. Modern methods include female sterilization, male sterilization, pill, intrauterine device (IUD), injectables, implants, male condom, diaphragm, emergency contraception, and lactational amenorrhea method (LAM). Traditional methods include periodic abstinence (rhythm method), withdrawal, and folk methods.

Table 4 shows that knowledge of contraception is high among currently married women. Almost all respondents know at least one method of contraception. Knowledge of a modern method is also nearly universal (98 percent). The pill and injectables are known by almost all married women (95 and 96 percent, respectively). The next most well known modern methods are implants and IUD (86 and 84 percent, respectively). Periodic abstinence is the most widely known traditional method (39 percent).

Table 4 Knowledge of contraceptive methods by background characteristics

Percentage of currently married women age 15-49 know any contraceptive method by background characteristics, Indonesia 2007

Background characteristic	Modern method											Traditional method			Number of women		
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	IUD	Injectables	Implants	Male condom	Intra-vag/dia-phragm	Emergency contraception	LAM	Any traditional method	Periodic abstinence		Withdrawal	Folk method
Age																	
15-19	97.2	97.0	41.5	20.2	92.1	57.2	93.1	73.0	62.4	6.3	2.7	12.6	24.8	14.8	16.8	3.9	814
20-24	98.9	98.7	58.2	30.2	95.9	76.7	97.5	82.9	76.2	11.0	5.6	19.4	42.5	33.3	28.1	4.1	3,952
25-29	99.1	98.8	64.8	39.2	96.6	84.8	98.1	87.6	80.5	13.5	7.3	24.5	51.8	41.3	36.4	5.0	5,585
30-34	99.3	99.0	70.1	42.3	96.9	88.0	98.4	88.9	82.2	14.6	7.5	26.5	53.9	43.7	38.4	6.0	5,765
35-39	99.1	98.9	71.9	43.7	95.6	88.1	97.0	90.4	78.6	14.8	6.2	24.9	52.3	42.0	35.1	7.7	5,704
40-44	98.1	97.8	68.5	42.0	93.7	85.5	95.0	86.9	74.7	16.0	6.7	24.7	48.5	39.6	33.4	6.9	4,899
45-49	96.7	96.3	63.9	39.0	91.6	81.4	92.8	81.1	68.3	12.5	5.2	19.1	41.0	31.4	27.5	6.1	4,211
Residence																	
Urban	99.5	99.5	78.0	51.6	98.1	91.9	98.7	90.9	88.7	20.4	9.1	33.4	63.0	55.1	44.8	7.0	12,842
Rural	97.9	97.5	57.6	30.6	93.0	78.2	95.0	83.2	68.4	9.0	4.5	16.1	38.0	26.8	25.1	5.3	18,089
Education																	
No education	91.9	89.5	37.7	17.2	76.8	55.5	82.2	62.0	37.4	3.3	1.8	9.2	16.9	7.9	8.7	5.4	2,002
Some primary Completed	97.3	97.0	50.5	22.7	91.2	74.0	93.6	78.3	57.6	6.2	2.8	10.8	25.9	15.9	15.6	4.0	5,112
primary	98.9	98.8	62.0	33.0	95.6	82.1	97.1	86.1	72.9	7.7	3.3	15.0	37.6	26.2	23.5	4.8	9,511
Some secondary	99.5	99.5	70.7	40.2	98.1	87.7	98.9	90.5	86.3	13.3	6.2	24.2	53.8	42.3	35.9	6.2	6,494
Secondary +	99.9	99.8	84.8	63.0	99.2	96.6	99.4	94.8	96.5	29.0	14.0	44.6	79.9	73.1	60.9	8.8	7,810
Living children																	
0	95.9	95.8	55.5	35.3	92.7	73.7	92.9	76.2	73.9	13.2	7.0	18.7	43.7	36.0	31.4	4.1	2,488
1-2	99.2	99.0	67.7	40.0	96.4	85.5	97.4	87.9	79.2	14.2	6.6	24.8	49.6	40.2	34.2	5.7	17,447
3-4	98.8	98.4	69.0	41.8	94.7	85.5	96.6	88.0	76.2	14.3	6.2	23.8	49.8	39.1	33.8	7.1	8,396
5+	96.5	96.1	56.4	31.2	90.3	77.6	93.8	80.5	65.6	9.5	5.2	15.8	40.3	28.2	27.1	6.1	2,600
Total	98.6	98.3	66.1	39.4	95.1	83.9	96.5	86.4	76.8	13.7	6.4	23.3	48.4	38.5	33.3	6.0	30,931

LAM = Lactational amenorrhea method

Current Use of Family Planning

Table 5 shows that 61 percent of married women age 15-49 in Indonesia use a method of family planning. Most of these women use a modern method of contraception (57 percent), with use of traditional methods being quite limited. As in the 2002-03 IDHS, the most popular modern methods are injectables (32 percent), the pill (13 percent), and IUD (5 percent). Three percent of women each use female sterilization and implants. The injectables continue to gain popularity (28 percent in 2002-03 to 32 percent in 2007). Over the same period, the use of other methods, either stayed at the same level (the pill 13 percent), or slightly declined (e.g., female sterilization and implants from 4 percent to 3 percent, and IUD from 6 percent to 5 percent).

Use of contraception varies by the women's characteristics. The rate is low among women in the youngest and oldest age groups, and peaks at age 30-34. Older women are less likely to use a modern method and more likely to use traditional methods. Permanent methods (female and male sterilization) or long-term methods like the IUD are more popular among older women.

Table 5 Current use of contraception

Percent distribution of currently married women by contraceptive method currently used, according to background characteristics, Indonesia 2007

Background characteristic	Modern method										Traditional method					Number of women	
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	IUD	Injectables	Implants	Male condom	LAM	Any traditional method	Periodic abstinence	Withdrawal	Folk method	Not currently using		Total
Age																	
15-19	46.8	46.2	0.0	0.0	10.0	0.0	34.8	1.0	0.2	0.2	0.7	0.1	0.5	0.1	53.2	100.0	814
20-24	61.5	59.9	0.0	0.0	12.0	1.3	43.9	2.1	0.6	0.1	1.6	0.3	1.1	0.2	38.5	100.0	3,952
25-29	64.1	60.7	0.2	0.0	14.1	2.6	40.3	2.5	1.0	0.0	3.3	0.9	2.1	0.3	35.9	100.0	5,585
30-34	69.1	64.7	1.3	0.1	14.9	4.2	39.3	3.1	1.7	0.1	4.4	1.7	2.2	0.5	30.9	100.0	5,765
35-39	68.6	63.0	3.8	0.2	16.3	6.2	30.7	4.1	1.6	0.0	5.5	2.7	2.4	0.5	31.4	100.0	5,704
40-44	59.6	54.6	6.6	0.5	12.9	7.9	21.7	3.0	2.0	0.0	5.0	2.1	2.5	0.5	40.4	100.0	4,899
45-49	42.1	38.2	7.4	0.5	7.9	8.1	11.8	1.5	1.0	0.0	3.9	1.2	2.2	0.5	57.9	100.0	4,211
Residence																	
Urban	62.5	57.1	4.0	0.2	13.9	6.7	28.0	1.8	2.4	0.0	5.3	2.3	2.6	0.4	37.5	100.0	12,842
Rural	60.6	57.5	2.3	0.2	12.8	3.6	34.5	3.5	0.5	0.0	3.0	0.9	1.7	0.4	39.4	100.0	18,089
Education																	
No education	42.3	40.1	2.5	0.8	7.1	3.6	21.7	4.2	0.3	0.0	2.2	0.1	0.9	1.2	57.7	100.0	2,002
Some primary	54.0	51.5	4.1	0.2	11.7	3.8	27.9	3.3	0.4	0.0	2.6	0.5	1.6	0.5	46.0	100.0	5,112
Compl. primary	64.0	61.1	2.2	0.2	14.7	3.7	36.9	2.9	0.4	0.0	2.9	0.7	1.9	0.3	36.0	100.0	9,511
Some secondary	65.5	61.4	2.8	0.1	13.9	3.3	37.8	2.4	1.1	0.0	4.1	1.6	2.3	0.3	34.5	100.0	6,494
Secondary +	64.4	57.8	3.6	0.2	13.6	8.8	26.0	2.1	3.4	0.1	6.6	3.4	2.8	0.3	35.6	100.0	7,810
Living children																	
0	8.3	7.9	0.0	0.0	3.5	0.0	4.1	0.0	0.1	0.0	0.4	0.1	0.3	0.0	91.7	100.0	2,488
1-2	68.1	64.3	1.1	0.1	14.5	5.4	38.7	3.0	1.4	0.0	3.7	1.5	1.9	0.3	31.9	100.0	17,447
3-4	67.7	62.5	6.7	0.3	15.1	5.9	29.6	3.2	1.6	0.0	5.2	1.9	2.6	0.7	32.3	100.0	8,396
5+	46.7	41.5	7.1	0.5	8.0	2.7	19.3	2.7	1.0	0.0	5.2	1.4	3.3	0.5	53.3	100.0	2,600
Total	61.4	57.4	3.0	0.2	13.2	4.9	31.8	2.8	1.3	0.0	4.0	1.5	2.1	0.4	38.6	100.0	30,931

Note: If more than one method is used, only the most effective method is considered in this tabulation.
LAM = Lactational amenorrhea method.

Contraceptive use varies by residence. Urban women are not only more likely to use contraception, but also tend to use long-term methods like female sterilization and IUD. Rural women, on the other hand, are more likely to use injectables and implants.


Contraceptive use increases with women's level of education. While less than half of married women with no formal education are currently using a method, at least 64 percent of women who completed primary or higher education are using a method of family planning.

The association between contraceptive use and a woman's parity is obvious in Table 5. As expected, women with children are more likely to use contraception than women with no children. Whereas few women with no children are using a family planning method, two in three women with one to four children are using contraception.

It has been observed in previous IDHS surveys that traditional methods, especially periodic abstinence, are more popular among urban women, better educated women, and women with higher parity.

Figure 2 shows the trends in contraceptive use between 1991 and 2007. There has been a gradual and steady increase in contraceptive prevalence rate among currently married women in Indonesia from 50 percent in 1991 to 61 percent in 2002-03. In the past five years, however, the increase has been minimal (60 percent in 2002-03 to 61 percent in 2007).

Figure 2 Trends in Contraceptive Use among Currently Married Women, Indonesia 1991-2007



E. Unmet Need for Family Planning Services

Table 6 presents data on the level of unmet need, met need, and the total demand for family planning services for currently married women age 15-49 by background characteristics. Nine percent of currently married women in Indonesia have an unmet need for family planning; 4 percent are in need because they want no more children and 5 percent because they would like to delay the next birth for two or more years. The total met need for family planning (i.e., the proportion currently using contraception) is 61 percent; 36 percent of married women are using contraception to limit the number of children and 25 percent for spacing purposes. Taking both the met and unmet need into account, the total demand for family planning among currently married women in Indonesia is 70 percent, of which 87 percent satisfied. If this demand is satisfied, the contraceptive prevalence in Indonesia among currently married women will be 71 percent instead of 61 percent.

The total unmet need for family planning services in Indonesia has remained unchanged since the 1997 IDHS (BPS and ORC Macro, 2003). The overall demand for family planning in 2002-03 was 70 percent, of which 88 percent had been satisfied.

Table 6 Need and demand for family planning among currently married women

Percentage of currently married women age 15-49 with unmet need for family planning, percentage with met need for family planning, the total demand for family planning, and the percentage for the demand for contraception that is satisfied, by background characteristics, Indonesia 2007

Background characteristic	Unmet need for family planning ¹			Met need for family planning (currently using) ²			Total demand for family planning			Percentage of demand satisfied	Number of women
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total		
Age											
15-19	7.3	2.5	9.8	41.8	5.1	46.8	49.1	7.6	56.6	82.7	814
20-24	7.9	1.9	9.8	51.3	10.2	61.5	59.4	12.1	71.5	86.3	3,952
25-29	7.7	1.9	9.6	44.5	19.5	64.1	52.5	21.5	74.0	87.1	5,585
30-34	5.3	3.9	9.2	31.5	37.6	69.1	36.9	41.6	78.5	88.3	5,765
35-39	2.7	6.8	9.5	14.6	54.0	68.6	17.4	61.0	78.4	87.9	5,704
40-44	1.0	8.6	9.6	4.1	55.5	59.6	5.1	64.1	69.2	86.1	4,899
45-49	0.5	5.5	6.0	1.4	40.7	42.1	1.9	46.3	48.2	87.5	4,211
Residence											
Urban	4.0	4.7	8.8	23.7	38.8	62.5	27.9	43.6	71.5	87.7	12,842
Rural	4.5	4.7	9.2	26.1	34.5	60.6	30.7	39.3	70.0	86.8	18,089
Education											
No education	3.0	7.7	10.6	8.7	33.6	42.3	11.7	41.3	53.0	80.0	2,002
Primary	3.8	5.3	9.1	22.4	38.1	60.5	26.3	43.6	69.9	86.9	14,623
Secondary	5.2	3.9	9.0	31.0	34.1	65.2	36.4	38.0	74.5	87.9	12,169
More than secondary	4.2	2.8	7.0	24.9	38.4	63.2	29.1	41.2	70.4	90.0	2,135
Total	4.3	4.7	9.1	25.1	36.3	61.4	29.5	41.1	70.6	87.2	30,931
Total ³	4.2	4.7	8.8	25.2	36.9	62.0	29.4	41.6	71.1	87.6	29,857

¹ Unmet need for spacing includes pregnant women whose pregnancy was mistimed; amenorrheic women who are not using family planning and whose last birth was mistimed, or whose last birth was unwanted but now say they want more children; and fecund women who are neither pregnant nor amenorrheic, who are not using any method of family planning, and say they want to wait 2 or more years for their next birth. Also included in unmet need for spacing are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted; amenorrheic women who are not using family planning, whose last child was unwanted and who do not want any more children; and fecund women who are neither pregnant nor amenorrheic, who are not using any method of family planning, and who want no more children

² Using for spacing is defined as women who are using some method of family planning and say they want to have another child or are undecided whether to have another. Using for limiting is defined as women who are using and who want no more children. Note that the specific methods used are not taken into account here

³ Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua).

F. Fertility Preferences

To gauge future fertility preferences, in the 2007 IDHS currently married respondents were asked whether or not they wanted another child and, if so, how soon. Overall, half of married women stated that they want no more children and three percent have been sterilized. Four in ten married women would like to have another child; 14 percent want another child within two years, 24 percent would prefer to wait two or more years, and three percent could not decide on the timing (Table 6). Thus, currently, eight in ten married women want either to space their next birth or to end childbearing. This represents the proportion of women who are potentially in need of some method of family planning.

Table 7 also shows that the desire to limit childbearing increases rapidly with the number of living children women have; 83 percent of women with no children want to have a child soon compared with 8 percent of women with two children. On the other hand, the proportion wanting no more children increases from 15 percent among women with one child to 60 percent among those with two children and 80 percent or higher among women with five or more children.

Table 7 Fertility preferences by number of living children

Percent distribution of currently married women age 15-49 by desire for children, according to number of living children, Indonesia 2007

Desire for children	Number of living children ¹							Total
	0	1	2	3	4	5	6+	
Have another soon ²	83.2	20.4	7.7	3.5	2.4	1.3	0.8	13.8
Have another later ³	6.3	54.8	21.0	9.6	6.0	3.6	1.9	24.1
Have another, undecided when	2.4	5.4	3.7	2.1	1.1	0.9	0.5	3.3
Undecided	1.1	3.2	5.0	4.5	3.7	3.4	5.0	4.0
Want no more	3.9	14.9	59.8	72.5	76.7	80.4	81.2	50.2
Sterilized ⁴	0.1	0.3	2.0	6.4	7.8	8.8	6.1	3.3
Declare infecund	3.0	0.9	0.8	1.1	1.8	1.3	4.0	1.2
Missing	0.0	0.1	0.1	0.2	0.5	0.2	0.5	0.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of women	1,859	8,239	9,566	5,789	2,823	1,315	1,340	30,931

¹ Includes current pregnancy
² Wants next birth within 2 years
³ Wants to delay next birth for 2 or more years
⁴ Includes both male and female sterilization

G. Maternity Care

Proper care during pregnancy and delivery are important for the health of both the mother and the baby in the 2007 IDHS, women who had given birth in the five years preceding the survey were asked a series of questions about maternal and child health care. For antenatal care (ANC), the interviewer was instructed to record all responses if more than one source of antenatal care was mentioned; however, in Table 8 only the provider with the highest qualifications is considered. For 93 percent of last births in the five years preceding the survey, antenatal care was received from a medically trained person (Table 8). Women age 20-34 are more likely than younger and older women to receive antenatal care from a health professional. Antenatal care from a medically trained person is higher among women with lower-order births, who may also be younger. Antenatal care coverage is higher in urban areas than in rural areas (98 percent and 90 percent, respectively). Antenatal care has a strong positive association with the woman's education, increasing from 63 percent among the small number of women with no education to 99 percent among women with more than secondary education.

Tetanus toxoid (TT) injections are given during pregnancy to prevent neonatal tetanus, a major cause of death among infants in many developing countries. In the 2007 IDHS, for the last birth since January 2002, the mothers were asked whether they had received tetanus toxoid injections while pregnant. Table 8 indicates that mothers received at least one TT injection during pregnancy for 73 percent of their most recent births. Mothers age 20-34 years, mothers of lower-order births, those living in urban areas, and better educated mothers are more likely to have received TT injections during pregnancy.

The third column in Table 8 shows the percentage of women who received iron supplementation when they were pregnant with their most recent birth in the five years preceding the survey. Overall, 77 percent of mothers reported receiving iron tablets. Again, women age 20-34, those with lower-order births, women living in urban areas, and those with better education are more likely to have received iron tablets.

Unlike ANC, which was collected only for the last live birth, information on delivery care was collected for all births in the five years preceding the survey. As with antenatal care, the interviewer was instructed to record all responses if more than one person assisted during delivery, with only the most

highly qualified person considered in Table 8. More than seven in ten births in the five years before the survey, the mothers were assisted by trained medical personnel during delivery. Differentials by background characteristics of the mother are similar to those for antenatal care.


The last column in Table 8 shows that 40 percent of births in the five years preceding the survey were delivered in a health facility. As with tetanus toxoid coverage, antenatal care, and delivery assistance, women 20-34, those with lower parity, urban women and better educated women are more likely than others to receive better medical services during delivery.

Table 8 Maternal care indicators							
Percentage of women age 15-49 who had a live birth in the five years preceding the survey who received antenatal care from a health professional for the most recent birth, and for the most recent birth received at least one tetanus toxoid injection during pregnancy and given iron tablets during pregnancy, and among all live births in the five years before the survey, percentage delivered by a health professional and percentage delivered in a health facility, according to background characteristics, Indonesia 2007							
Background characteristic	Percentage with antenatal care from health professional	Percentage given at least one tetanus toxoid injection during pregnancy	Percentage given iron tablets or syrup during pregnancy	Number of women	Percentage delivered by a health professional ¹	Percentage delivered in a health facility	Number of births
Mother's age at birth							
<20	90.8	66.4	74.6	1,385	62.6	35.8	1,716
20-34	94.2	74.7	78.9	10,552	74.8	47.4	12,482
35+	90.1	69.2	71.2	2,106	71.1	46.2	2,306
Mother's age at birth (5 year age groups)							
<20	90.8	66.4	74.6	1,385	62.6	35.8	1,716
20-24	93.9	74.8	78.8	3,796	72.1	43.9	4,573
25-29	94.5	73.6	79.4	3,731	76.0	49.8	4,340
30-34	94.3	75.8	78.4	3,026	76.7	49.1	3,569
35-39	93.0	72.2	74.6	1,572	74.3	48.7	1,745
40-44	82.5	61.0	62.7	466	64.3	39.6	488
45-49	75.1	55.9	51.9	67	40.3	31.7	73
Birth order							
1	95.2	75.1	80.6	4,856	79.6	54.4	5,855
2-3	94.4	75.2	80.0	6,568	73.4	45.6	7,529
4-5	90.4	68.2	68.8	1,860	65.4	34.5	2,207
6+	78.1	52.2	54.9	759	46.2	24.1	913
Residence							
Urban	97.7	77.5	84.0	5,897	87.6	70.3	6,835
Rural	90.1	69.8	72.5	8,145	62.7	28.9	9,669
Education							
No education	62.6	32.1	42.3	457	31.3	15.4	578
Some primary	82.2	55.4	60.6	1,677	44.7	22.0	1,996
Completed primary	92.4	71.1	73.3	4,106	62.3	31.0	4,759
Some secondary	96.4	79.1	81.9	3,543	79.3	48.7	4,132
Secondary +	99.1	81.2	87.9	4,260	94.0	71.2	5,038
Total	93.3	73.0	77.3	14,043	73.0	46.1	16,504
Total ²	93.7	73.3	77.6	13,643	73.7	46.6	16,056

¹ Doctor, obgyn, nurse, midwife, or village midwife
² Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua). If the respondent mentioned more than one person attending during delivery, only the most qualified is considered in this tabulation.

Figure 3 shows that the maternal care indicators related to care during pregnancy have not changed much since the 2002-03 IDHS. However, improvement is observed in delivery characteristics. The proportion of births delivered by a health professional increased from 66 percent to 73 percent, and the proportion of births delivered in a health facility increased from 40 percent to 46 percent.

**Figure 3 Trends in Maternal Health Care Indicators
Indonesia 2002-03 and 2007**



Note: Live births in the five years preceding the survey

H. Vaccination of Children

In the 2007 IDHS, mothers were asked to show the interviewer the health cards of all children born since January 2002. The interviewer then copied from the cards the dates of each vaccination received. If a child never received a health card or if the mother was unable to show the card to the interviewer, the mother was asked what vaccinations the child had received. Questions were asked for each vaccine type. In this report, a child was considered fully vaccinated if he or she had received a BCG vaccination against tuberculosis; three doses of DPT vaccine to prevent diphtheria, pertussis, and tetanus; three doses of polio vaccine (Polio 1-3); and one dose of measles vaccine. The results presented here are based on both health card information and, for those children without a card, information provided by the mother.

Table 9 pertains to vaccination coverage of children age 12 to 23 months, the age by which they should have received all vaccinations. Mothers were able to produce health cards for 37 percent of these children, which is an increase from 31 percent recorded in the 2002-03 IDHS. Based on both the health cards and the mothers' reports, 59 percent of children have received all of the recommended vaccinations. Again, this is a notable gain from 51 percent recorded in the 2002-03 IDHS. Nine percent of children had received no vaccinations and the remaining 33 percent of children were partially vaccinated.

Table 9 shows vaccination coverage by type of vaccine and dropout for DPT and polio vaccinations. Overall, 85 percent of children receive BCG, 84 percent receive the first dose of DPT vaccine, and 89 percent received the first dose of polio vaccine. However, only 67 percent of children

complete the DPT doses and only 56 percent receive the third dose of polio. Coverage of vaccination against measles is 76 percent.

Vaccination coverage differs slightly by gender of child, but varies substantially by other background characteristics. Lower-order births and children in urban areas are more likely than other children to have completed the vaccination schedule. For example, 68 percent of urban children compared with 52 percent of rural children had received all basic vaccinations. Full immunization coverage improves with mother's level of education, from 19 percent for children whose mothers have no education to 73 percent for children whose mothers have had more than secondary education.

Table 9. Vaccinations by background characteristics

Percentage of children age 12-23 months who received specific vaccines at any time before the survey (according to vaccination card or mother's report), and percentage with a vaccination card seen, by background characteristics, Indonesia 2007


Background characteristic	DPT			Polio				Measles	All basic vaccinations ¹	No vaccinations	Hepatitis B			Percentage with a vaccination card seen	Number of children	
	BCG	1	2	3	1	2	3				4	1	2			3
Sex																
Male	86.1	84.5	74.7	65.6	88.4	82.0	72.2	54.4	75.1	56.3	8.9	80.5	71.5	59.4	37.9	1,622
Female	84.7	84.3	76.8	67.9	89.2	83.2	74.8	56.7	77.9	61.0	8.2	80.4	72.0	61.2	35.6	1,472
Birth order																
1	89.1	89.0	80.0	72.7	90.5	85.0	77.4	60.2	78.8	63.2	6.0	84.0	75.1	65.8	41.9	1,139
2-3	85.4	84.8	76.8	66.3	89.8	84.2	74.4	55.5	76.8	58.6	8.6	82.3	73.3	59.6	36.8	1,382
4-5	79.2	76.5	68.7	59.8	84.4	76.6	67.6	49.7	72.6	53.3	12.6	71.8	64.9	55.6	30.4	416
6+	75.7	68.7	53.3	44.8	79.8	66.4	51.7	36.3	66.0	39.0	17.1	61.5	51.3	39.1	16.7	158
Residence																
Urban	92.0	91.3	83.6	74.8	94.4	89.5	82.2	64.0	82.0	67.5	4.7	88.8	81.0	70.0	38.1	1,274
Rural	80.8	79.5	70.2	61.0	84.9	77.7	67.3	49.5	72.5	52.3	11.3	74.6	65.2	53.5	35.9	1,820
Education																
No education	58.9	47.8	41.7	28.1	65.3	45.5	32.3	17.8	49.0	18.7	31.7	44.5	38.9	21.1	18.7	68
Some primary	71.1	61.9	52.7	43.5	76.4	68.5	57.5	36.2	63.5	37.3	20.3	60.6	47.5	36.6	23.8	352
Completed primary	80.3	81.3	70.1	59.9	85.7	77.2	65.4	47.1	71.0	52.1	11.3	75.5	64.1	51.5	38.3	898
Some secondary	87.6	87.6	77.9	69.2	90.8	85.0	75.8	62.3	76.0	60.6	6.1	82.4	75.3	65.1	40.8	754
Secondary +	95.0	95.0	89.1	81.2	95.9	92.8	87.1	67.1	87.8	72.8	2.5	92.6	86.3	75.3	38.2	1,022
Total	85.4	84.4	75.7	66.7	88.8	82.6	73.5	55.5	76.4	58.6	8.6	80.5	71.7	60.3	36.8	3,094
Total ²	85.9	85.0	76.3	67.4	89.4	83.5	74.4	56.1	76.8	59.2	8.0	81.1	72.5	61.1	37.0	3,014

¹ BCG, measles, and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

² Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua).

Figure 4 shows the trend in immunization coverage and health card coverage since the 2002-03 IDHS. The 2007 survey recorded substantial gains in coverage for all vaccines.

Figure 4 Trends in Vaccination Coverage among Children Age 12-23 Months, Indonesia 2002-03 and 2007



I. Childhood Illnesses

In the 2007 IDHS, several questions were included to obtain information on the prevalence and treatment practices for acute respiratory infection and diarrhea, illnesses that are known to contribute significantly to early childhood mortality. It should be noted that the morbidity data collected are subjective because they are based on a mother's perception of illness without validation by medical personnel.

The prevalence of ARI was estimated by asking mothers whether their children under age five had been ill with a cough accompanied by short, rapid breathing in the two weeks preceding the survey. In the 2007 IDHS, for each child under five years of age, mothers also were asked if the child had experienced an episode of diarrhea in the two weeks prior to the survey. Overall, 11 percent of children under age five were reported to have symptoms consistent with ARI during the two weeks prior to the survey, and 14 percent were reported to have had diarrhea.

Table 10 shows the treatment of children who were reported by their mothers to have symptoms of ARI and children with diarrhea in the two weeks preceding the survey. For 77 percent of children with ARI symptoms, advice or treatment was sought from a health provider. Children under six months were less likely than children in other age groups to receive treatment. Treatment of ARI varies little by the child's sex. Urban children are more likely to have received treatment for their ARI symptoms than rural children (84 percent and 73 percent, respectively). Treatment of children with ARI increases with their mother's education; 42 percent for children whose mother has no formal education were likely to have been taken to a provider compared with 87 percent for children whose mothers have more than secondary education.

Treatment or advice was sought for 58 percent of children with diarrhea, and 35 percent were given solution prepared from packets of oral rehydration salts (ORS). Mother's education plays a role in whether advice or treatment is sought for the diarrhea. Better educated mothers are more likely than less educated mothers to seek advice for their children.

Table 10 also shows the percentage of children with diarrhea who were given oral rehydration therapy, which includes solution prepared from ORS, recommended homemade fluids and increased fluids. Overall, 61 percent of children with diarrhea were given ORT. ORT is less likely to be given to young children (under 6 months old). Otherwise, the proportions receiving ORT do not vary markedly by other characteristics

Table 10 Treatment for acute respiratory infection, fever, and diarrhea

Among children under five years who were sick with a cough accompanied by short, rapid breathing (symptoms of acute respiratory infection - ARI) or fever in the two weeks preceding the survey, percentage for whom treatment was sought from a health facility or provider, and among children under five years who were sick with diarrhea during the two weeks preceding the survey, percentage for whom treatment was sought from a health facility or provider, percentage given a solution made from oral rehydration salt (ORS) packets, and percentage given any oral rehydration therapy (ORT), by background characteristics, Indonesia 2007

Background characteristic	Children with symptoms of ARI or with fever		Children with diarrhea			
	Percentage for whom treatment was sought from a health facility/provider ¹	Number with ARI/fever	Percentage for whom treatment was sought from a health facility/provider ¹	Percentage given solution from ORS packet	Percentage given any ORT ²	Number with diarrhea
Age in months						
<6	65.5	402	35.9	6.6	33.4	187
6-11	80.1	739	61.1	28.0	51.7	302
12-23	81.1	1,325	64.2	40.2	67.9	640
24-35	76.6	1,181	60.1	37.7	65.1	482
36-47	76.6	991	50.9	35.1	59.7	306
48-59	73.8	900	58.1	42.7	68.0	261
Sex						
Male	76.4	2,904	58.2	35.4	63.7	1,217
Female	77.4	2,634	57.4	33.7	57.5	963
Residence						
Urban	83.6	2,166	63.9	33.4	58.7	799
Rural	72.5	3,372	54.4	35.4	62.2	1,381
Education						
No education	42.4	216	34.4	23.3	49.8	97
Some primary	66.5	713	49.7	31.6	60.9	318
Completed primary	72.6	1,683	50.5	32.5	59.6	683
Some secondary	81.5	1,479	63.9	40.3	64.8	558
Secondary +	87.4	1,446	70.4	35.3	60.7	522
Total	76.9	5,539	57.9	34.7	60.9	2,180
Total ³	76.5	5,382	57.2	34.6	60.4	2,111

¹ Excludes pharmacy, shop, traditional practitioner
² Includes ORS, recommended home fluids, or increased fluids
³ Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua)

J. Infant Feeding Practices

Breast milk contains all the nutrients needed by children in the first 6 months of life. Supplementing breast milk before the child is 6 months of age is discouraged because it increases the likelihood of contamination and hence, risks of diarrheal disease. At a later stage of the baby's development, breast milk should be supplemented by other liquids and eventually by solid or mushy food to provide adequate nourishment.

The 2007 IDHS collected data on infant feeding for all children born in the three years preceding the survey and living with their mothers. Table 11 shows that 32 percent of infants under age 6 months are being exclusively breastfed. For the rest, breast milk is supplemented with other milk (18 percent), plain water (9 percent), non-milk liquids or juices (2 percent), or with solid or semi-solid food (30 percent). By age 6-9 months, 14 percent of infants are no longer being breastfed, and 75 percent receive complementary food.

Feeding of babies using a bottle with a nipple is not recommended at any age. However, the 2007 IDHS findings show that this practice has increased substantially since the 2002-03 IDHS. Overall, in 2007, 25 percent of babies under two months of age are being given this type of feeding compared with 14 percent in 2002-03. The proportion of babies who use a bottle with a nipple for children under six months in 2007 also is higher than that in 2002-03 (28 percent and 17 percent, respectively).

Table 11 Breastfeeding status by age

Percent distribution of youngest children under three years living with the mother by breastfeeding status, and percentage of children under three years using a bottle with a nipple, according to age in months, Indonesia 2007

Age in months	Not breast-feeding	Exclusively breastfed	Breastfeeding and consuming:				Total	Number of children	Percentage using a bottle with a nipple ¹	Number of children
			Plain water only	Non-milk liquids/ juice	Other milk	Comple-mentary food				
<2	4.6	48.3	5.8	0.6	28.6	12.2	100.0	479	25.1	486
2-3	10.5	34.4	9.6	1.7	16.5	27.2	100.0	590	30.2	599
4-5	9.7	17.8	10.6	2.6	11.2	48.1	100.0	595	27.8	601
6-7	13.7	7.2	5.1	0.6	3.3	70.0	100.0	596	24.5	607
8-9	13.6	1.7	2.4	0.5	1.7	80.1	100.0	592	29.2	608
10-11	17.6	0.7	1.4	1.3	0.9	78.1	100.0	496	28.1	504
12-15	20.1	0.6	2.2	0.4	0.1	76.5	100.0	1,090	33.4	1,119
16-19	27.6	0.2	1.3	0.1	0.5	70.4	100.0	924	34.7	980
20-23	49.7	0.0	0.0	0.5	0.0	49.7	100.0	915	37.3	995
24-27	61.2	0.3	0.2	0.7	0.0	37.5	100.0	986	34.0	1,107
28-31	72.1	0.0	0.0	0.5	0.1	27.3	100.0	906	33.6	1,038
32-35	77.4	0.0	0.0	0.0	0.0	22.6	100.0	856	29.3	1,017
<6	8.5	32.4	8.9	1.7	18.1	30.4	100.0	1,664	27.9	1,686
6-9	13.7	4.5	3.8	0.6	2.5	75.0	100.0	1,188	26.9	1,215
<6 ^a	8.5	32.8	9.2	1.6	18.2	29.7	100.0	1,617	27.9	1,638
6-9 ^a	13.8	4.5	3.9	0.5	2.6	74.7	100.0	1,162	27.1	1,189

Note: Breastfeeding status refers to a 24-hour period (yesterday and the past night). Children who are classified as *breastfeeding and consuming plain water only* consume no liquid or solid supplements. The categories of not breastfeeding, exclusively breastfed, breastfeeding and consuming plain water, water-based liquids/juice, other milk, and complementary foods (solids and semi-solids) are hierarchical and mutually exclusive, and their percentages add to 100 percent. Thus, children who receive breast milk and non-milk liquids and who do not receive complementary foods are classified in the non-milk liquid category even though they may also get plain water. Any children who get complementary food are classified in that category as long as they are breastfeeding as well.

¹ Based on all children under three years

^a Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua).

K. Childhood Mortality

An important objective of the 2007 IDHS was to measure levels and trends in mortality among children. The childhood mortality rates presented in Table 12 are estimated directly from information obtained in the birth history section of the Women's Questionnaire on each child's birth date, survivorship status, and the age at death for children who died. The rates are defined as followed:

- Neonatal mortality: the probability of death in the first month

- Postneonatal mortality: the difference between infant mortality and neonatal mortality
- Infant mortality: the probability of death before the first birthday
- Child mortality: the probability of death between the first and fifth birthdays
- Under-five mortality: the probability of death before the fifth birthday.


The rates shown in Table 12 were calculated for three five-year periods before the survey. For the most recent five-year period, the under-five mortality rate was 45 deaths per 1,000 live births. Looking at the age pattern of mortality during the five-year period prior to the survey, more than three-fourths of all deaths took place during the first year of the child's life. In turn, the majority of infant deaths occurred during neonatal period.

Years preceding the survey	Approximate calendar year	Neonatal mortality (NN)	Postneonatal mortality (PNN) ¹	Infant mortality (_{1q0})	Child mortality (_{4q1})	Under-five mortality (_{5q0})
0-4	2003-2007	20	15	34	11	45
5-9	1998-2002	24	21	44	15	59
10-14	1988-1997	28	25	53	17	69
0-4 ^a	2003-2007	19	15	34	10	44
5-9 ^a	1998-2002	24	21	45	14	58
10-14 ^a	1988-1997	28	25	53	17	69

¹ Computed as the difference between the infant and neonatal mortality rates
^a Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua)

Figure 5 shows the trends in infant and under-five mortality since the 1991 IDHS. The figure shows that the decline in childhood mortality has slowed down in recent years, as is the case in populations with low mortality rates.

**Figure 5 Trends in Early Childhood Mortality Rates
Indonesia 1991-2007**



L. Knowledge of HIV/AIDS

Since 1994, the Government of Indonesia has been promoting a National AIDS Strategy, a collaborative effort by the government, nongovernmental organizations, the private sector, and the community. The strategy promotes healthy life, safe sex, safe injection, condom use and supporting people living with HIV/AIDS.

In the 2007 IDHS, female and male respondents were asked whether they had heard of AIDS, and if so, what were their perceptions regarding prevention and treatment of the disease. Table 12 shows that 61 percent of ever-married women and 71 percent of currently married men reported having heard of AIDS. These rates are practically the same as those recorded in the 2002-03 IDHS (59 percent for women and 73 percent for men, respectively).

Data in Table 13 show that knowledge of AIDS among women varies according to background characteristics. Women age 20-39, married women, women who live in urban areas, and better educated women are more likely than other women to have heard of AIDS. The knowledge level among married men follows the pattern OF WOMEN, with levels markedly higher among urban and better educated men.

Table 13 Knowledge of AIDS				
Percentage of ever-married women and currently married men who have heard of AIDS by background characteristics, Indonesia 2007				
Background characteristic	Women		Men	
	Have heard of AIDS	Number	Have heard of AIDS	Number
Age				
15-19	52.4	845	45.9	29
20-24	68.8	4,094	68.7	432
25-29	71.8	5,771	77.4	1,116
30-39	65.4	12,024	79.4	3,097
40-49	47.3	10,160	68.3	2,930
50-54	na	0	53.7	1,155
Marital status				
Married	61.8	30,931	71.4	8,758
Divorced/widowed	49.3	1,964	na	0
Residence				
Urban	77.3	13,745	85.5	3,728
Rural	49.3	19,150	61.0	5,030
Education				
No education	9.4	2,270	17.7	361
Some primary	27.8	5,572	39.6	1,605
Completed primary	52.8	10,077	63.7	2,339
Some secondary	77.6	6,781	84.6	1,721
Secondary +	94.3	8,193	95.5	2,727
Total	61.0	32,895	71.4	8,758
Total ¹	61.4	31,850	71.7	8,472

na = Not applicable
¹ Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua).

Knowledge of Ways to Reduce the Risk of Getting AIDS Virus

Although many women and men in Indonesia have a basic knowledge of AIDS, the proportions aware of ways in which the risk of infection can be reduced are generally low. Overall, 42 percent of women see limiting sex to one uninfected partner as a means of reducing the risk of transmission, 37 percent agree that abstaining from having sex would reduce the chances of infection, and 35 percent say that consistent use of condoms would reduce the chances of infection. Knowledge among men is slightly higher than that for women. The corresponding proportions are 52 percent for limiting sex to one uninfected partner, 49 percent for using condoms, and 43 percent for abstaining from having sex.

Results in Table 14 show that knowledge of AIDS prevention among women and men varies according to their background characteristics. Women age 20-39, married women, women who live in the urban areas, and better-educated women are more likely than other women to know ways to reduce the risk of getting the AIDS virus. The knowledge level among married men follows the women's pattern, high among urban and better-educated men.

Table 14 Knowledge of ways to reduce the risk of getting the AIDS virus

Percentage of ever-married women age 15-49 and currently married men age 15-54 who, in response to prompted questions, say that people can reduce the risk of getting the AIDS virus by using condoms every time they have sexual intercourse, by having one uninfected sex partner who has no other partners, and by abstaining from sexual intercourse, by background characteristics, Indonesia 2007

Background characteristic	Percentage of women who say HIV can be prevented by:					Number of women	Percentage of men who say HIV can be prevented by:				
	Using condoms ¹	Limiting sexual intercourse to one uninfected partner ²	Using condoms and limiting sexual intercourse to one uninfected partner ^{1,2}	Abstaining from sexual intercourse	Using condoms ¹		Limiting sexual intercourse to one uninfected partner ²	Using condoms and limiting sexual intercourse to one uninfected partner ^{1,2}	Abstaining from sexual intercourse	Number of men	
Age											
15-19	27.5	32.4	21.0	27.2	845	*	*	*	*	29	
20-24	40.3	47.2	33.4	42.6	4,094	49.8	49.7	42.8	39.6	432	
25-29	40.8	49.2	34.0	42.4	5,771	52.6	56.7	45.4	44.4	1,116	
30-39	39.3	46.0	33.6	40.3	12,024	56.0	59.3	47.5	49.1	3,097	
40-49	26.6	32.4	22.5	27.3	10,160	46.5	49.5	38.2	40.9	2,930	
50-54	na	na	na	na	0	32.3	38.0	28.2	30.9	1,155	
Marital status											
Married	35.9	42.8	30.3	37.1	30,931	48.9	52.4	41.3	42.9	8,758	
Divorced/widowed	28.4	32.7	23.8	29.3	1,964	na	na			0	
Residence											
Urban	48.9	56.4	42.4	49.3	13,745	62.2	66.6	53.7	52.7	3,728	
Rural	25.8	32.0	20.9	27.5	19,150	39.1	41.8	32.2	35.6	5,030	
Education											
No education	4.3	5.2	3.1	4.7	2,270	3.2	4.5	2.6	3.6	361	
Some primary	11.2	15.0	8.7	12.3	5,572	21.6	22.9	16.8	18.5	1,605	
Completed primary	26.0	32.2	20.6	27.7	10,077	36.8	41.1	29.0	35.0	2,339	
Some secondary	44.9	53.8	37.5	47.3	6,781	59.6	62.1	49.3	51.4	1,721	
Secondary +	64.5	73.6	56.8	64.0	8,193	74.9	79.5	66.5	63.7	2,727	
Total	35.5	42.2	29.9	36.6	32,895	48.9	52.4	41.3	42.9	8,758	
Total ³	35.9	42.8	30.3	37.1	30,931	32.3	38.0	28.2	30.9	1,155	

Note: An asterisk indicates that an estimate is based on fewer than 25 unweighted cases.

na = Not applicable.

¹ Using condoms every time they have sexual intercourse

² Partner who has no other partners

³ Excludes 5 provinces which were not included in the 2002-03 IDHS (NAD, Maluku, North Maluku, Papua, and West Papua).

Knowledge of Voluntary Counseling and Testing (VCT)

In the 2007 IDHS, female and male respondents who had heard of AIDS were asked also whether they were aware of the existence of voluntary HIV testing preceded with counseling. Table 15.1 shows that only 8 percent of ever-married women reported having heard of VCT. Data in Table 15.1 show that knowledge of VCT among women varies according to the woman's background characteristics. Knowledge of VCT among women age 15-19 is very limited (3 percent). Women age 20-39, women who live in the urban areas, and women who completed secondary education are more likely than other women to have heard of VCT. Knowledge of VCT decreases with increasing number of children; women with no children are the most likely than women with children to have heard of VCT.

Table 15.1 Knowledge about voluntary counseling and testing (VCT): Women

Percentage of ever-married women age 15-49 who know about voluntary HIV counseling and testing (VCT) and percentage who know specific VCT providers, by background characteristics, Indonesia 2007

Background characteristic	Ever-married women		Among ever-married women who know about VCT, percentage who know specific providers												Number of ever-married women who know about VCTI
	Percentage who know about VCT	Number of ever-married women	Public						Private						
			Public hospital	Public health center	Public clinic	Public VCT clinic	Other public	Private hospital	Private clinic	Private VCT clinic	Private doctor	Private nurse/midwife	Other private medical	Other	
Age															
15-19	2.9	845	(53.2)	(35.2)	(0.0)	(6.9)	(0.0)	(9.4)	(0.0)	(0.0)	(6.8)	(6.6)	(0.0)	(2.9)	24
20-24	7.1	4,094	55.2	8.5	1.0	6.2	0.7	14.3	2.0	3.4	12.4	8.6	3.2	7.8	292
25-29	8.9	5,771	65.0	13.2	3.5	10.8	0.4	13.9	3.1	4.0	7.2	3.1	3.7	6.3	514
30-34	8.4	6,020	70.9	13.6	0.4	7.6	1.0	13.4	0.5	3.3	7.7	2.0	2.7	5.2	505
35-39	9.1	6,004	75.8	16.8	0.9	6.3	1.7	18.7	0.9	3.1	4.3	2.3	1.5	5.1	545
40-44	6.1	5,365	79.0	12.7	0.2	8.7	2.1	16.0	0.3	4.5	7.5	0.8	0.4	4.6	325
45-49	6.1	4,795	71.2	14.5	2.7	10.1	1.9	11.2	0.3	4.2	6.4	0.4	2.0	4.5	291
Residence															
Urban	12.4	13,745	72.2	12.0	1.7	8.8	1.2	15.8	1.0	3.8	5.5	1.6	2.2	5.0	1,705
Rural	4.1	19,150	64.9	17.7	1.0	7.0	1.2	12.6	1.8	3.3	10.8	5.1	2.5	6.7	792
Education															
No education	0.3	2,270	*	*	*	*	*	*	*	*	*	*	*	*	7
Some primary	1.3	5,572	74.1	17.2	0.0	6.2	0.0	11.9	0.0	1.1	6.1	6.7	0.0	8.8	72
Completed primary	2.6	10,077	72.5	22.1	0.4	1.3	0.7	9.6	0.1	0.7	8.6	4.7	0.1	6.1	264
Some secondary	6.7	6,781	66.3	16.6	0.6	3.8	0.1	9.1	1.9	3.8	7.5	3.0	2.4	6.4	453
Secondary +	20.8	8,193	70.1	11.6	1.9	10.6	1.6	17.3	1.3	4.1	6.9	2.2	2.7	5.1	1,700
Living children															
0	10.6	2,687	64.3	11.3	0.9	12.6	0.5	10.6	0.7	5.4	11.6	3.0	2.7	7.0	285
1-2	8.1	18,545	69.2	13.4	1.7	8.0	0.8	16.1	1.8	3.6	7.7	3.0	2.6	5.6	1,511
3-4	6.8	8,908	72.9	16.5	0.4	6.5	2.6	14.5	0.3	3.4	3.9	2.2	1.3	4.9	605
5+	3.5	2,754	76.9	9.4	5.7	9.9	1.5	8.6	0.0	0.4	7.1	0.8	3.1	3.8	97
Total	7.6	32,895	69.8	13.8	1.5	8.2	1.2	14.8	1.2	3.7	7.2	2.7	2.3	5.5	2,497

Note: Figures in parentheses are based on 25-49 unweighted cases. An asterisk indicates that an estimate is based on fewer than 25 unweighted cases.

Meanwhile, only 7 percent of married men reported having heard of VCT. Data in Table 15.2 show that knowledge of VCT among men also varies according to the man's background characteristics. Knowledge of VCT among oldest men is lower than younger men (4.6 percent). Men age 30-349, men who live in the urban areas, and men who completed secondary education are more likely than other women to have heard of VCT. Knowledge of VCT decreases with increasing number of children; men with no children are the most likely than men with children to have heard of VCT.

Percentage of ever-married women and married men who know specific VCT providers from public hospital is quite high, more than 60 percent.

Table 15.2 Knowledge about voluntary counseling and testing (VCT): Men

Percentage of currently married men age 15-54 who know about voluntary HIV counseling and testing (VCT) and percentage who know specific VCT providers, by background characteristics, Indonesia 2007

Background characteristic	Currently married men		Among currently married men who know about VCT, percentage who know specific providers											Number of currently married men who know about VCT		
	Percent-age who know about VCT	Number of ever-married men	Public					Private					Other private medical		Other	
			Public hospital	Public health center	Public clinic	Public VCT clinic	Other public	Private hospital	Private clinic	Private VCT clinic	Private doctor	Private nurse/midwife				
Age																
15-19	(1.5)	29	*	*	*	*	*	*	*	*	*	*	*	*	*	0
20-24	7.2	432	(55.9)	(3.7)	(2.9)	(14.6)	(5.6)	(7.7)	(0.0)	(1.1)	(1.8)	(0.0)	(14.2)	(22.9)	31	
25-29	7.3	1,116	57.4	5.2	2.2	14.7	3.9	13.8	2.0	5.4	1.3	0.0	8.8	12.6	82	
30-34	9.9	1,418	55.2	18.6	0.9	7.4	9.6	15.4	0.9	7.2	5.3	0.7	11.3	21.2	140	
35-39	6.6	1,679	63.2	20.1	0.0	11.6	4.3	7.7	0.2	5.5	1.7	1.3	8.2	17.4	110	
40-44	6.1	1,570	56.6	14.5	1.0	5.4	8.6	9.3	0.0	1.1	5.3	0.0	12.5	8.9	95	
45-49	7.6	1,359	75.6	10.4	6.3	6.9	0.7	11.8	1.6	2.8	17.5	4.1	9.3	10.9	103	
50-54	4.6	1,155	76.1	14.2	0.0	6.5	2.6	4.8	0.0	1.6	2.2	0.0	2.5	9.5	53	
Residence																
Urban	10.6	3,728	63.5	13.8	0.8	11.3	6.3	11.5	1.0	3.8	5.4	1.2	7.5	15.3	397	
Rural	4.3	5,030	60.3	14.2	3.7	4.8	3.8	9.9	0.3	4.8	6.4	0.9	13.5	14.1	217	
Education																
No education	0.1	361	*	*	*	*	*	*	*	*	*	*	*	*	0	
Some primary	0.5	1,605	*	*	*	*	*	*	*	*	*	*	*	*	9	
Completed primary	2.0	2,339	(67.9)	(16.5)	(0.9)	(3.7)	(3.1)	(3.5)	(2.3)	(12.4)	(10.3)	(0.4)	(12.0)	(3.5)	47	
Some secondary	5.5	1,721	49.8	7.7	0.5	10.2	5.8	10.4	0.9	0.7	3.2	0.5	16.1	19.0	95	
Secondary +	16.8	2,727	63.8	14.4	2.3	9.5	5.6	11.7	0.6	4.1	6.0	1.2	8.3	15.5	459	
Living children																
0	7.5	723	53.1	14.1	4.5	2.1	4.9	13.0	2.2	0.3	0.0	0.0	21.9	15.2	54	
1-2	7.4	4,855	60.7	10.9	2.0	10.3	5.9	12.0	0.6	5.1	6.1	1.4	8.6	17.3	362	
3-4	7.4	2,411	66.3	18.7	0.3	8.7	4.8	8.6	0.1	3.5	6.9	0.8	8.5	10.8	177	
5+	2.8	769	(80.4)	(24.1)	(5.4)	(7.5)	(3.5)	(7.1)	(5.2)	(2.8)	(4.1)	(0.0)	(5.7)	(7.0)	21	
Total	7.0	8,758	62.4	13.9	1.8	9.0	5.4	10.9	0.8	4.2	5.7	1.1	9.6	14.9	614	

Note: Figures in parentheses are based on 25-49 unweighted cases. An asterisk indicates that an estimate is based on fewer than 25 unweighted cases.

REFERENCES

Badan Pusat Statistik-Statistics Indonesia (BPS) and ORC Macro. 2003. *Indonesia Demographic and Health Survey 2002-2003*. Calverton, Maryland, USA: BPS and ORC Macro.

APPENDIX

Table A.1 Background characteristics of respondents by province

Percent distribution of women and men by province, weighted and unweighted numbers, Indonesia 2007

Province	Women			Men		
	Weighted percent	Weighted number	Unweighted number	Weighted percent	Weighted number	Unweighted number
Sumatera						
DI Aceh	1.6	514	929	1.6	137	245
North Sumatera	4.5	1,487	1,126	4.2	370	277
West Sumatera	1.7	570	905	1.6	137	217
Riau	1.5	494	991	1.5	130	243
Jambi	1.1	367	874	1.1	95	231
South Sumatera	2.8	928	1,055	2.8	241	289
Bengkulu	0.6	211	753	0.6	53	197
Lampung	2.9	963	920	3.1	271	265
Bangka Belitung	0.6	194	815	0.6	52	222
Riau Islands	0.4	140	731	0.4	36	184
Java						
DKI Jakarta	4.5	1,471	1,722	4.7	408	497
West Java	16.9	5,545	1,693	16.5	1,444	432
Central Java	16.4	5,383	1,450	17.3	1,517	425
DI Yogyakarta	1.7	551	1,110	1.7	146	305
East Java	18.0	5,924	1,485	17.8	1,561	387
Banten	4.0	1,310	1,413	3.9	344	357
Bali and Nusa Tenggara						
Bali	1.8	587	1,302	2.0	174	409
West Nusa Tenggara	2.1	705	964	2.2	194	272
East Nusa Tenggara	1.9	627	821	2.0	172	236
Kalimantan						
West Kalimantan	1.9	628	933	1.8	162	236
Central Kalimantan	0.9	294	792	0.9	82	223
South Kalimantan	1.7	550	953	1.5	128	237
East Kalimantan	1.4	475	837	1.5	132	218
Sulawesi						
North Sulawesi	1.1	373	894	1.2	102	229
Central Sulawesi	1.0	339	818	1.0	89	210
South Sulawesi	3.2	1,067	1,217	3.0	259	280
Southeast Sulawesi	0.8	259	767	0.7	60	172
Gorontalo	0.5	163	884	0.5	46	224
West Sulawesi	0.4	139	757	0.5	41	226
Maluku and Papua						
Maluku	0.5	168	805	0.5	44	222
North Maluku	0.4	129	754	0.4	36	194
West Papua	0.3	89	702	0.3	24	188
Papua	0.8	251	723	0.8	70	209
Total	100.0	32,895	32,895	100.0	8,758	8,758

Table A.2 Fertility by province

Total fertility rate for the three years preceding the survey, percentage of women age 15-49 currently pregnant, and mean number of children ever born to women age 40-49 years, by province, Indonesia 2007

Province	Total fertility rate	Percentage of women age 15-49 currently pregnant	Mean number of children ever born to women age 40-49
Sumatera			
DI Aceh	3.1	6.1	4.2
North Sumatera	3.8	3.6	4.7
West Sumatera	3.4	3.8	4.3
Riau	2.7	4.6	4.0
Jambi	2.8	5.3	3.6
South Sumatera	2.7	3.1	4.2
Bengkulu	2.4	3.9	4.0
Lampung	2.5	4.0	4.3
Bangka Belitung	2.5	5.5	4.2
Riau Islands	3.1	4.7	3.3
Java			
DKI Jakarta	2.1	3.8	3.2
West Java	2.6	4.1	3.7
Central Java	2.3	3.5	3.3
DI Yogyakarta	1.8	4.4	2.5
East Java	2.1	2.6	2.8
Banten	2.6	3.4	4.6
Bali and Nusa Tenggara			
Bali	2.1	3.5	2.5
West Nusa Tenggara	2.8	4.9	4.1
East Nusa Tenggara	4.2	6.2	4.2
Kalimantan			
West Kalimantan	2.8	5.1	4.0
Central Kalimantan	3.0	7.1	3.6
South Kalimantan	2.6	5.7	3.7
East Kalimantan	2.7	5.7	3.7
Sulawesi			
North Sulawesi	2.8	4.1	2.8
Central Sulawesi	3.3	4.0	3.9
South Sulawesi	2.8	4.1	3.7
Southeast Sulawesi	3.3	5.6	4.3
Gorontalo	2.6	3.8	3.2
West Sulawesi	3.5	6.3	4.4
Maluku and Papua			
Maluku	3.9	5.1	4.4
North Maluku	3.2	6.5	4.3
West Papua	3.4	4.7	4.3
Papua	2.9	4.2	3.8
Total	2.6	3.9	3.5

Table A.3 Knowledge of contraceptive methods by province

Percentage of currently married women age 15-49 who know any contraceptive method, by province, Indonesia 2007

Province	Any method	Any modern method	Modern method									Traditional method				Number of women	
			Female sterilization	Male sterilization	Pill	IUD	Injectables	Im-plants	Male condom	Intravag/Dia-phragm	Emergency contra-ception	LAM	Any tra-ditional method	Periodic absti-nence	With-drawal		Folk method
Sumatera																	
DI Aceh	95.8	95.5	28.3	20.5	90.5	61.7	92.9	59.5	62.0	13.2	9.2	18.0	31.6	21.3	20.0	7.7	472
North Sumatera	95.5	95.4	65.3	33.6	90.1	81.0	93.8	83.1	79.6	19.7	18.9	37.5	73.4	58.5	65.3	6.8	1,389
West Sumatera	99.3	99.3	71.4	22.0	96.5	93.9	98.3	95.0	88.0	13.9	8.7	27.2	62.1	42.3	48.3	6.0	532
Riau	97.2	96.5	51.6	29.1	93.5	79.8	93.5	82.0	82.5	12.1	7.1	28.0	60.8	44.1	48.1	3.3	474
Jambi	99.2	99.1	48.0	33.0	97.4	83.1	98.0	88.2	75.5	10.4	3.6	11.9	40.4	23.6	15.4	18.9	346
South Sumatera	99.0	99.0	77.9	34.5	97.6	82.8	98.5	94.2	79.9	12.1	3.0	14.7	40.3	34.1	23.6	2.9	871
Bengkulu	99.6	99.6	82.9	47.3	99.0	96.0	99.2	96.8	88.2	15.2	7.4	19.7	41.8	31.9	23.7	8.0	200
Lampung	99.6	99.6	72.5	50.6	99.2	93.1	99.4	95.2	87.2	16.6	6.9	19.6	60.9	49.1	42.9	5.2	925
Bangka Belitung	99.2	99.2	55.3	35.9	97.3	79.0	96.6	81.6	79.7	11.0	4.6	14.5	41.9	33.6	23.7	4.1	182
Riau Islands	98.8	98.8	54.9	37.8	96.4	87.2	96.4	84.4	89.6	15.8	11.1	38.8	59.8	52.5	38.0	4.1	134
Java																	
DKI Jakarta	99.9	99.9	86.3	74.4	99.8	97.9	99.7	96.3	97.8	30.4	13.0	47.0	80.2	77.7	54.8	3.7	1,352
West Java	99.9	99.9	62.3	52.1	99.3	88.5	99.2	87.2	78.8	12.6	4.1	12.2	36.8	30.9	19.7	7.3	5,243
Central Java	99.4	99.4	77.0	49.5	94.5	86.8	97.4	90.1	79.8	12.3	4.4	27.6	46.8	38.8	31.9	2.6	5,158
DI Yogyakarta	100.0	100.0	90.2	69.8	99.7	98.7	99.9	96.9	97.2	33.3	9.2	32.4	83.6	74.0	71.5	7.2	517
East Java	98.5	98.4	72.3	22.3	94.3	81.7	95.3	86.1	71.6	9.5	5.4	25.2	44.4	32.1	33.2	6.8	5,525
Banten	99.4	99.3	64.7	43.0	96.7	83.2	98.5	90.7	78.3	11.6	5.7	21.6	39.2	33.7	28.0	4.6	1,231
Bali and Nusa Tenggara																	
Bali	99.5	99.5	81.1	59.0	95.9	96.7	97.8	70.7	81.7	30.1	11.3	32.5	69.0	62.2	56.8	2.2	564
West Nusa Tenggara	98.1	97.8	57.8	32.6	90.2	84.8	96.1	87.0	61.7	9.7	5.2	13.9	28.8	23.3	15.9	5.1	636
East Nusa Tenggara	89.9	89.8	46.9	28.3	79.5	68.3	87.4	70.8	47.1	6.9	4.3	14.0	47.6	38.5	31.8	4.8	577
Kalimantan																	
West Kalimantan	97.1	97.0	33.8	20.7	94.0	67.5	93.7	66.9	65.9	7.6	3.8	8.7	23.3	17.3	13.6	3.0	590
Central Kalimantan	100.0	99.8	41.7	28.6	98.4	74.8	97.5	79.8	76.1	16.9	14.5	24.2	52.6	43.1	31.2	8.9	280
South Kalimantan	99.3	99.2	53.7	27.6	98.9	80.9	97.5	88.0	80.2	12.7	2.0	16.6	50.1	29.0	31.8	17.8	507
East Kalimantan	99.2	99.1	68.2	44.3	97.1	88.9	98.4	91.6	89.1	18.7	7.7	43.0	66.1	57.7	47.7	10.3	455
Sulawesi																	
North Sulawesi	99.7	99.7	72.3	41.4	98.5	92.2	98.4	94.6	85.1	21.6	10.0	29.9	60.5	56.5	31.0	2.2	360
Central Sulawesi	99.7	99.7	53.6	20.1	97.2	84.6	98.3	91.3	71.3	14.9	16.1	25.4	48.4	37.8	31.5	8.8	319
South Sulawesi	97.1	96.4	39.6	17.0	91.9	61.1	93.2	76.2	60.1	9.5	3.7	17.4	51.3	27.4	41.4	4.6	967
Southeast Sulawesi	97.5	97.2	43.8	18.5	91.8	67.9	93.6	81.8	64.4	14.5	8.6	23.1	46.8	38.2	27.7	5.1	242
Gorontalo	99.7	99.7	47.6	22.8	98.3	94.4	97.8	95.5	60.1	17.6	2.1	14.3	38.9	36.0	19.7	1.6	152
West Sulawesi	97.3	97.3	36.4	11.8	93.3	63.9	95.1	77.9	58.1	8.8	6.1	23.4	42.0	29.0	28.0	3.4	131
Maluku and Papua																	
Maluku	93.1	91.2	41.8	26.8	84.5	70.5	89.9	77.5	59.8	10.5	5.5	13.5	41.5	32.7	27.0	5.0	157
North Maluku	97.2	96.5	40.6	18.1	89.7	71.8	94.4	84.6	60.4	12.4	7.7	21.1	39.3	31.0	25.4	9.3	120
West Papua	93.2	92.7	49.0	34.7	88.9	74.5	88.8	72.2	77.9	20.1	7.9	37.0	43.6	37.0	26.1	4.9	83
Papua	84.9	61.9	31.5	15.7	51.6	37.8	54.8	40.4	44.1	6.4	4.7	14.6	65.7	22.2	19.2	41.6	242
Total	98.6	98.3	66.1	39.4	95.1	83.9	96.5	86.4	76.8	13.7	6.4	23.3	48.4	38.5	33.3	6.0	30,931

LAM = Lactational amenorrhea method

Table A.4 Current use of contraception by province

Percent distribution of currently married women by contraceptive method currently used, according to province, Indonesia 2007

Province	Modern method										Traditional method				Total	Number of women	
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	IUD	Injectables	Implants	Male condom	LAM	Any traditional method	Periodic abstinence	Withdrawal	Folk method			Not currently using
Sumatera																	
DI Aceh	47.4	45.4	0.6	0.0	9.3	1.3	33.2	0.4	0.5	0.1	2.0	0.6	0.8	0.6	52.6	100.0	472
North Sumatera	54.2	42.6	7.4	0.0	11.7	2.1	17.4	1.9	2.1	0.1	11.5	2.8	7.9	0.9	45.8	100.0	1,389
West Sumatera	59.9	52.8	2.6	0.0	8.7	5.8	29.9	3.9	1.9	0.0	7.2	1.8	5.1	0.2	40.1	100.0	532
Riau	56.7	52.8	2.5	0.0	14.7	1.6	31.3	1.7	1.0	0.0	4.0	1.6	2.2	0.2	43.3	100.0	474
Jambi	65.2	62.5	2.4	0.0	18.4	1.5	34.3	4.7	1.2	0.0	2.7	2.0	0.4	0.3	34.8	100.0	346
South Sumatera	64.8	62.6	2.3	0.0	10.1	0.9	44.1	4.8	0.5	0.0	2.1	1.3	0.7	0.2	35.2	100.0	871
Bengkulu	74.0	70.4	1.5	0.1	13.0	1.7	46.9	5.4	1.8	0.0	3.6	1.2	1.8	0.5	26.0	100.0	200
Lampung	71.1	66.0	1.3	0.1	14.6	2.5	42.3	3.9	1.4	0.0	5.0	1.8	3.1	0.2	28.9	100.0	925
Bangka Belitung	67.8	64.7	1.6	0.0	26.2	1.6	31.6	1.8	1.9	0.0	3.1	1.4	1.7	0.1	32.2	100.0	182
Riau Islands	57.6	54.0	2.2	0.0	17.6	3.0	27.3	1.1	2.8	0.1	3.5	2.3	1.0	0.1	42.4	100.0	134
Java																	
DKI Jakarta	60.1	56.4	2.7	0.4	13.8	6.5	27.2	2.1	3.6	0.3	3.7	2.2	1.4	0.1	39.9	100.0	1,352
West Java	61.1	60.3	1.5	0.4	19.4	5.1	31.0	1.3	1.6	0.0	0.8	0.6	0.2	0.0	38.9	100.0	5,243
Central Java	63.7	60.0	4.9	0.6	8.7	4.0	37.6	2.6	1.6	0.0	3.8	1.9	1.8	0.1	36.3	100.0	5,158
DI Yogyakarta	66.9	54.8	3.5	0.2	6.8	13.9	21.7	2.0	6.6	0.1	12.0	4.1	7.7	0.3	33.1	100.0	517
East Java	66.1	62.3	3.9	0.0	12.1	7.9	33.7	4.2	0.5	0.0	3.8	1.3	2.0	0.5	33.9	100.0	5,525
Banten	57.4	55.4	2.9	0.3	9.9	4.4	35.7	1.5	0.7	0.0	2.0	0.8	1.2	0.0	42.6	100.0	1,231
Bali and Nusa Tenggara																	
Bali	69.4	65.4	2.9	0.1	7.7	23.8	26.7	1.2	2.9	0.0	4.0	2.4	1.5	0.1	30.6	100.0	564
West Nusa Tenggara	54.8	52.2	2.3	0.2	7.0	4.6	33.5	4.3	0.4	0.0	2.7	0.6	0.6	1.4	45.2	100.0	636
East Nusa Tenggara	42.1	30.1	2.3	0.0	4.3	2.2	19.4	1.6	0.2	0.0	12.0	5.2	6.1	0.7	57.9	100.0	577
Kalimantan																	
West Kalimantan	62.7	61.2	2.4	0.0	15.5	2.2	38.5	1.7	0.9	0.0	1.5	0.4	0.3	0.8	37.3	100.0	590
Central Kalimantan	66.5	65.2	0.8	0.1	23.2	1.0	37.9	1.7	0.3	0.3	1.3	0.9	0.2	0.2	33.5	100.0	280
South Kalimantan	64.4	63.2	1.3	0.0	29.9	1.1	26.8	3.4	0.7	0.0	1.2	0.5	0.5	0.2	35.6	100.0	507
East Kalimantan	59.2	55.4	2.4	0.1	20.9	2.4	24.6	2.9	2.1	0.0	3.9	1.0	1.5	1.3	40.8	100.0	455
Sulawesi																	
North Sulawesi	69.3	66.7	1.6	0.0	23.1	5.9	29.2	6.6	0.3	0.0	2.6	2.2	0.3	0.1	30.7	100.0	360
Central Sulawesi	63.6	59.8	2.4	0.0	21.8	2.7	28.6	3.9	0.3	0.0	3.8	1.7	1.3	0.8	36.4	100.0	319
South Sulawesi	53.4	42.9	1.3	0.0	12.1	1.2	25.2	2.8	0.2	0.1	10.5	1.5	8.4	0.6	46.6	100.0	967
Southeast Sulawesi	50.7	44.4	1.7	0.0	16.3	0.9	19.7	5.1	0.6	0.1	6.2	3.0	3.1	0.2	49.3	100.0	242
Gorontalo	60.1	58.8	1.5	0.0	17.8	9.1	19.1	10.9	0.0	0.2	1.4	1.1	0.1	0.1	39.9	100.0	152
West Sulawesi	45.4	44.5	1.2	0.0	19.4	1.4	19.6	2.5	0.4	0.0	0.9	0.5	0.3	0.1	54.6	100.0	131
Maluku and Papua																	
Maluku	34.1	29.4	2.8	0.0	4.2	1.3	18.5	2.0	0.6	0.0	4.8	1.3	1.5	2.0	65.9	100.0	157
North Maluku	48.8	46.2	1.9	0.0	7.3	1.0	31.0	5.0	0.1	0.0	2.5	1.0	0.2	1.3	51.2	100.0	120
West Papua	39.6	37.5	2.8	0.4	6.8	1.3	23.9	2.3	0.0	0.0	2.1	0.7	0.2	1.1	60.4	100.0	83
Papua	38.3	24.5	2.5	0.2	5.9	1.3	11.5	3.2	0.0	0.0	13.8	0.9	2.1	10.8	61.7	100.0	242
Total	61.4	57.4	3.0	0.2	13.2	4.9	31.8	2.8	1.3	0.0	4.0	1.5	2.1	0.4	38.6	100.0	30,931

Note: If more than one method is used, only the most effective method is considered in this tabulation.
LAM = Lactational amenorrhea method.

Table A.5 Unmet need and the demand for family planning among currently married women by province

Percentage of currently married women age 15-49 with unmet need for family planning, percentage with met need for family planning, the total demand for family planning, and the percentage for the demand for contraception that is satisfied, by province, Indonesia 2007

Province	Unmet need for family planning ¹			Met need for family planning (currently using) ²			Total demand for family planning			Percentage of demand satisfied	Number of women
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total		
Sumatera											
DI Aceh	8.9	3.0	12.0	29.3	18.1	47.4	38.3	21.1	59.4	79.8	472
North Sumatera	5.1	7.2	12.3	15.2	38.9	54.2	20.8	46.3	67.1	81.6	1,389
West Sumatera	6.3	5.0	11.2	24.1	35.9	59.9	30.4	40.9	71.3	84.2	532
Riau	5.5	3.6	9.1	25.6	31.1	56.7	31.3	34.7	66.1	86.2	474
Jambi	3.5	3.6	7.0	30.0	35.2	65.2	33.7	38.7	72.5	90.3	346
South Sumatera	3.4	4.0	7.4	27.1	37.7	64.8	30.5	41.7	72.2	89.8	871
Bengkulu	2.7	3.4	6.1	31.0	43.0	74.0	33.9	46.5	80.4	92.4	200
Lampung	2.4	3.1	5.5	31.3	39.8	71.1	33.9	43.0	76.8	92.8	925
Bangka Belitung	1.9	1.3	3.2	33.6	34.2	67.8	35.5	35.5	71.0	95.5	182
Riau Islands	5.3	7.1	12.3	22.6	34.9	57.6	28.1	42.0	70.1	82.4	134
Java											
DKI Jakarta	3.2	3.7	6.9	23.3	36.9	60.1	26.5	40.6	67.1	89.7	1,352
West Java	4.6	5.4	10.0	26.1	35.0	61.1	30.8	40.5	71.3	85.9	5,243
Central Java	2.7	4.7	7.4	24.0	39.8	63.7	26.8	44.5	71.3	89.7	5,158
DI Yogyakarta	2.9	3.9	6.8	21.1	45.7	66.9	24.5	50.0	74.4	90.8	517
East Java	3.2	5.1	8.2	24.3	41.8	66.1	27.5	47.1	74.6	88.9	5,525
Banten	5.6	3.4	9.0	33.1	24.3	57.4	38.8	27.6	66.5	86.4	1,231
Bali and Nusa Tenggara											
Bali	2.2	3.6	5.8	13.9	55.5	69.4	16.1	59.1	75.2	92.3	564
West Nusa Tenggara	8.6	4.3	12.9	26.4	28.4	54.8	35.2	32.7	67.8	81.0	636
East Nusa Tenggara	9.8	7.7	17.4	20.9	21.2	42.1	31.0	29.1	60.2	71.0	577
Kalimantan											
West Kalimantan	4.7	3.0	7.7	30.9	31.8	62.7	35.6	34.8	70.4	89.1	590
Central Kalimantan	3.7	2.1	5.7	31.8	34.8	66.5	35.5	36.9	72.4	92.1	280
South Kalimantan	3.3	2.9	6.2	36.6	27.8	64.4	39.9	30.7	70.6	91.2	507
East Kalimantan	3.4	4.3	7.7	25.0	34.3	59.2	28.5	38.6	67.0	88.6	455
Sulawesi											
North Sulawesi	2.2	3.9	6.1	24.1	45.1	69.3	26.3	49.1	75.5	91.9	360
Central Sulawesi	4.0	4.3	8.3	27.9	35.7	63.6	32.5	40.1	72.6	88.6	319
South Sulawesi	9.2	4.6	13.9	24.8	28.6	53.4	34.1	33.3	67.4	79.4	967
Southeast Sulawesi	8.5	4.4	12.9	27.6	23.0	50.7	36.4	27.6	64.0	79.9	242
Corontalo	4.9	1.8	6.6	23.2	36.9	60.1	28.2	38.7	66.9	90.1	152
West Sulawesi	12.3	5.0	17.4	26.7	18.7	45.4	39.1	23.7	62.8	72.4	131
Maluku and Papua											
Maluku	8.9	13.5	22.4	13.2	21.0	34.1	22.1	34.5	56.6	60.3	157
North Maluku	6.7	6.3	13.0	25.3	23.5	48.8	32.0	29.8	61.8	78.9	120
West Papua	12.2	4.3	16.6	18.4	21.2	39.6	30.6	25.5	56.2	70.5	83
Papua	7.7	8.1	15.8	19.5	18.8	38.3	27.2	26.9	54.1	70.8	242
Total	4.3	4.7	9.1	25.1	36.3	61.4	29.5	41.1	70.6	87.2	30,931

¹ Unmet need for spacing includes pregnant women whose pregnancy was mistimed; amenorrheic women who are not using family planning and whose last birth was mistimed, or whose last birth was unwanted but now say they want more children; and fecund women who are neither pregnant nor amenorrheic, who are not using any method of family planning, and say they want to wait two or more years for their next birth. Also included in unmet need for spacing are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted; amenorrheic women who are not using family planning, whose last child was unwanted and who do not want any more children; and fecund women who are neither pregnant nor amenorrheic, who are not using any method of family planning, and who want no more children

² Using for spacing is defined as women who are using some method of family planning and say they want to have another child or are undecided whether to have another. Using for limiting is defined as women who are using and who want no more children. Note that the specific methods used are not taken into account here

Table A.6 Maternal care indicators by province

Percentage of women who had a live birth in the five years preceding the survey who received specific maternal health services during pregnancy for the most recent birth, and among all live births in the five years before the survey, percentage delivered by a health professional and percentage delivered in a health facility, by province, Indonesia 2007

Province	Percentage with antenatal care from health professional	Percentage given at least one tetanus toxoid injection during pregnancy	Percentage given iron tablets or syrup during pregnancy	Number of women	Percentage delivered by a health professional ¹	Percentage delivered in a health facility	Number of births
Sumatera							
DI Aceh	89.2	56.0	57.0	269	72.5	24.8	324
North Sumatera	89.3	33.7	62.7	803	84.5	35.0	1,197
West Sumatera	96.0	81.8	77.3	304	80.5	63.5	383
Riau	93.2	64.5	65.9	243	84.9	43.6	290
Jambi	84.6	68.9	70.9	169	69.8	26.1	186
South Sumatera	91.3	65.5	69.6	424	67.5	33.4	491
Bengkulu	93.7	76.9	78.8	100	72.3	12.1	111
Lampung	95.4	78.4	81.3	409	69.8	44.7	452
Bangka Belitung	94.0	77.1	78.5	93	81.5	43.2	103
Riau Islands	93.8	54.7	77.5	76	91.6	76.3	93
Java							
DKI Jakarta	99.5	76.4	80.3	649	97.3	88.4	741
West Java	95.3	81.4	82.1	2,328	68.2	44.6	2,600
Central Java	97.3	81.8	83.5	2,109	83.0	53.1	2,308
DI Yogyakarta	99.3	85.8	94.1	179	95.8	86.8	201
East Java	93.2	68.1	80.7	1,947	77.5	65.5	2,178
Banten	86.3	64.1	70.1	599	52.1	38.1	695
Bali and Nusa Tenggara							
Bali	98.8	87.9	92.8	225	92.6	90.8	253
West Nusa Tenggara	95.3	70.8	87.4	347	64.3	32.2	412
East Nusa Tenggara	87.1	76.9	83.4	375	46.2	20.7	507
Kalimantan							
West Kalimantan	90.6	68.6	54.4	312	62.2	33.7	374
Central Kalimantan	91.0	75.7	72.8	138	68.1	14.1	160
South Kalimantan	93.0	69.9	75.7	249	75.6	19.4	289
East Kalimantan	93.9	83.2	81.3	218	75.5	45.8	262
Sulawesi							
North Sulawesi	95.9	86.6	88.5	166	87.3	55.0	191
Central Sulawesi	90.7	80.8	75.7	192	59.6	19.0	243
South Sulawesi	92.2	82.5	71.9	500	58.8	30.6	631
Southeast Sulawesi	91.3	80.1	58.9	144	56.6	8.4	192
Gorontalo	88.5	72.1	67.3	68	53.6	21.7	82
West Sulawesi	86.6	77.5	59.1	75	43.8	12.6	103
Maluku and Papua							
Maluku	70.3	57.0	57.0	99	32.8	12.4	143
North Maluku	88.0	85.7	84.1	71	45.9	18.0	93
West Papua	80.4	64.3	65.1	45	57.7	39.0	62
Papua	69.0	50.5	56.3	117	46.3	26.2	152
Total	93.3	73.0	77.3	14,043	73.0	46.1	16,504

¹ Doctor, obgyn, nurse, midwife, or village midwife

Table A.7 Vaccinations by province

Percentage of children age 12-23 months who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report), and percentage with a vaccination card seen, by province, Indonesia 2007

Province	DPT			Polio				Measles	All ¹	No vacci-nations	Hepatitis B			Percentage with vacci-nation card	Number of children	
	BCG	1	2	3	1	2	3				4	1	2			3
Sumatera																
DI Aceh	63.5	58.4	46.8	33.7	67.5	54.3	46.9	22.0	50.1	26.8	26.0	50.3	38.3	27.4	16.2	63
North Sumatera	66.2	65.3	52.8	39.2	81.0	72.3	60.7	31.6	52.4	32.8	15.5	58.4	48.6	38.1	28.2	231
West Sumatera	87.1	85.2	73.6	69.0	91.4	84.5	75.2	65.5	72.7	60.2	5.1	81.3	76.5	71.9	36.6	71
Riau	73.3	74.5	69.7	52.1	74.5	66.9	56.8	39.3	68.9	41.4	20.6	73.7	58.7	49.3	26.8	60
Jambi	71.7	75.1	65.4	55.2	75.1	72.4	64.3	44.1	61.7	44.8	24.9	66.3	54.9	41.5	34.3	35
South Sumatera	91.0	88.0	80.3	67.8	93.1	84.3	70.9	43.6	73.0	54.6	5.8	84.5	73.4	57.5	32.8	80
Bengkulu	89.6	86.4	78.8	61.6	95.7	89.6	79.0	62.3	81.0	54.9	1.4	79.0	72.4	61.6	33.8	21
Lampung	93.4	94.1	83.8	78.5	96.8	94.0	86.4	68.1	83.5	67.0	0.9	90.9	81.9	72.5	42.4	110
Bangka Belitung	76.7	74.8	71.9	69.3	80.1	76.7	70.3	53.1	68.9	59.3	16.1	71.1	66.1	64.3	35.5	21
Riau Islands	82.8	82.8	73.0	68.6	90.9	84.7	75.3	62.0	82.4	62.5	8.2	79.5	72.6	64.3	23.2	20
Java																
DKI Jakarta	88.4	87.0	80.8	77.4	90.9	85.9	82.6	51.5	79.7	71.5	8.7	87.1	81.1	67.3	27.8	133
West Java	89.6	89.1	78.4	70.0	91.6	84.6	74.9	59.1	81.2	63.9	5.7	85.6	77.6	66.0	45.7	543
Central Java	95.7	96.5	93.2	88.3	98.2	93.8	91.5	76.7	87.1	74.7	0.9	94.9	90.1	77.2	49.1	430
DI Yogyakarta	100.0	100.0	99.8	97.0	100.0	100.0	100.0	95.6	95.2	93.8	0.0	100.0	94.5	88.1	56.7	35
East Java	87.1	87.2	79.7	70.7	89.3	83.3	74.7	62.4	80.3	64.8	6.3	82.9	71.2	63.3	39.4	353
Banten	80.4	74.7	63.1	48.8	92.0	79.3	63.1	35.7	76.6	37.4	5.5	72.0	61.4	48.3	17.5	123
Bali and Nusa Tenggara																
Tenggara																
Bali	94.3	95.9	87.9	77.3	91.9	89.4	86.7	71.6	85.5	72.2	3.4	92.3	86.1	78.2	46.5	49
West Nusa Tenggara	86.3	85.3	77.9	62.0	87.5	84.4	67.9	49.6	80.3	55.7	11.4	82.6	74.4	59.4	27.4	78
East Nusa Tenggara	86.5	80.8	67.3	52.6	88.1	83.0	58.1	36.0	77.2	45.7	10.0	69.1	53.8	36.9	25.4	95
Kalimantan																
West Kalimantan	81.2	81.1	68.1	63.5	76.4	70.3	63.6	50.9	68.2	50.1	17.2	70.4	65.6	58.5	31.1	73
Central Kalimantan	80.7	77.7	58.1	50.1	91.0	85.0	56.0	38.3	83.3	42.2	4.7	80.3	63.3	47.6	15.1	28
South Kalimantan	81.3	76.2	62.5	55.0	82.4	77.3	62.3	55.2	62.1	50.8	16.4	69.8	60.8	52.9	32.0	48
East Kalimantan	85.9	87.1	83.2	75.0	89.1	86.8	80.6	73.3	82.4	69.2	10.9	83.7	76.1	67.3	46.2	53
Sulawesi																
North Sulawesi	96.2	94.0	89.7	81.8	94.5	93.6	88.4	59.9	86.2	76.1	3.8	87.9	78.4	58.6	38.4	38
Central Sulawesi	80.5	81.2	75.2	57.3	85.6	79.6	72.1	45.6	72.3	50.3	12.2	75.4	63.9	46.3	36.5	43
South Sulawesi	79.8	76.7	69.2	61.8	79.1	76.0	67.8	53.6	69.0	55.1	17.8	73.8	66.9	54.1	27.5	112
Southeast Sulawesi	85.4	83.6	72.5	65.9	88.6	78.0	70.6	47.6	79.5	64.1	9.6	80.0	73.1	60.5	37.1	37
Corontalo	83.2	81.5	68.4	58.6	80.4	75.0	70.3	50.5	69.3	55.2	13.4	75.7	67.5	54.7	29.5	16
West Sulawesi	80.6	76.2	70.3	59.2	85.1	79.1	64.7	46.1	74.9	53.0	14.5	74.8	66.0	50.6	37.0	15
Maluku and Papua																
Maluku	67.7	67.4	55.8	44.2	69.3	56.2	45.2	33.1	58.4	39.7	27.3	58.6	50.3	39.2	17.6	24
North Maluku	73.9	63.8	55.4	41.5	79.7	62.1	49.3	34.3	70.2	37.1	20.3	72.2	59.0	35.2	41.0	14
West Papua	61.0	62.4	52.5	35.2	72.4	66.2	55.8	24.1	55.3	23.7	26.7	52.6	40.2	31.4	21.6	12
Papua	72.0	70.2	52.1	40.9	72.0	55.8	38.0	32.6	63.0	32.6	26.7	65.4	44.1	35.4	25.5	29
Total	85.4	84.4	75.7	66.7	88.8	82.6	73.5	55.5	76.4	58.6	8.6	80.5	71.7	60.3	36.8	3,094

¹ BCG, measles and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

Table A.8 Treatment for acute respiratory infection, fever, and diarrhea by province

Among children under five years who were sick with a cough accompanied by short, rapid breathing (symptoms of acute respiratory infection - ARI) or fever in the two weeks preceding the survey, percentage for whom treatment was sought from a health facility or provider, and among children under five years who were sick with diarrhea during the two weeks preceding the survey, percentage for whom treatment was sought from a health facility or provider, percentage given a solution made from oral rehydration salt (ORS) packets, and percentage given any oral rehydration therapy (ORT), by province, Indonesia 2007

Province	Children with symptoms of ARI or with fever		Children with diarrhea			
	Percentage for whom treatment was sought from a health facility/provider ¹	Number with ARI/fever	Percentage for whom treatment was sought from a health facility/provider ¹	Percentage given solution from ORS packet	Percentage given any ORT ²	Number with diarrhea
Sumatera						
DI Aceh	91.7	129	75.7	36.2	67.8	60
North Sumatera	74.9	420	50.8	25.0	46.8	181
West Sumatera	73.5	158	54.0	37.1	81.0	53
Riau	84.6	102	57.3	45.9	56.8	47
Jambi	81.1	50	77.7	55.3	80.4	27
South Sumatera	75.6	123	53.6	49.9	66.5	69
Bengkulu	80.7	45	49.8	33.8	67.3	22
Lampung	75.2	126	(63.5)	(34.0)	(59.5)	47
Bangka Belitung	85.4	34	(76.4)	(53.9)	(71.2)	6
Riau Islands	87.1	32	64.5	45.7	58.9	13
Java						
DKI Jakarta	88.2	178	69.2	48.2	71.0	50
West Java	71.3	939	53.5	30.1	56.8	455
Central Java	82.2	665	53.3	23.2	52.2	211
DI Yogyakarta	86.6	53	*	*	*	11
East Java	83.5	755	64.6	32.0	62.5	279
Banten	66.9	238	61.3	27.3	46.0	68
Bali and Nusa Tenggara						
Bali	88.9	56	86.5	54.3	67.8	23
West Nusa Tenggara	72.5	174	47.6	43.2	68.8	70
East Nusa Tenggara	80.0	199	61.8	57.5	83.0	73
Kalimantan						
West Kalimantan	71.6	128	56.1	41.4	62.3	55
Central Kalimantan	73.1	58	70.8	48.5	66.5	33
South Kalimantan	70.7	111	56.4	34.4	58.9	43
East Kalimantan	81.7	86	48.8	39.6	61.4	35
Sulawesi						
North Sulawesi	84.5	64	69.3	33.0	67.2	26
Central Sulawesi	65.5	108	52.0	34.2	68.4	37
South Sulawesi	73.1	213	55.4	32.8	60.0	71
Southeast Sulawesi	55.1	74	48.6	33.9	64.1	26
Corontalo	62.4	37	59.6	38.7	75.6	13
West Sulawesi	64.0	46	56.2	35.2	64.3	21
Maluku and Papua						
Maluku	53.9	40	27.0	20.9	65.1	13
North Maluku	76.4	36	55.4	35.5	64.4	12
West Papua	86.9	13	78.3	56.7	69.6	8
Papua	83.3	47	79.4	57.7	73.6	22
Total	76.9	5,539	57.9	34.7	60.9	2,180

¹ Excludes pharmacy, shop, traditional practitioner

² Includes ORS, recommended home fluid, or increased fluids

Table A.9 Early childhood mortality rates by province

Neonatal, postneonatal, infant, child, and under-five mortality rates for the 10-year period preceding the survey, by province, Indonesia 2007

Province	Neonatal mortality (NN)	Postneonatal mortality (PNN) ¹	Infant mortality (₁ q ₀)	Child mortality (₄ q ₁)	Under-five mortality (₅ q ₀)
Sumatera					
DI Aceh	14	11	25	21	45
North Sumatera	24	22	46	22	67
West Sumatera	34	13	47	16	62
Riau	28	8	37	11	47
Jambi	23	15	39	9	47
South Sumatera	25	17	42	11	52
Bengkulu	17	29	46	20	65
Lampung	27	16	43	13	55
Bangka Belitung	20	19	39	8	46
Riau Islands	18	25	43	16	58
Java					
DKI Jakarta	15	13	28	9	36
West Java	19	19	39	10	49
Central Java	14	12	26	6	32
DI Yogyakarta	15	3	19	3	22
East Java	21	14	35	10	45
Banten	25	21	46	13	58
Bali and Nusa Tenggara					
Bali	14	19	34	4	38
West Nusa Tenggara	34	38	72	21	92
East Nusa Tenggara	31	26	57	24	80
Kalimantan					
West Kalimantan	23	23	46	14	59
Central Kalimantan	13	17	30	4	34
South Kalimantan	39	19	58	19	75
East Kalimantan	16	11	26	12	38
Sulawesi					
North Sulawesi	24	11	35	9	43
Central Sulawesi	28	31	60	10	69
South Sulawesi	22	19	41	12	53
Southeast Sulawesi	16	25	41	21	62
Gorontalo	22	31	52	18	69
West Sulawesi	46	27	74	25	96
Maluku and Papua					
Maluku	25	34	59	37	93
North Maluku	32	19	51	24	74
West Papua	21	16	36	26	62
Papua	24	17	41	25	64

¹ Computed as the difference between the infant and neonatal mortality rates

Table A.10. Knowledge of AIDS by province

Percentage of ever-married women and currently married men who have heard of AIDS, by province, Indonesia 2007

Province	Ever-married women		Currently married men	
	Percentage who have heard of AIDS	Number of women	Percentage who have heard of AIDS	Number of men
Sumatera				
DI Aceh	43.4	514	56.6	137
North Sumatera	60.9	1,487	82.1	370
West Sumatera	66.2	570	72.4	137
Riau	67.6	494	85.7	130
Jambi	54.0	367	78.6	95
South Sumatera	51.5	928	46.4	241
Bengkulu	56.7	211	66.0	53
Lampung	62.0	963	72.9	271
Bangka Belitung	58.2	194	75.8	52
Riau Islands	68.8	140	89.1	36
Java				
DKI Jakarta	91.0	1,471	97.2	408
West Java	67.5	5,545	77.0	1,444
Central Java	63.3	5,383	72.2	1,517
DI Yogyakarta	81.7	551	93.7	146
East Java	55.2	5,924	65.0	1,561
Banten	56.8	1,310	59.4	344
Bali and Nusa Tenggara				
Bali	72.9	587	83.9	174
West Nusa Tenggara	42.5	705	57.4	194
East Nusa Tenggara	35.6	627	51.5	172
Kalimantan				
West Kalimantan	45.1	628	71.2	162
Central Kalimantan	55.7	294	66.6	82
South Kalimantan	66.1	550	85.6	128
East Kalimantan	69.3	475	66.0	132
Sulawesi				
North Sulawesi	76.7	373	83.3	102
Central Sulawesi	52.5	339	66.9	89
South Sulawesi	48.0	1,067	57.1	259
Southeast Sulawesi	51.8	259	81.2	60
Gorontalo	41.2	163	53.0	46
West Sulawesi	42.4	139	41.8	41
Maluku and Papua				
Maluku	55.6	168	58.7	44
North Maluku	46.9	129	64.6	36
West Papua	60.0	89	89.3	24
Papua	56.4	251	66.0	70
Total	61.0	32,895	71.4	8,758

Table A.11 Knowledge of ways to reduce the risk of getting the AIDS virus by province

Percentage of ever-married women and currently married men who, in response to prompted questions, say that people can reduce the risk of getting the AIDS virus by using condoms every time they have sexual intercourse, by having one uninfected sex partner who has no other partners, and by abstaining from sexual intercourse, by background characteristics, Indonesia 2007

Province	Ever-married women					Currently married men				
	Percentage using condoms	Percentage limiting sexual intercourse to one uninfected partner	Percentage limiting sexual intercourse to one uninfected partner	Percentage abstaining from sexual intercourse	Number of women	Percentage using condoms	Percentage limiting sexual intercourse to one uninfected partner	Percentage limiting sexual intercourse to one uninfected partner	Percentage abstaining from sexual intercourse	Number of men
Sumatera										
DI Aceh	15.8	21.7	11.5	16.2	514	27.4	28.7	17.7	30.8	137
North Sumatera	33.8	36.6	28.0	33.1	1,487	63.2	55.4	47.7	49.3	370
West Sumatera	39.5	53.2	35.0	47.1	570	40.3	49.5	32.5	42.1	137
Riau	38.4	44.2	31.9	38.7	494	60.7	69.5	57.0	58.2	130
Jambi	24.7	32.3	19.4	23.8	367	52.6	49.5	42.0	50.5	95
South Sumatera	22.7	32.4	19.5	27.8	928	28.8	31.8	25.2	29.7	241
Bengkulu	36.1	38.8	29.5	36.4	211	45.1	57.0	42.5	46.5	53
Lampung	34.4	46.2	31.5	39.0	963	36.7	53.5	31.7	30.9	271
Bangka Belitung	35.8	41.2	30.1	30.2	194	34.4	23.6	18.2	29.8	52
Riau Islands	38.5	44.8	32.5	41.2	140	63.7	69.2	54.0	50.1	36
Java										
DKI Jakarta	61.7	73.6	57.8	66.3	1,471	85.6	86.4	78.4	64.9	408
West Java	41.8	52.0	36.5	47.0	5,545	51.5	64.1	47.4	47.2	1,444
Central Java	34.7	43.3	28.9	34.2	5,383	47.7	42.8	33.7	41.3	1,517
DI Yogyakarta	63.5	70.1	60.2	56.0	551	81.6	91.0	79.5	80.9	146
East Java	35.8	37.3	29.1	35.9	5,924	43.6	48.1	37.8	36.0	1,561
Banten	27.8	34.5	22.4	25.0	1,310	39.8	41.6	34.4	36.8	344
Bali and Nusa Tenggara										
Bali	39.0	42.4	30.4	33.4	587	74.5	76.1	69.4	47.1	174
West Nusa Tenggara	19.7	20.2	12.4	18.8	705	22.5	26.7	17.9	16.8	194
East Nusa Tenggara	16.0	20.3	12.7	19.1	627	38.6	44.3	35.9	38.4	172
Kalimantan										
West Kalimantan	22.5	27.7	19.4	22.7	628	42.8	52.0	36.5	45.5	162
Central Kalimantan	36.2	42.6	30.6	41.0	294	45.7	48.9	36.6	37.4	82
South Kalimantan	30.1	48.0	25.6	32.4	550	71.2	77.0	66.6	73.5	128
East Kalimantan	38.6	41.6	30.1	36.0	475	51.2	56.1	47.5	52.9	132
Sulawesi										
North Sulawesi	44.7	53.2	36.6	41.2	373	65.5	59.6	52.0	44.2	102
Central Sulawesi	23.6	27.6	16.3	21.4	339	37.9	43.7	32.4	44.4	89
South Sulawesi	27.5	32.0	21.8	28.4	1,067	40.5	42.7	36.1	43.3	259
Southeast Sulawesi	22.5	30.3	17.4	23.3	259	54.4	50.9	34.0	40.7	60
Corontalo	16.0	20.7	12.1	18.4	163	32.1	26.2	22.1	26.5	46
West Sulawesi	19.2	22.1	13.9	19.1	139	26.3	25.9	20.7	30.5	41
Maluku and Papua										
Maluku	35.5	44.3	31.0	43.4	168	33.3	35.5	27.6	28.9	44
North Maluku	14.5	16.3	9.1	13.4	129	36.9	43.6	29.6	25.0	36
West Papua	31.6	35.5	23.9	32.4	89	71.1	58.0	49.9	51.1	24
Papua	32.7	31.0	23.3	31.8	251	41.0	37.8	26.3	39.5	70
Total	35.5	42.2	29.9	36.6	32,895	48.9	52.4	41.3	42.9	8,758

na = Not applicable

¹Using condoms every time they have sexual intercourse

²Partner who has no other partners

Table A.12 Knowledge about voluntary counseling and testing (VCT)

Percentage of ever-married women age 15-49 who know about voluntary HIV counseling and testing (VCT) and percentage who know specific VCT providers, by province, Indonesia 2007

Province	Ever-married women		Among ever-married women who know about VCT, percentage who know specific providers											Number of ever-married women who know VCT	
	Percentage who know about VCT	Number of currently married women	Public					Private					Other		
			Public hospital	Public health center	Public clinic	Public VCT clinic	Other public	Private hospital	Private clinic	Private VCT clinic	Private doctor	Private nurse/midwife			Other private medical
Sumatera															
DI Aceh	4.0	514	(77.2)	(18.7)	(2.1)	(21.2)	(0.7)	(0.0)	(2.9)	(10.5)	(3.2)	(0.0)	(0.0)	(0.0)	21
North Sumatera	9.9	1,487	64.7	12.3	2.2	10.7	0.0	25.5	2.0	9.4	6.9	5.6	4.3	3.4	147
West Sumatera	13.9	570	76.8	18.1	4.4	5.5	4.0	8.9	0.0	1.1	10.3	3.2	0.9	6.3	79
Riau	5.5	494	67.2	16.3	0.0	3.6	0.9	8.9	0.0	5.7	1.5	1.7	3.9	10.8	27
Jambi	2.8	367	*	*	*	*	*	*	*	*	*	*	*	*	10
South Sumatera	3.7	928	(74.4)	(7.5)	(0.0)	(12.4)	(0.0)	(11.1)	(0.0)	(0.0)	(2.2)	(2.2)	(0.0)	(16.2)	34
Bengkulu	7.9	211	75.8	11.8	2.2	7.0	0.0	1.9	0.0	0.0	4.3	0.1	0.0	6.0	17
Lampung	4.0	963	(47.0)	(17.0)	(0.0)	(5.4)	(0.0)	(14.0)	(3.4)	(2.8)	(19.8)	(2.8)	(6.9)	(19.9)	38
Bangka Belitung	7.3	194	78.9	10.6	3.7	16.6	0.0	5.5	0.0	10.0	2.5	0.0	0.0	1.7	14
Riau Islands	7.4	140	(44.4)	(24.4)	(0.0)	(2.5)	(1.7)	(11.0)	(0.0)	(4.7)	(2.4)	(0.0)	(14.2)	(9.4)	10
Java															
DKI Jakarta	26.6	1,471	78.3	5.8	0.4	11.7	0.5	21.1	0.2	2.4	1.7	1.2	1.6	1.3	391
West Java	8.3	5,545	67.6	17.1	0.9	5.1	3.1	19.6	1.7	3.3	6.5	2.4	1.6	10.6	461
Central Java	5.7	5,383	65.8	10.3	0.0	8.9	1.2	13.2	0.0	6.7	8.0	1.2	3.5	2.7	304
DI Yogyakarta	15.8	551	80.3	17.5	1.2	4.0	3.7	28.9	1.8	2.4	7.1	3.9	4.4	0.0	87
East Java	5.0	5,924	56.0	19.4	4.0	4.2	0.0	6.4	3.2	0.0	16.1	7.6	2.7	8.6	294
Banten	5.5	1,310	49.4	16.0	2.8	6.4	0.0	8.0	4.8	1.1	5.1	2.2	1.8	14.8	72
Bali and Nusa Tenggara															
Bali	5.2	587	78.7	14.6	0.0	4.6	0.0	10.7	0.3	3.7	6.2	5.0	2.7	3.3	30
West Nusa Tenggara	7.9	705	81.9	15.6	1.4	12.5	0.0	15.1	0.0	7.7	12.7	1.6	0.0	3.2	56
East Nusa Tenggara	2.7	627	*	*	*	*	*	*	*	*	*	*	*	*	17
Kalimantan															
West Kalimantan	7.4	628	85.3	23.2	5.0	9.8	0.0	3.9	2.8	7.1	2.0	2.0	0.0	0.0	46
Central Kalimantan	3.9	294	(69.5)	(18.1)	(3.8)	(14.0)	(3.2)	(5.4)	(0.0)	(13.1)	(13.4)	(7.7)	(0.0)	(1.9)	12
South Kalimantan	5.0	550	54.3	22.1	0.0	4.8	0.0	0.0	0.0	4.3	17.5	2.2	6.1	4.6	27
East Kalimantan	9.7	475	79.2	10.4	0.0	4.9	4.6	3.3	1.3	0.0	2.7	0.0	2.3	1.7	46
Sulawesi															
North Sulawesi	19.7	373	89.8	8.4	2.1	15.4	0.6	11.3	1.7	6.2	8.4	1.6	0.4	0.8	74
Central Sulawesi	8.7	339	89.1	3.1	1.5	19.9	0.0	6.2	0.0	3.9	4.3	0.0	0.0	0.0	30
South Sulawesi	6.0	1,067	78.0	16.8	0.0	6.8	0.0	9.8	0.0	2.7	2.8	1.4	4.2	0.0	64
Southeast Sulawesi	7.9	259	78.2	20.9	3.7	5.2	0.0	9.0	0.0	0.0	7.5	0.0	2.0	2.7	20
Gorontalo	6.2	163	72.6	15.5	0.0	13.4	0.0	4.6	2.0	0.0	8.6	0.0	0.0	0.0	10
West Sulawesi	4.4	139	(64.6)	(16.6)	(0.0)	(7.3)	(0.0)	(4.6)	(0.0)	(0.0)	(14.2)	(6.4)	(0.0)	(7.8)	6
Maluku and Papua															
Maluku	7.9	168	85.5	13.0	1.3	4.5	0.0	21.8	0.0	0.0	8.3	0.0	0.0	5.1	13
North Maluku	3.2	129	*	*	*	*	*	*	*	*	*	*	*	*	4
West Papua	21.2	89	74.7	4.9	0.0	18.5	0.0	13.0	0.0	15.0	0.4	0.0	0.0	2.2	19
Papua	6.1	251	(83.7)	(20.1)	(4.8)	(4.6)	(0.0)	(8.2)	(0.0)	(0.0)	(4.4)	(4.4)	(2.2)	(4.0)	15
Total	7.6	32,895	69.8	13.8	1.5	8.2	1.2	14.8	1.2	3.7	7.2	2.7	2.3	5.5	2,497

MEASURE DHS Preliminary Reports

Turkey 1998	December	1998	English
Ghana 1998	May	1999	English
Guatemala 1997	June	1999	Spanish
Guinea 1999	October	1999	French
Kazakhstan 1999	December	1999	English/Russian
Tanzania 1999	February	2000	English
Zimbabwe 1999	March	2000	English
Bangladesh 1999-2000	June	2000	English
Egypt 2000	June	2000	English
Ethiopia 2000	July	2000	English
Haiti 2000	September	2000	French
Cambodia 2000	November	2000	English
Turkmenistan 2000	January	2001	English
Malawi 2000	February	2001	English
Rwanda 2000	February	2001	French
Azerbaijan 2000	March	2001	English
Gabon 2000	March	2001	French
Uganda 2001	May	2001	English
Mauritania 2001	July	2001	French
Nepal 2001	August	2001	English
Mali 2001	December	2001	French
Nicaragua 2001	December	2001	Spanish
Benin 2001	February	2002	French
Eritrea 2002	September	2002	English
Zambia 2002	October	2002	English
Jordan 2002	November	2002	English
Dominican Republic 2002	January	2003	Spanish
Uzbekistan 2002	May	2003	English
Indonesia 2003	August	2003	English
Nigeria 2003	October	2003	English
Kenya 2003	December	2003	English
Indonesia (young adult) 2002	December	2003	English
Jayapura City, Indonesia (young adult) 2002	December	2003	English
Philippines 2003	January	2004	English
Bolivia 2003	March	2004	Spanish
Mozambique 2003	April	2004	English
Burkina Faso 2002-03	May	2004	French
Ghana 2003	June	2004	English
Morocco 2003-04	June	2004	French
Bangladesh 2004	September	2004	English
Cameroon 2004	October	2004	French
Madagascar 2003-04	October	2004	French
Chad 2004	February	2006	French
Kenya (SPA) 2004	March	2006	English
Peru Continuous 2004-05	April	2006	Spanish
Tanzania 2004-05	May	2006	English
Uganda (AIS) 2004-05	June	2006	English
Malawi 2004	August	2006	English
Senegal 2006	August	2006	French
Guinea 2006	August	2006	French
Lesotho 2004	September	2006	English
Egypt 2006	September	2006	English
Rwanda 2006	November	2006	French
Ethiopia 2006	November	2006	English
Moldova 2006	November	2006	English/Romanian
Vietnam (AIS) 2006	February	2006	English/Vietnamese
Armenia 2005	March	2006	English
Congo (Brazzaville) 2005	March	2006	French
Côte d'Ivoire (AIS) 2005	June	2006	French
Cambodia 2005	July	2006	English
Haiti 2005-06	July	2006	French
Zimbabwe 2005-06	August	2006	English
Niger 2006	August	2006	French
Niger (Intervention zones) 2006	October	2006	French
Nepal 2006	October	2006	English
Uganda 2006	November	2006	English
Tanzania (SPA) 2006	January	2007	English
Benin 2006	March	2007	French
Azerbaijan 2006	April	2007	English
Mali 2006	April	2007	French
Pakistan 2006-07	June	2007	English
Swaziland 2006-07	June	2007	English
Liberia 2007	July	2007	English
Democratic Republic of Congo 2007	December	2007	French
Bangladesh 2007	December	2007	English
Rwanda (SPA) 2007	December	2007	English/French
Jordan 2007	January	2008	English/Arabic
Uganda (SPA) 2007	March	2008	English
Ukraine 2007	June	2008	English
Indonesia 2007	July	2008	English
Indonesia (young adult) 2007	July	2008	English

MEASURE DHS Preliminary Reports are distributed to a limited number of recipients needing early access to survey findings and are not available for general distribution. The national implementing agency is responsible for in-country distribution; MEASURE DHS is responsible for external distribution. Publication of MEASURE DHS final reports meant for general distribution is expected 9 to 12 months after publication of the preliminary report.