

Healthy Timing & Spacing of Pregnancy (HTSP) In Indonesia

HTSP is an intervention to help women and families delay or space their pregnancies to achieve the healthiest outcomes for women, newborns, infants, and children, *within the context of free and informed choice*

Summary

Key research findings regarding the risks of closely spaced pregnancies:

- Recent USAID-sponsored research found that unhealthy pregnancy spacing is associated with multiple adverse outcomes for mothers and newborns.
- Becoming pregnant too soon after a previous birth, miscarriage, or abortion places mothers and newborns at a higher risk of health complications—or even death.
- Early pregnancy (when mother is younger than 18) the mothers and their newborns face increased risks of health complications compared to women 20-24 years old.

In Indonesia

- Among married women of reproductive age (MWRA) 13% of births are spaced <2 years apart
- Among mothers between 15 and 19 years old, 33% of births are spaced <2 years apart

Considerations for program design:

- When women and their families are advised of the benefits of pregnancy delay and spacing, use of family planning increases substantially.
- How can improved HTSP advocacy, behavior change communication, counseling, and community outreach play a role in helping women achieve the longer intervals they want and reduce the risk of the adverse health outcomes?

Consider these findings in relation to the overall risks to mothers and newborns in Indonesia.

Annual number of births*	4,495,000
Annual number of neonatal deaths**	81,000
Neonatal mortality rate*	18
Annual number of infant deaths (includes neonatal)**	126,000
Infant mortality rate*	28
Annual number of <5 deaths (includes both infant and neonatal deaths)*	162,000
Child mortality rate*	36
Annual number of low birth weight infants**	404,550
% of infants with low birth weight*	9
Annual number of maternal deaths**	10,330
Maternal mortality rate (UNICEF adjusted #)*	230
Lifetime risk of maternal death when a woman becomes pregnant*	1 in 150

Healthy timing and spacing of pregnancy may help reduce the consequences of these health risks for mothers and newborns.

Source: *UNICEF, 2007 State of the World's Children
** Calculated

Introduction

Recent research from developing countries shows that unhealthy timing or spacing of pregnancies is linked to increased risk of multiple adverse health outcomes. Table 1 shows how it can affect the health of the mother and the newborn.

Following a pregnancy that occurred quickly after a previous birth, the risk of a child dying is at least twice as high as that for longer intervals.

An infant born after a short interval has increased chances of:

- Being born pre-term
- Having below normal weight at birth
- Being small for gestational age

A woman who becomes pregnant too quickly following a previous birth, or induced abortion or miscarriage, faces higher risks of:

- Anemia
- Premature rupture of membranes
- Abortion
- Miscarriage
- Death

Table 1. Risks of Adverse Health Outcomes After Very Short Interval Pregnancy, Compared to the Reference Group Interval Used in the Selected Study

RISKS WHEN PREGNANCY OCCURS 6 MONTHS AFTER A LIVE BIRTH		
Adverse Outcome	Increased Risk	
Induced Abortion	650%	
Miscarriage	230%	
Newborn Death (<9 mos.)	170%	
Maternal Death	150%	
Preterm Birth	70%	
Stillborn	60%	
Low Birth Weight	60%	
RISKS WHEN PREGNANCY OCCURS QUICKLY AFTER AN ABORTION OR MISCARRIAGE, COMPARED TO RISKS AFTER WAITING 6 MONTHS		
Increased Risk After 1-2 Mos. Interval	Increased Risk After 3-5 Mos. Interval	
Low Birth Weight	170%	140%
Maternal Anemia	160%	120%
Preterm Birth	80%	40%
<small>Sources: Conde-Agudelo, et al, 2000, 2005, 2006; Da Vanzo, et al, 2004; Razzaque, et al, 2005; Rutstein, 2005.</small>		

World Health Organization (WHO) Recommendations

Based on a review of six USAID-supported studies, WHO produced a policy brief in 2006 on birth spacing which included the following preamble and recommendations:

Preamble

Individuals and couples should consider health risks and benefits along with other circumstances such as their age, fecundity, fertility aspirations, access to health services, child-rearing support, social and economic circumstances, and personal preferences in making choices for the timing of the next pregnancy.

Recommendation for spacing after a live birth

After a live birth, the recommended interval before attempting the next pregnancy is at least 24 months in order to reduce the risk of adverse maternal, perinatal and infant outcomes.

Recommendation for spacing after an abortion

After a miscarriage or induced abortion, the recommended minimum interval to next pregnancy is at least six months in order to reduce risks of adverse maternal and perinatal outcomes.

Source: World Health Organization, 2006 Report of a WHO Technical Consultation on Birth Spacing

In addition to the two recommendations in the WHO policy brief, ESD operational messages include a 3rd message for adolescents – “for adolescents, the recommended time period for the first pregnancy is at least 18 years of age or older.”

Within the context of informed choice about spacing or limiting future pregnancies, *for those who choose to space*, the above recommendations in the 2006 WHO Policy Brief, in conjunction with the preamble should be incorporated into counseling and behavior change communication messages.

Birth Spacing Trends and Demand and Unmet Need for Spacing

High percentages of mothers and newborns are at risk of adverse health outcomes due to too closely spaced pregnancies

13% of pregnancies to MWRA occur in less than the recommended interval. For mothers 15 to 19 years old, 33% occur in less than a two year interval. International data show that there is a 265% increased risk of a child death associated with births of less than 24 month intervals and a 300% increased risk of a neonatal, post-neonatal or infant death associated with births of less than 18 month intervals.


Indonesia is making progress in addressing its birth spacing issues

Between 1987 and 2003, the portion of births occurring after an interval that less than 24 months after a previous birth, declined by about 48%.

In Indonesia younger women have unmet need for family planning

Younger women face unnecessary health risks on both the *timing* and *spacing* of pregnancies. The rate of adolescent pregnancies has remained basically unchanged and 33% of births to 15-19 year olds are spaced less than 2 years apart.

According to the 2002/03 DHS, 57% of women of any reproductive age in Indonesia use a modern contraceptive method. Unmet family planning need for spacing among the total population of married women of any reproductive age is low (4%). In the younger age cohorts (15-29 years), the unmet need for spacing (at about 7%) is almost twice as high as that found in the entire reproductive age population. There is even demand among zero-parity women; for example, 27% of the 15-19 year olds with spacing demand in Indonesia have not yet had a child (Jansen 2005).



Spacing Patterns and Programmatic Implications

A major reproductive health issue

All available data and information indicate that the higher risks of short birth intervals and early pregnancies represent a major reproductive health issue in Indonesia. The problem of short birth intervals is even more pronounced in younger women, among whom the highest risks from very short birth intervals are more common. Although contraceptive prevalence has increased from 1987, it is likely that the majority of existing demand for spacing still goes unmet and the high frequency of short birth intervals demonstrates how much further family planning services must progress to better address birth spacing needs.

Strategies are needed

To reduce the current number of pregnancies that occur less than recommended intervals, couples will need easier access to spacing services that are responsive to their circumstances. Since most of the births occurring among women 15-19 years old are first births, the main issue for this age cohort is the timing of the first pregnancy. Communication, counseling and services for adolescents should focus on the health risks associated with the timing of a first pregnancy and birth.

Program research findings to date

Operations and survey research indicate that when women and their families are advised of the health and quality of life benefits of pregnancy delay and spacing, use of family planning increases substantially. One study conducted in India found that use of contraception for delaying the first child increased from 5% to 20% over 4 years in the intervention area, and from 4% to 8 % in the control area. The same study found that use of contraception for spacing the second child increased from 14 to 33% in the intervention area and from 10 to 20% in the control area. A project in Egypt, implemented from 2003-2005, which improved the quality of services, and included HTSP counseling and community outreach among influential leaders, was found to have increased the contraceptive prevalence rate from 50 to 80% among all married women of reproductive age and from 38 to 73 % among young, low-parity women. Similar studies are underway in Egypt, India, Yemen, Nepal, and Bangladesh.

Implications for Program Design

These statistics inform the following recommendations:

- **Present HTSP data to decision-makers**, to advance understanding of the role of pregnancy timing and spacing in the health of mothers and newborns in Indonesia.
- **Implement HTSP behavior change communication** and counseling interventions as an integral risk prevention strategy in all family planning, child and maternal health communications and client counseling protocols.
- Ensure that the two 2006 **WHO pregnancy spacing recommendations**, as well as information on the specific health benefits associated with the healthy timing and spacing of pregnancy are included in all communications and protocols.
- **Develop or strengthen** pregnancy delay or spacing services and communication activities for young (15-24 years) clients.
- **Strengthen service-delivery** and expand service outreach options **for adolescents who are married or un-married**, including those who have not yet had children.
- To achieve a more balanced method mix, **help families understand** that long-acting and intermediate methods (IUDs and injectables) are safe, and can effectively help them achieve their spacing preferences.
- Use this HTSP Profile in advocacy with national policymakers and district health teams, to **advocate for increased resources for HTSP interventions**.

Definitions

Unmet need for spacing: The percentage of currently married women who want to wait before having their next birth and are not using any method of family planning. Included with the unmet need for spacing are pregnant women whose pregnancy was mistimed; amenorrheic women whose last birth was mistimed; and, fecund women who are neither pregnant nor amenorrheic, not using any method of family planning, and who want to wait two or more years for their next birth. Unmet need for spacing also includes fecund women who are not using any contraception and are unsure whether they want to have another child or who want another child but are unsure when to have the next birth.

Demand for family planning for spacing: The sum of current contraceptive prevalence for spacing (including currently pregnant or amenorrheic women whose pregnancy or last birth was the result of a contraceptive failure) and the unmet need for spacing.

References

Conde-Agudelo, A. and J. M. Belizán

2000 Maternal Morbidity and Mortality Associated with Interpregnancy Interval: a Cross Sectional Study. *British Medical Journal*, 321:1255-1259 (18 November).

Conde-Agudelo, A., J. Belizán, R. Breman, S. Brockman and A. Rosas-Bermudez

2005 Effect of the Interpregnancy Interval After an Abortion on Maternal and Perinatal Health in Latin America. *International Journal of Obstetrics and Gynecology*, Vol. 89, Supplement 1 (April), S34-S40.

Conde-Agudelo, A., A. Rosas-Bermúdez and A. C. Kafury-Goeta

2006 Birth Spacing and Risk of Adverse Perinatal Outcomes: A Meta-analysis. *Journal of the American Medical Association*, 295:1809-1823.

Da Vanzo, J., A. Razzaque, M. Rahman, L. Hale, K. Ahmed, M. A. Khan, G. Mustafa and K. Gauzia

2004 The Effects of Birth Spacing on Infant and Child Mortality, Pregnancy Outcomes, and Maternal Morbidity and Mortality in Matlab, Bangladesh. Rand Corporation: Rand Labor and Population Working Paper Series, WR-198 (October).

Demographic and Health Survey

1989 Indonesia National Contraceptive Prevalence Survey 1987. Columbia, Maryland: Central Bureau of Statistics; National Family Planning Coordinating Board; Institute for Resource Development/Westinghouse.

1992 Indonesia Demographic and Health Survey 1991. Columbia, Maryland: Central Bureau of Statistics; National Family Planning Coordinating Board; Ministry of Health; Macro International, Inc.

1995 Indonesia Demographic and Health Survey 1994. Calverton, Maryland: Central Bureau of Statistics; State Ministry of Population/National Family Planning Coordinating Board; Ministry of Health; Macro International, Inc.

1998 Indonesia Demographic and Health Survey 1997. Calverton, Maryland: Central Bureau of Statistics; State Ministry of Population/National Family Planning Coordinating Board; Ministry of Health; Macro International, Inc.

2003 Indonesia Demographic and Health Survey 2002-2003. Calverton Maryland: Statistics Indonesia; National Family Planning Coordinating Board; Ministry of Health; ORC Macro.

Jansen, W.

2005 Existing Demand for Birth Spacing in Developing Countries: Perspectives from Household Survey

Data. *International Journal of Obstetrics and Gynecology*, Vol. 89, Supplement 1 (April), S50-S60.

Razzaque, A., J. Da Vanzo, M. Rahman, K. Gausia, L. Hale, M. A. Khan, A.H.M.G. Mustafa
2005 Pregnancy Spacing and Maternal Morbidity in Matlab, Bangladesh. *International Journal of Obstetrics and Gynecology*, Vol. 89, Supplement 1 (April), S41-S49.

Rutstein, S.

2005 Effects of Preceding Birth Intervals on Neonatal, Infant and Under-Five Years Mortality and Nutritional Status in Developing Countries: Evidence from the Demographic and Health Surveys. *International Journal of Obstetrics and Gynecology*, Vol. 89, Supplement 1 (April), S7-S24.

UNICEF

2006 State of the World's Children 2007: The Double Divide of Gender Equality. New York: United Nations Children's Fund.

World Health Organization

2006 Report of a WHO Technical Consultation on Birth Spacing. World Health Organization, Geneva, Switzerland. Available at:
<http://www.who.int/makingpregnancysafer/publications/policybriefbirthspacing.pdf>.

World Health Organization

2006 Policy brief on Birth Spacing – Report from a World Health Organization Technical Consultation. *WHO Department of Reproductive Health and Research and Department of Making Pregnancy Safer*.

Acknowledgement

Prepared by William H. Jansen, Ph.D.

The Extending Service Delivery (ESD) Project, funded by the United States Agency for International Development (USAID) Bureau for Global Health, is designed to address unmet need for family planning (FP) and increase the use of reproductive health /family planning (RH/FP) services at the community level, especially among underserved populations, in order to improve health and socioeconomic development. To accomplish its mission, ESD strengthens global learning and application of best practices; increases access to community-level RH/FP services; and improves capacity for supporting and sustaining RH/FP services. ESD works closely with USAID missions to devise tailored strategies that meet the RH/FP service delivery needs of specific countries. A five-year Leader with Associates Cooperative Agreement, ESD is managed by Pathfinder International in partnership with IntraHealth International, Management Sciences for Health, and Meridian Group International, Inc. Additional technical assistance is provided by Adventist Development and Relief Agency International, the Georgetown University Institute for Reproductive Health, and Save the Children.

Contact information:

Extending Service Delivery Project
1201 Connecticut Avenue, NW, Suite 700
Washington, D.C. 20036
Tel. 202-775-1977
Fax. 202-775-1988
esdmail@esdproj.org

This publication was made possible through support provided by the Office of Population and Reproductive Health, Bureau for Global Health, U.S. Agency for International Development, under the terms of Award No. GPO-A-00-05-00027-00. The opinions expressed herein are those of the author(s) and do not necessarily reflect the views of the U.S. Agency for International Development.