


INNOVATIONS IN TROPICAL REFORESTATION

*Casuarinas:  
Nitrogen-Fixing Trees  
for Adverse Sites*



The cover art depicting a representative casuarina branch, cane, and seed is by Priscilla Fawcett. It is reprinted from P. B. Tomlinson's *The Biology of Trees Native to Tropical Florida*, courtesy of the author.



*National Academy Press*

The National Academy Press was created by the National Academy of Sciences to publish the reports issued by the Academy and by the National Academy of Engineering, the Institute of Medicine, and the National Research Council, all operating under the charter granted to the National Academy of Sciences by the Congress of the United States.



INNOVATIONS IN TROPICAL REFORESTATION

*Casuarinas:  
Nitrogen-Fixing Trees  
for Adverse Sites*

*(Panel on Casuarina)* Report of an Ad Hoc Panel of the  
Advisory Committee on Technology Innovation  
Board on Science and Technology for International Development  
Office of International Affairs  
National Research Council

NATIONAL ACADEMY PRESS  
Washington, D.C. 1984

**NOTICE:** The project that is the subject of this report was approved by the Governing Board of the National Research Council, whose members are drawn from the Councils of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine. The members of the committee responsible for the report were chosen for their special competences and with regard for appropriate balance.

This report has been reviewed by a group other than the authors according to the procedures approved by a Report Review Committee consisting of members of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine.

The National Research Council was established by the National Academy of Sciences in 1916 to associate the broad community of science and technology with the Academy's purposes of furthering knowledge and of advising the federal government. The Council operates in accordance with general policies determined by the Academy under the authority of its congressional charter of 1863, which establishes the Academy as a private, nonprofit, self-governing membership corporation. The Council has become the principal operating agency of both the National Academy of Sciences and the National Academy of Engineering in the conduct of their services to the government, the public, and the scientific and engineering communities. It is administered jointly by both Academies and the Institute of Medicine. The National Academy of Engineering and the Institute of Medicine were established in 1964 and 1970, respectively, under the charter of the National Academy of Sciences.

The Board on Science and Technology for International Development (BOSTID) of the Office of International Affairs addresses a range of issues arising from the ways in which science and technology in developing countries can stimulate and complement the complex processes of social and economic development. It oversees a broad program of bilateral workshops with scientific organizations in developing countries and conducts special studies. BOSTID's Advisory Committee on Technology Innovation publishes topical reviews of technical processes and biological resources of potential importance to developing countries.

This report has been prepared by an ad hoc panel of the Advisory Committee on Technology Innovation, Board on Science and Technology for International Development, Office of International Affairs, National Research Council. Program costs for the study were provided by the Office of the Science Advisor, Agency for International Development, under Grant No. DAN/5538-G-SS-1023-00.



## STUDY PARTICIPANTS

### Australia

- MATTHEW ALEXANDRA, Brunswick, Victoria  
BRYAN A. BARLOW, Herbarium Australiense, Division of Plant Industry, Commonwealth Scientific and Industrial Research Organization (CSIRO), Canberra, Australian Capital Territory (A.C.T.)  
DOUGLAS J. BOLAND, Division of Forest Research, CSIRO, Canberra, A.C.T.  
T. H. BOOTH, Division of Water and Land Resources, CSIRO, Canberra, A.C.T.  
C. J. BOROUGH, Division of Forest Research, CSIRO, Canberra, A.C.T.  
GLYNN D. BOWEN, Division of Soils, CSIRO, Glen Osmond, South Australia  
MICHAEL CHUK, Forestry Unit, Conservation Commission of the Northern Territory, Alice Springs, Northern Territory  
PETER D. COYNE, Australian National Parks and Wildlife Service, Norfolk Island, New South Wales  
PETER M. DAVIDSON, Airport Engineering Services Branch, Department of Transport, Canberra, A.C.T.  
MAXWELL F. C. DAY, Deakin, A.C.T.  
JOHN C. DORAN, Division of Forest Research, CSIRO, Canberra, A.C.T.  
A. FLEMING, Department of Forestry, Australian National University, Canberra, A.C.T.  
A. H. GIBSON, Division of Plant Industry, CSIRO, Canberra, A.C.T.  
NORMAN HALL, Division of Forest Research, CSIRO, Canberra, A.C.T. (retired)  
HEATHER HOWARD, Division of Forest Research, CSIRO, Canberra, A.C.T.  
R. DOUGLAS JOHNSTON, Division of Forest Research, CSIRO, Canberra, A.C.T.  
A. C. LAWRIE, Department of Botany, Monash University, Clayton, Victoria  
JOHN LEIGH, Division of Plant Industry, CSIRO, Canberra, A.C.T.  
STEPHEN J. MIDGLEY, Division of Forest Research, CSIRO, Canberra, A.C.T.  
HENRY A. NIX, Division of Water and Land Resources, CSIRO, Canberra, A.C.T.  
PAUL REDDELL, Division of Soils, CSIRO, Glen Osmond, South Australia  
J. REDMOND, Australian Development Assistance Bureau, Department of Foreign Affairs, Canberra, A.C.T.  
IAN RILEY, School of Environmental and Life Sciences, Murdoch University, Murdoch, Western Australia  
JOHN SAUNDERS, Division of Water and Land Resources, CSIRO, Canberra, A.C.T.  
KENNETH R. SHEPHERD, Department of Forestry, Australian National University, Canberra, A.C.T.  
JOHN W. TURNBULL, Division of Forest Research, CSIRO, Canberra, A.C.T.  
K. B. WALSH, Department of Botany, University of Queensland, St. Lucia, Queensland

### Egypt

- M. H. EL-LAKANY, Forestry and Wood Technology Department, Faculty of Agriculture, Alexandria University, Alexandria

**India**

S. KONDAS, Tamil Nadu Forest Plantation Corporation Ltd., Tiruchirapalli,  
Tamil Nadu

J. S. P. YADAV, Central Soil Salinity Research Institute, Karnal, Haryana

**Papua New Guinea**

ANDREW AATAIA, Provincial Forest Officer, Goroka, Eastern Highlands Province

K. THIAGALINGAM, Department of Agriculture, University of Papua New  
Guinea, Port Moresby

**Philippines**

SATURNINA C. HALOS, Forest Research Institute, College, Laguna

**Senegal**

Y. DOMMERGUES, Office de la Recherche Scientifique et Technique Outre-Mer,  
Dakar

**Sri Lanka**

K. VIVEKANANDAN, Forestry Department, Colombo

**Thailand**

PRAVIT CHITTACHUMNONK, Silviculture Research Sub-Division, Royal Forests  
Department, Bangkok

**USA**

EDWARD S. AYENSU, Office of Biological Conservation, Smithsonian Institution,  
Washington, D.C. (*Chairman, National Research Council Casuarina Panel*)

H. W. EBSENSHADE, International Tree Crops Institute, Winters, California

THOMAS F. GEARY, USDA Forest Service, Southeastern Forest Experimental  
Station, Lehigh Acres, Florida

DAVID NORTHINGTON, Texas Tech University, Lubbock, Texas

JOHN G. TORREY, Maria Moors Cabot Foundation for Botanical Research,  
Harvard University, Petersham, Massachusetts

**Zimbabwe**

M. CHIHAMBAKWE, Mafungabusi Forest, Gokwe

L. J. MULLIN, Forest Research Centre, Harare

\* \* \*

NOEL D. VIETMEYER, Professional Associate, Board on Science and Technology  
for International Development, *Casuarina Study Director*

**National Research Council Staff**

F. R. RUSKIN, *BOSTID Editor*

MARY JANE ENGQUIST, *Staff Associate*

CONSTANCE REGES, *Administrative Secretary*

# Preface

This book highlights eighteen species of casuarina, a group of underexploited Australasian trees that could have exceptional potential in reforesting difficult terrain in many parts of the world. The idea for the study arose at a 1980 meeting of the Commission for the Application of Science to Agriculture, Forestry, and Aquaculture—a commission of the International Council of Scientific Unions—when Max Day was asked to propose a project in forestry for Third World countries, emphasizing trees for firewood, soil protection, and fodder production.

Dr. Day's suggestion for a joint study of casuarinas interested the U.S. National Research Council (NRC) because in 1980 it had briefly mentioned the promise of casuarinas in its report, *Firewood Crops: Shrub and Tree Species for Energy Production*.

It is not our purpose here to recommend casuarinas over all other possible reforestation species. No single species or group of species can provide the answer to the vast problem of tropical deforestation. Instead, we want to encourage broad consideration for increased planting of casuarinas.

Whenever governments, researchers, corporations, or private landowners are attempting to replenish their tropical lands with trees, they first should carefully assess native species; then the best-known fast-growing species such as pines, eucalypts, and gmelina; and finally, the lesser-known exotics such as leucaena, casuarinas, acacias, and calliandra. (These lesser-known species and others are described in companion reports listed below.) When all such species are included in trials, the best combination for a given site will become apparent, misplaced enthusiasm will be avoided, and some of today's obscure species, such as most of these casuarinas, may become major resources.

This study was done in cooperation with the Commonwealth Scientific and Industrial Research Organization (CSIRO) of Australia, which convened a meeting of casuarina experts in Canberra, Australia,

in August 1981.\* Some 43 delegates attended from 10 countries. After the conference, the study participants stayed for two days of discussion and the drafting of the statements that led to this report.

The participants are grateful to the organizations that made this study possible. CSIRO hosted the conference; the Australian Development Assistance Bureau sponsored a participant from Papua New Guinea; the International Development Research Centre (Ottawa, Canada) provided travel support for the participants from India, Thailand, Egypt, and Papua New Guinea; the Government of Zimbabwe supported the two participants from that country; and the U.S. Agency for International Development supported the participants from the United States, Philippines, Senegal, and Norfolk Island.

The Advisory Committee on Technology Innovation of the NRC Board on Science and Technology for International Development (see page 113) is assessing scientific and technological advances that might prove especially applicable to problems of developing countries.

This report is one of an ACTI series *Innovations in Tropical Reforestation*. Other titles include:

- *Leucaena: Promising Forage and Tree Crop for the Tropics* (1977)
- *Firewood Crops: Shrub and Tree Species for Energy Production Volume I* (1980)
- *Sowing Forests from the Air* (1981)
- *Mangium and Other Fast-Growing Acacias for the Humid Tropics* (1983)
- *Calliandra: A Versatile Small Tree for the Humid Tropics* (1983)
- *Firewood Crops: Shrub and Tree Species for Energy Production Volume II* (1983)

Information on promising fast-growing, nitrogen-fixing trees is also contained in *Tropical Legumes: Resources for the Future* (1979). An updated edition of the 1977 leucaena book is in preparation.

These activities are supported largely by the U.S. Agency for International Development (AID). This study was sponsored by AID's Office of the Science Advisor, which also made possible the free distribution of this report in developing countries and to members of the development community.

How to cite this report:

National Research Council. 1984. *Casuarinas: Nitrogen-Fixing Trees for Adverse Sites*. National Academy Press, Washington, D.C.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Experiences with Casuarinas</b>	<b>16</b>
<b>3</b>	<b>The Plants</b>	<b>33</b>
<b>4</b>	<b>Management</b>	<b>40</b>
<b>5</b>	<b>Uses</b>	<b>46</b>
<b>6</b>	<b>Best-Known Species</b>	<b>55</b>
<b>7</b>	<b>Other Promising Species</b>	<b>67</b>
<b>8</b>	<b>Recommendations and Research Needs</b>	<b>91</b>

## Appendixes


<b>A</b>	<b>Selected Readings</b>	<b>99</b>
<b>B</b>	<b>Research Contacts</b>	<b>106</b>
<b>C</b>	<b>Index of Species</b>	<b>112</b>


	<b>Advisory Committee on Technology Innovation</b>	<b>113</b>
	<b>Board on Science and Technology for</b>	
	<b>International Development (BOSTID)</b>	<b>113</b>
	<b>BOSTID Publications</b>	<b>115</b>

The drawings on pages 66 and 90 are by M. Flackton, reprinted from *Forest Flora of New South Wales* by Joseph Henry Maiden.

INNOVATIONS IN TROPICAL REFORESTATION

*Casuarinas:  
Nitrogen-Fixing Trees  
for Adverse Sites*







# 1

## Introduction

### TROPICAL DEFORESTATION

We are, says a recent U.S. Government report, “in transition from global forest wealth to global forest poverty.” Europe and North America began a century ago to balance cutting and regrowth carefully, but in the great equatorial greenbelt—at once among the richest and most fragile of resources—poverty and lack of effective controls have combined to unleash a voracious appetite for trees. Two-thirds of Latin America’s original forest are gone or seriously depleted. Half of Africa’s woodlands have disappeared. Thailand has lost one-quarter of its forest in the last 10 years, and the Philippines, one-seventh in the last 5. Spreading patches of the Amazon, Central Africa, and the Himalayan foothills have taken on the barren look of North Africa and the Middle East, both of which once boasted verdant stands of trees.

The issue goes far beyond millions of acres of lost trees. Land haphazardly cleared for crops or grazing—the single leading cause of deforestation—typically replaces rich jungle with virtually useless shrubs or sharp-edged grasses.

For one-half of the world’s population, the principal fuel is firewood, and the hills are stripped for it. More than 90 percent of the wood cut in Africa (nearly 2 million hectares a year) is burned as fuel. In East Africa, as trees thin out, women spend up to six days a week collecting wood simply to keep their families in cooking fuel; without it, food supplies are useless. In West African cities, where it often costs more to heat a pot than fill it, organized syndicates use trucks, donkeys, and porters to spread the destruction far into the bush.

Ultimately, the only sure way to stop deforestation is to minimize its causes—lack of fuel, fodder, and farmland for the third of humanity that lives on the edge of starvation. For most sites, this means growing



Casuarinas are broad-leafed plants (angiosperms) with foliage like conifers (gymnosperms). Casuarina foliage, with its "needles" and cones, is often mistaken for some form of pine. However, casuarinas are easily distinguished because their needles are segmented and can be pulled apart into small pieces. Picture shows *Casuarina equisetifolia* subspecies *incana*, photographed at Green Island on the Queensland coast in northern Australia. (D. J. Boland)

more trees. And to make any impact, it will take hardy, adaptable, and vigorous trees that can colonize the harsh, eroded land and begin its return to productivity. This is where casuarinas may find a role.

## CASUARINAS

Casuarinas\* comprise a group of about 80 species of shrubs and trees that are primarily native to the southern hemisphere, mostly to

\*The generic name was invented by Rumphius in the 1650s and applied by Linnaeus. It alludes to the drooping, filamentous branchlets which resemble the plumage of the cassowary bird (*Casuaris* species). The Australian pronunciation "caz-yew-a-reena" seems preferable to "caz-yew-a-rye-na," although both are used. These trees are commonly known as sheoaks (Australia), agoho (Philippines), Australian pine (Caribbean), and by many other local common names. In 1982 taxonomist L.A.S. Johnson proposed splitting the family Casuarinaceae into four genera: *Casuarina*, *Allocasuarina*, *Gynosperma*, and a yet unnamed genus. (See Johnson, 1982.) In this report we have decided to retain the traditional nomenclature. Johnson's proposed names are, however, included with the appropriate species in chapters 6 and 7.

Australia, where they occur in tropical, subtropical, and temperate coastal regions as well as in the arid inland. A few species are native to Indo-Pacific areas from Peninsular Malaysia to Polynesia.

One species, *Casuarina equisetifolia*, has been used widely as a seaside ornamental tree and to stabilize coastal sand dunes because it thrives in sand and saline conditions. Two others, *Casuarina cunninghamiana* and *Casuarina glauca*, have also been planted outside their native habitats. But there are at least 15 more casuarinas with apparent promise, particularly as firewood crops. They deserve wider recognition.

## APPEARANCE

Some casuarinas are small diffuse or spreading shrubs only a few centimeters high, while others are tall, erect, and graceful evergreens with open, feathery crowns. Some attain heights of 50 m and diameters



In addition to their foliage, casuarina trees often have the form and shape of conifers, further compounding the confusion with pines. Picture shows hybrid of *Casuarina equisetifolia* and *Casuarina junghuhniana*, Bangkok, Thailand. (Pravit Chittachumnonk)

of 1 m. \* Many, however, are only 15–25 m tall. Viewed from a distance, their small cones and long, drooping branchlets, which look like pine needles, make them look like conifers.

## USES

The casuarinas produce high-quality fuelwood that burns with great heat and has been called the best firewood in the world. It splits easily, burns readily even when green, has low ash content, and makes excellent charcoal. It is useful for both domestic and industrial fuel and was a prime fuel for Australian bakeries until they began to be electrified, only 50 years ago.

Casuarina wood was one of Australia's first exports: British ships that took convicts to the colony of New South Wales often loaded with casuarina and red cedar for the return voyage. It is a very dense wood, but because it splits and warps on drying it is difficult to use for lumber or furniture. However, in the form of poles and beams it is used for construction because it is very strong. Small items such as roofing shingles and turnery products are also made from it. The wood can be pulped for paper, but so far the cost of grinding up this hard, dense wood has kept it from becoming an industrial pulpwood. Nevertheless, in Egypt, several particle board factories rely on casuarina, almost the only trees to grow in Cairo's arid environs.

Casuarina trees, taken collectively, have many other uses. Among the various species there are casuarinas capable of stabilizing shifting sand dunes, stabilizing eroding hillslopes, and reclaiming marshy soils that are periodically inundated by fresh, or even brackish, water. Many make useful shade trees, windbreaks, and shelterbelts; the shape and form of their foliage makes them excellent for these purposes, and most species are fairly wind firm and often retain branches to ground level. They also withstand clipping and, in Hawaii, for example, they are frequently trimmed as hedges and yard trees. Casuarinas are poor forage sources, but in times of extreme drought Australian farmers use them as emergency animal feed. In Madagascar the bark, reported to contain 6–18 percent tannin, has been used extensively for tanning leather.

## ADVANTAGES

On suitable sites casuarinas may grow as rapidly as the fastest-growing tropical trees. In parts of the Philippines *Casuarina equisetifolia* has been known to outgrow *Leucaena leucocephala* and *Gmelina arborea*. On granitic uplands in Puerto Rico *Casuarina equisetifolia*

\*All diameters reported in this book were measured at "breast height" (1.3 m; 4.5 ft).



*Casuarinas* are noted for fast growth, especially on favorable sites. This *Casuarina equisetifolia* is 26 months old, grown at Port-au-Prince, Haiti, in a trial of provenances gathered throughout the Caribbean. (A. van Wingerden)

grows faster than *Pinus caribaea*. In India *Casuarina equisetifolia* saplings have been measured as growing 3 m a year.

Most casuarinas are easy to propagate. They usually set seeds in abundance, and for the majority of species, the seeds store well and germinate readily. Probably all casuarinas can be propagated vegetatively. A sterile hybrid of *Casuarina equisetifolia* and *C. junghuhniana* is propagated almost entirely by cuttings in Thailand and India, and it seems likely that other species can also be propagated this way, but



Casuarinas have remarkable ability to grow on harsh sites and toxic soils. This page, top: *Casuarina decaisneana* in the Simpson desert in the heart of Australia where few, if any, other trees can tolerate the extreme heat, searing drought, and frequent ground fires. (M. Chuk) Bottom: *Casuarina deplancheana* thriving in soils so rich in aluminum and iron that they are toxic to most plants, New Caledonia. (Y. Dommergues) Facing page, top: One-year-old *Casuarina equisetifolia* planted in sterile sand dunes on the coast of Senegal. (Y. Dommergues) Bottom: *Casuarina equisetifolia* rehabilitating quarry site and limestone spoil from cement factory, Mombasa, Kenya. (R. D. Haller)





Casuarinas are renowned for an ability to thrive in areas exposed to salt spray, waterlogging, and even inundation by the sea. This page, top: *Casuarina equisetifolia* growing down to the high water mark, Kona Coast, Hawaii, USA. (C. S. Hodges) Bottom: *Casuarina cunninghamiana* on the banks of the Cotter River, near Canberra, Australia. (D. J. Boland) Facing page, top: *Casuarina equisetifolia* behind the beach, south coast of Thailand. (Pravit Chittachumnonk) Bottom: *Casuarina equisetifolia*, close to the high tide mark, west coast of Florida, USA. (C. J. Anderson)





most have not been tried. Cuttings have the advantage of being a rapid, sure method of perpetuating the traits of superior trees and also of allowing trees of a single sex to be planted so as to avoid weedy regeneration through the spread of natural seedlings.

With their hardy nature, casuarinas need little care. They are fairly resistant to pests and suffer few major diseases. Most species also tolerate extreme heat. *Casuarina decaisneana*, for example, grows in parts of central Australia where summer temperatures may reach 47°C.

In most cases, young casuarina seedlings compete aggressively with weeds. In the Philippines *Casuarina equisetifolia* is recognized as one of the best trees for planting in sites covered by *Imperata* grass, a tenacious weed that renders large areas of the tropics useless for agriculture. In the Highlands of Papua New Guinea *Casuarina oligodon* is also planted in areas infested with *Imperata*. This ability of casuarina trees to withstand competition from grasses in their early years could give them an advantage over eucalypts, particularly in Africa and Asia, where grass competition has caused some extensive failures of eucalypt plantings.


Casuarinas seem good candidates for planting in barren or polluted areas. They have the basic physiology for survival in diverse situations, and they often serve to colonize and revegetate poor sites. Many species will grow on soils of low fertility, some thriving on light soils, others on heavy soils. Along India's west coast as well as in Madagascar, for instance, casuarina is found on laterite—the infertile, red, iron-rich, leached soil widely found in the tropics. In Uruguay, Brazil, Senegal, and elsewhere, *Casuarina equisetifolia* thrives in very deep sandy soils. In Argentina *Casuarina cunninghamiana* has been observed colonizing bare limestone. At a lime-grinding factory southwest of Cairo, Egypt, casuarinas grow right in the lime, where no other tree survives. In Kenya *Casuarina equisetifolia* grows well around a cement works, and in Malaysia it colonizes sterile tin tailings. In Hawaii *Casuarina equisetifolia* is growing well in sterile pumice.

Some casuarinas are salt tolerant. Near Bangkok, Thailand, there are commercial plantings of the hybrid between *Casuarina equisetifolia* and *Casuarina junghuhniana* in salt-marsh areas sometimes inundated with saline water. *Casuarina obesa* occupies saline sites in the wheat belt of Western Australia. *Casuarina glauca* grows naturally on estuarine lowlands of New South Wales and Queensland that are flooded with brackish tidal water. And both *Casuarina equisetifolia* and *Casuarina glauca* thrive in dunes at or near the seaside—often directly in the path of ocean spray. *Casuarina equisetifolia* even grows on the shores of Hilo Bay, Hawaii, on tidal rocks often submerged in salt water.

## SYMBIOSES

The outstanding ability of various casuarinas to grow vigorously on poor soils is due partly to their unusual symbiosis with an actinomycete, *Frankia*, that enables them to use nitrogen directly from the atmosphere. Casuarinas belong to a group of more than 170 species of actinomycete-nodulated woody plants that fix atmospheric nitrogen. In moist soils at certain sites in Australia and elsewhere, the roots of one-third or more of the trees of a given *Casuarina* species are well nodulated by this nitrogen-fixing, bacteria-like microorganism.

*Frankia* infects root hairs of fine casuarina roots, forming nodules (swellings). These nodules are woody and perennial and can form large masses in the root system. The nitrogen that the actinomycete in the nodules fixes enables the casuarinas to grow well in soils that otherwise would be too deficient in nitrogen to sustain plant growth. The amounts of nitrogen they fix is roughly comparable to the amounts fixed by




Nodules on 1-year-old *Casuarina equisetifolia*. Casuarinas form large root nodules. These house symbiotic microorganisms that provide the plant with usable nitrogenous compounds. Although casuarinas can grow without the helpful microbes, the development of nitrogen-fixing nodules fosters faster growth, enriches soil with nitrogen, benefits growing plants, and helps casuarina trees become established on previously unproductive lands. (Y. Dommergues)


legumes with their *Rhizobium* symbionts. In sand dunes in the Cap Vert peninsula of Senegal, nitrogen in the soil around casuarinas has increased annually at rates of about 60 kg per hectare. This fixed nitrogen is added to the soil through decay of dead rootlets and leafy litter.

Casuarina roots also have symbiotic relationships with three other types of microorganisms. Two forms of mycorrhizal fungi enter the root and facilitate the uptake of minerals, notably phosphorus, and some trace elements. Furthermore, a range of unidentified microorganisms interact with the plant roots to produce dense mats of "proteoid roots," the large surface area of which probably also helps to absorb phosphorus and other vital minerals. Experiments have shown that proteoid roots can double the growth of plants in soils that are very low in phosphorus.

With the combination of all four symbionts, it seems hard to imagine plants better suited than casuarinas to the reforestation of low-fertility lands.



Opened root nodule of *Casuarina glauca*. The microorganism in casuarina root nodules is a filamentous soil actinomycete, *Frankia*. Its ability to "fix" nitrogen from the air, thereby providing casuarinas with nitrogenous "fertilizer," is one reason for the trees' ability to survive on barren land. (Electron Microscope Centre, University of Queensland)



A limitation of some casuarinas is their tendency to spread by root suckers. The profuse root suckering of *Casuarina glauca*, for example, makes it a nuisance in some areas. Here the tree is colonizing a degraded site in Haiti, where the spreading is considered a blessing because little else can survive. (T. F. Geary)

## LIMITATIONS

Care must be taken when introducing a casuarina to a new area. Some species are so innately vigorous that they can get out of hand and could develop into major pests by “escaping” into natural areas. This is already happening in southern Florida (USA), where *Casuarina equisetifolia* and *Casuarina glauca* in many instances are becoming undesirable weeds. *Casuarina glauca* is also a pest in Hawaiian pastures, where it spreads by root suckering. Because of this, casuarinas are not recommended for planting in areas where their wood is not needed.

Along the edges of roads and farm fields, as well as in lawns, root suckers are a nuisance and have to be cut back continually. On the other hand, in many situations root suckering is an advantage—in fuelwood plantations, for example, the cut trees rapidly regenerate from root sprouts and do not have to be replanted. This is particularly true in eroded “blowouts” where the suckering species can quickly occupy the whole area.



Another limitation is that the wood from most *Casuarina* species is not suited to many lumber uses because it cracks and warps on drying. (M. H. El-Lakany)

Casuarinas have still other limitations. Because the branchlets are often rich in selenium, silica, and salt, the copious leaf litter under casuarinas may be toxic to nearby plants. On some sites the thick litter under the trees may contribute to acidification or salinization of the soil, and casuarinas, therefore, may not be good companion crops. In such cases, the effects are probably similar to those of pine trees.\*

Although most casuarinas are well adapted to survive fires (some by producing shoots from thick woody rootstocks, others by coppice shoots from buds in the stem), a few are fire sensitive. *Casuarina equisetifolia*, for example, is readily killed even by light fires.

Without their *Frankia* symbiont, casuarinas are often disappointing when planted as exotics.

\*Casuarinas do not seem to be universally bad neighbors. Indian scientists, for instance, have found that under casuarina stands infertile soil is enriched and there is an overall improvement of the site. (Information from S. Kondas.) And in Papua New Guinea casuarinas are widely used to improve soil. This also applies to most natural systems in Australia in which *Casuarina* species grow.

## CONCLUSION

Casuarinas occur in a wide range of habitats, from rainforest to desert, from seashore to high mountaintop, and from cool temperate regions to the hot humid tropics. They have modest site requirements and most are rapid-growing, trouble-free trees. Many may be salt tolerant, others are especially wind firm, and all seem adaptable to poor soils. Casuarina trees form root nodules in symbiotic association with the soil actinomycete *Frankia* that fixes atmospheric nitrogen. They also form symbioses with at least two different groups of fungi, which endow various species with the ability to extract phosphorus and other minerals from even the poorest soils.

Casuarinas can be used for many purposes, from amenity planting, land reclamation, shelterbelts, and dune stabilization to the production of shingles, particle board, tannin, timber and roundwood, and perhaps paper pulp. The wood makes outstanding fuel.

During the past century, most casuarinas have been largely neglected in comparison with many Australian *Eucalyptus* species that have become established forest resources in many of the world's warm regions. Now casuarinas warrant increased attention. One day they may complement eucalypts as a global wood resource, especially as they have the advantage of fixing nitrogen and seem to tolerate competition from weedy grasses better than eucalypts.

## 2

# Experiences with Casuarinas

This chapter briefly mentions how various casuarinas are grown and used in almost 20 different regions. Although Australia has had the most extensive involvement with the trees, its experience is dealt with under the species themselves in chapters 6 and 7.

## ASIA

### CHINA\*

A new great wall has appeared in China—a green shelterbelt along the southern coast covering an area of more than 1 million hectares, built mainly with *Casuarina equisetifolia*.

In 1949, immediately after the People's Republic of China was established, the government made efforts to stimulate forestry development. People were mobilized and large afforestation projects started. These included forests primarily established for protection purposes, such as shelterbelts to stop the encroachment of coastal and desert sand dunes and to diminish the force of strong winds.

Since 1954, vast plantings of *Casuarina equisetifolia* have been established along the coast fronting the South China Sea. Much of the coast there comprises bare dunes that formerly were constantly moving inland, destroying arable land. This belt of casuarina stretches for 3,000 km and varies from 0.5 to 5 km in width. Most of it is in the province of Guangdong, but it extends into Fujian province and the Guanxi autonomous region.

\*Information from J. W. Turnbull.





*Casuarina equisetifolia*, planted for fuelwood and sand-dune stabilization at Wu Yang People's Commune, Guangdong Province, China. With 44 km of coastline along the South China Sea, this commune suffered greatly from sand dunes invading agricultural land. Now, some 1,500 hectares of *Casuarina* shelterbelt have been planted along the beachfront. Fallen cones, small branches, and leaf litter are collected for fuel as often as once every 2 weeks (see rake marks). These 13-year-old trees average more than 13 cm in diameter and 17 m in height. They are harvested for fuel and wood, but only in small clear-cut patches so as to retain sand stabilization and shelter for the farm fields and villages behind the dunes. (T. M. Catterson)

The plantations grow in a mild, subtropical climate between the latitudes of 20°N and 30°N.\* Strong northeasterly winds occur frequently, and typhoons in summer are common.

Forestry officials in the area promote the use of casuarina. New plantations are planted using seeds collected selectively from trees that are tall, straight, fast growing and disease free.

Seedlings can reach 3 m tall 1 year after planting, and trees in an average plantation are 7–8 m tall and 5–7 cm in diameter after 4 years. The rotation period ranges from 10 to 15 years, and the mean annual increment averages 4–5 m<sup>3</sup> per hectare.

The major benefits of this gigantic belt of *Casuarina equisetifolia* are the control of drifting sand, the sheltering of villages from coastal winds, and the production of wood for construction and fuel. The wood is used for house construction, especially for beams, as well as for boatbuilding and furniture. Fuel is a very important product for domestic use and for firing local brick kilns. In addition to branch wood and scrap wood that are used as fuel, each hectare of plantation annually provides up to 4 tons of litter and twigs that are continuously harvested for domestic fuel.

(In some areas the trees are attacked by a disease that causes the whole tree to wilt and die. This has been attributed to a bacterium, *Pseudomonas*, attacking the roots. *Casuarina glauca* is reportedly less susceptible.)

## MICRONESIA

*Casuarina equisetifolia* (gago) is a common native tree in Guam and most other high islands of Micronesia where it grows both along the coasts and in upland savannas. On Guam the spread of the trees in savannas is retarded by frequent fires, which kill them. It occurs naturally on both limestone and volcanic soils.

## PAPUA NEW GUINEA

Casuarinas are part of village life in the Highlands of Papua New Guinea. In the early 1950s the Department of Primary Industry (Office of Forests), together with various patrol officers, set up tree nurseries throughout the Highlands and people were taught the value of *Casuarina oligodon* and some other species. Since then the people have been growing casuarina, transforming the landscape from one predominantly of grasslands to an attractive and varied one with many trees.

\*Mean annual rainfall is about 1,500 mm, and there is a dry season from September to March. Mean annual temperature is 24°C, the absolute maximum is 37°C, and the area is frost free. The soils are loose sands with a pH of 5-7.



Eastern Highlands Province, Papua New Guinea. *Casuarina oligodon* is traditionally planted as a fallow crop and around villages. Soil samples collected after 5–10 years of this forest fallow show substantial increases in soil nitrogen and organic matter. (R. L. Parfitt)

In the early 1960s, as a small-scale research project, *Casuarina oligodon* was successfully underplanted beneath *Araucaria hunsteinii* and *Araucaria cunninghamii*. In recent years casuarinas have been used to revegetate copper mine spoil at Bougainville. And since before World War I, *Casuarina equisetifolia* has been an ornamental in Port Moresby, Wewak, Lae, Madang, Rabaul, and other coastal towns.

Casuarinas are also used as a cover crop for coffee. Coffee is a major cash crop in the Papua New Guinea Highlands, and *Casuarina oligodon* is a very useful shade tree because it lets light through to the crop and fixes nitrogen to improve the soil.

#### THAILAND

*Casuarina equisetifolia* is widespread in Thailand, and it has long been planted along coastlines to produce the poles used in building fishing traps as well as for fuelwood. A sterile hybrid between *Casuarina junghuhniana* and *Casuarina equisetifolia* was introduced to Thailand around 1900 as an ornamental tree. It soon received greater attention as a plantation species, owing to its fast growth, good stem form, and its adaptability to difficult site conditions. Both species are now used extensively to reclaim land abandoned after mining as well as for fuelwood production in village woodlots and for durable pilings for urban construction.



Rangsit, Thailand. Hybrid of *Casuarina equisetifolia* and *Casuarina junghuhniana* growing on heavy acid soils poorly suited to most other tree crops. Grown from cuttings, the trees are harvested after only 5 years for pilings, posts, and firewood. (Pravit Chittachumnonk)



Madras, India. *Casuarina equisetifolia* thrives at the seaside. In South India, it is grown on sterile sand dunes and is harvested for fuel and wood products on a 7–10 year rotation. (N. D. Vietmeyer)

## INDIA\*

*Casuarina equisetifolia* was introduced into Madras State in the 1860s to fuel steam locomotives. Villagers then spread it along the coasts of the Indian Peninsula, especially on the east coast. In due course, farmers brought it into the South Indian agricultural system by planting it around their homes and fields. Casuarinas now provide protection to exposed sand on the shores of South India and on the Andaman Islands.

In South India *Casuarina equisetifolia* is considered the best species for colonizing sand dunes. In certain parts it is used to check dunes near farmers' fields, fishermen's dwellings, roads, railways, and ports. It makes life less wearisome in the coastal sands, with their fierce winds, flying sand, and dust under the hot tropical sun.

This tree is now one of the main firewood species in peninsular India. There are some 39,000 hectares under cultivation in plantations. Normally harvested on a 7– to 15–year rotation, the species yields

\*Information from S. Kondas.

100–200 tons of fuelwood per hectare. The wood brings high prices. It is used for home cooking as well as for drying agricultural products such as cocoa, tea, copra, and tobacco. Today, *Casuarina equisetifolia* is being planted on a large scale in parts of Tamil Nadu, where the dry season is 5–6 months long and the soil is coastal sand, river alluvium, red loam, red gravelly soil, or hard laterite.

In the 1970s the *Casuarina junghuhniana* x *Casuarina equisetifolia* hybrid was brought to India from Thailand as fuel for the tea-drying industry. This hybrid is performing well. There is also an unidentified casuarina species growing sparsely as shade trees in tea gardens in the Nilgiri Hills in Madras State. This species has not spread much.

In India, when casuarinas are felled they are converted and marketed in at least four forms: stumps, thicker branches, finer branches with needles, and billets 1 m long. These products meet the various needs of both the rich and poor. The stumps are exceptionally good for making charcoal.

In certain parts of India well-to-do agriculturists plant a few hectares to casuarina when their daughters are young. A 5-year rotation provides sufficient dowry at the time of the marriage settlement.

## ISRAEL

*Casuarina equisetifolia* is used with much success for binding sand near the seashore. The casuarina plantations are widespread along roadsides and as windbreaks and small woodlots. Casuarina plantations in Israel were found to be mixed stands of *Casuarina cunninghamiana* and *Casuarina glauca*, with the former comprising 75 percent or more of the trees. Growth of *Casuarina glauca* is superior to that of *Casuarina cunninghamiana*.

In addition, 15 *Casuarina* species are being tested in 16 experimental plots.

## AFRICA

### EGYPT

*Casuarina* is the most important genus of trees in Egyptian forestry. Three species are grown: *Casuarina equisetifolia*, *Casuarina glauca*, and *Casuarina cunninghamiana*. A natural hybrid between the last two species has also been identified.

Along the coast, *Casuarina equisetifolia* is planted to protect houses against wind and salt. Inland, desert highways are protected by belts of *Casuarina glauca* and *Casuarina cunninghamiana*. And when farmers receive land from the government, the first thing they do is to



Egypt, southwest of Alexandria. A shelterbelt of *Casuarina cunninghamiana*, 12 years old, planted in a grid pattern to protect farmland from desert. Casuarinas, the most important genus of trees in Egypt, are widely planted for shade and shelter under harsh conditions. (International Development Research Centre)

plant casuarina shelterbelts. The casuarinas' popularity in Egypt can be judged from the fact that annual planting was 1 million seedlings in the mid-1970s, 4 million in 1980, and is projected at 10–15 million by 1990.

#### KENYA\*

*Casuarina equisetifolia* occurs along the Kenya Coast, often in sand dunes close to the sea. It has proved to be the best growing tree for reafforesting a formerly strip-mined limestone quarry of a cement factory near Mombasa. Approximately 40,000 trees were planted and vast numbers of seedlings are now colonizing the surrounding barren area. They are planted together with trees such as *Conocarpus lancifolius* and neem (*Azadirachta indica*). The estimated per hectare yield of wood is 120 tons and more in 5 years of growth. The charcoal and firewood produced from the casuarina is of superior quality. The straight stems are used for building poles and fence posts. (They must be treated to last if they are to be exposed to rain and high moisture.) The leaves are eaten by goats and eland. The humus under the trees rapidly increases as a result of the constant leaf fall. The dry leaves are eaten by the red-legged millipedes during the wet season, accelerating the process of humus production. In one part of the quarry 10 cm of humus has been produced in the last 10 years.

The trees are planted at 1 m x 1 m spacing, thinned out in the second and third year to 2.5 m x 2.5 m. The thinnings are sold for use in building mud-and-wattle houses.

*Casuarina equisetifolia* is also a popular amenity tree. About 10 years ago, establishment of the species started on plantation scale in Arabuka-Sokoke Forest Reserve. To date, there are about 350 hectares planted. The rate of growth is quite acceptable. In 9-year-old stands established at a spacing of 2.5 m x 2.5 m, the trees attained a mean height of 20 m and diameter of 12 cm. The wood is used mainly for building poles, fence posts, fuelwood, and charcoal.

*Casuarina cunninghamiana* and *Casuarina glauca* were introduced in 1908 and 1910, respectively. They are popular as ornamentals at an elevation of 1,600–1,800 m. Old trees of *Casuarina cunninghamiana* tend to be stag-headed.

*Casuarina junghuhniana*, introduced in 1956, has shown impressive performance. In a 26-year-old trial plot at an elevation of 2,100 m, trees have reached a mean dominant height of 25.5 m and diameter of 38.0 cm.

*Casuarina torulosa* was introduced in 1952. In a 30-year-old experimental plot on the same site as *Casuarina junghuhniana*, trees have reached a mean dominant height of 18.9 m and diameter of 29.6 cm.

\*Information from P. K. Arap Konuche and R. D. Haller.



## SENEGAL


Only one species, *Casuarina equisetifolia*, is successfully grown in coastal areas of Senegal. Introduced in 1925, it is now extensively used to stabilize coastal sand dunes between Dakar and St. Louis. These plantations now cover about 4,000 hectares and are being increased at the rate of 300–400 hectares per year.

The rainfall in the region is less than 500 mm annually. Brushwood, plastic fences, or fine nylon nets are first laid on the soil for temporary stabilization. Then, after rain has moistened the upper 40 cm of soil, the seedlings are planted. They receive no further attention. *Casuarina equisetifolia* is the only plant yet found that can survive this harsh treatment in the very poor soils and hostile climate.

## ZIMBABWE

*Casuarina cunninghamiana* was probably introduced into Zimbabwe during the early years of this century, and since then it has been widely used for street and roadside plantings and windbreaks. Over the years the species has been largely neglected, but foresters now look at it with new interest. Agroforestry projects using it are being started, and researchers hope that it can help relieve Zimbabwe's acute fuelwood shortage.

In 1964–1965 two small plots of *Casuarina junghuhniana* were established in a high-rainfall locale from a seed sample obtained from Kenya. The health and general performance of these plots has been good. In 1982 seed of *Casuarina decaisneana* was sown in trials of trees for the country's arid areas.



Dakar, Senegal. Along Senegal's northern coast, a belt of *Casuarina equisetifolia* has been planted in coastal sand dunes. This arid region has annual rainfall of less than 250 mm. The shifting dunes are temporarily stabilized with brush mats or fine nylon net before planting. (M. C. G. McD. Dow)



Mombasa, Kenya. Ten-year-old *Casuarina equisetifolia* on formerly strip-mined quarry floor. (R. D. Haller)

## THE AMERICAS

### ARGENTINA

Casuarinas, together with eucalypts and acacias, are common trees in Argentinean pampas and other dry areas of the country. To develop agriculture and livestock in these naturally treeless regions required the planting of trees for shade and to protect homesteads, animals, and crops from the strong, hot, and (in the north) salty winds. As a result, *Casuarina cunninghamiana* is very well known, although many people do not realize that it is not a pine.

In the Parana River delta, casuarinas are planted along the edges of rivers and creeks to protect the banks from erosion by waves. The



Fremantle, Haiti. *Casuarina glauca* on severely eroded hillside, planted by innovative Haitian farmer in loose rock using root suckers. (T. F. Geary)

intricate system of roots forms a strong wall against the water. The growth in these conditions is reported to be extraordinarily fast.

#### MEXICO

Casuarinas are now being planted around the huge saline Lake Texcoco in an effort to tame the dust storms that often overwhelm Mexico City.

#### HAITI\*

Casuarinas are not common in Haiti, but a small plantation of *Casuarina glauca* has grown well on a severely eroded hillside. The stand, which has nitrogen-fixing nodules and root suckers, is proliferating over the stony soil surface, and litter buildup is protecting against further erosion. Casuarinas appear to be good candidates for restoring Haiti's eroded mountain soils and for providing the fuelwood, charcoal, and poles so desperately needed in that country.

#### DOMINICAN REPUBLIC\*

In the dry zones of the Dominican Republic, and especially on alkaline soils, there are a few casuarina trees (thought to be *Casuarina*

\*Information from T. F. Geary.

*equisetifolia*) that have grown to a large size. A private company has made trial plantings of *Casuarina equisetifolia* on reclaimed strip-mined land in a humid zone and has found that it grows faster than *Eucalyptus camaldulensis*, *Pinus caribaea* var. *hondurensis*, and the native *Pinus occidentalis*.

*Casuarina equisetifolia* was a common street tree in the coastal city of Santo Domingo until hurricanes in 1979 blew the trees down. Trees 20–25 cm in diameter snapped off 3–5 m above ground, while larger trees were uprooted. Streets were clogged with debris from downed trees, and cleanup was costly. *Casuarina equisetifolia* is being replanted in some sections of the city, but it may not again become a common ornamental.

Because the Dominican Republic depends heavily on charcoal and wood for home cooking, and the native forests have been overexploited, casuarina plantations could become important to the fuelwood industry of the island.

#### PUERTO RICO

In Puerto Rico, *Casuarina equisetifolia* is the only casuarina commonly planted. It is found along the coast and, less commonly, in the



Bonao, Dominican Republic. One-year-old *Casuarina equisetifolia* on rehabilitated nickel strip mine. Soil is latosol, very deficient in nutrients. This casuarina displayed better height growth and required less weed control than eucalyptus, leucaena, and two pine species. (T. F. Geary)

lower mountain regions. It is recommended for beaches and windbreaks.\* Planting close to buildings is discouraged because a disease sometimes kills old trees and there is danger that they will blow over in hurricanes. Natural regeneration is rare in Puerto Rico because ants eat nearly all the seeds.

In screening trials of trees for commercial timber production on granitic uplands, *Casuarina equisetifolia* grew faster than *Pinus caribaea* var. *hondurensis* and most other fast-growing species tried. Only *Eucalyptus tereticornis* outpaced it.†

#### CALIFORNIA‡

Various casuarina species, particularly *Casuarina cunninghamiana* and *Casuarina stricta*, have been used for street trees and ornamentals in California. They have also been planted along public highways and, along one particularly windy stretch of road in the San Francisco Bay delta, have proved superior to pine as a windbreak. *Casuarina cunninghamiana* windbreaks are being tested on croplands in several regions of the state, including high desert and inland valley sites, as part of a state-funded windbreak-demonstration program. It is also among the tree species recommended by the state forestry department for planting for fuelwood. Growth rates as high as 2 m in 6 months have been recorded for *Casuarina cunninghamiana* planted adjacent to irrigated and fertilized cropland. A number of species are also being tested on salt-affected soils in the southern San Joaquin Valley, including *Casuarina decossata*, *Casuarina cristata*, *Casuarina glauca*, and *Casuarina helmsii*.

#### FLORIDA§

Australian pines, as the casuarinas are known locally, are common on the Florida coast, as well as along roads and property lines, on ditch and canal banks, and around buildings. The identities of the different species are uncertain, but it appears that all trees are *Casuarina equisetifolia*, *Casuarina glauca*, *Casuarina cunninghamiana*, or their various hybrids. Once these trees were highly regarded for landscaping, windbreaks, and shade because they grow well on acid and alkaline soils, sand dunes, calcareous rocky soils, muck, and many other soils. Now Floridians are not so enthusiastic.

Casuarina wood has no economic use in Florida, and several problems are associated with the trees. On fore-shore dunes the roots

\*Schubert, 1979.

†Geary and Briscoe, 1972.

‡Information from H. W. Esbenshade.

§Information from T. F. Geary and J. F. Morton.



Homestead, Florida. Windbreak of probable hybrid between *Casuarina equisetifolia* and *Casuarina glauca*. Very tall shelterbelts and roadside plantings of casuarinas are a common sight in South Florida; they dominate the skyline of Miami, for example. (T. F. Geary)

can prevent sea turtles from digging nests for their eggs, and sometimes turtles are trapped by these roots. The casuarinas have crowded out native vegetation in some parts of coastal southern Florida. Much of the pollen is airborne from December to April and is suspected of being the cause of respiratory complaints during these months. Giant specimens along city streets threaten adjacent dwellings in this hurricane-prone region. As a result, some efforts are being made to eradicate the tree from environmentally critical areas, and landscape plantings that are being removed in the process of highway widening are not being replaced.

*Casuarina equisetifolia* is naturalized in southern Florida, and over many decades the land it occupies has slowly increased. Along the coast it is appreciated for its shade and the pleasant sound when the wind whispers through its foliage. The spread of *Casuarina equisetifolia* into the interior of southern Florida is limited by the frequent fires and severe freezes that periodically eliminate it there. It is out of control, however, on some sandy coastal habitats.


*Casuarina glauca* is also found as an ornamental windbreak and is used for cattle shade in southern Florida. On the coast it is usually found near brackish waters, but inland is where it is more prominent. Because of its prolific root suckering, it is now considered a pest: some counties have laws banning its planting and the planting of other casuarinas. While its root suckers create "mini-jungles" and its wide-spreading root system disrupts pavements and lawns, the scarcity of

female trees in Florida fortunately restricts *Casuarina glauca* from getting entirely out of hand.

*Casuarina cunninghamiana* is less common in Florida, even though it is the most cold-hardy and the least objectionable. It is relatively slow growing and is intolerant of coastal sites.

#### HAWAII\*

Hawaii has perhaps tested more species of casuarina than any other location. Although there is estimated to be only 3,800 hectares of casuarina, 12 of the different species listed in this report are repre-



Hawaii. Casuarina shelterbelts are common around sugarcane fields.  
(C. S. Hodges)

\*Information from R. G. Skolmen.

sented.\* In addition, there are plantings of *Casuarina angularis*, *Casuarina nodiflora*, and an unknown species from Timor.

Locally known as "ironwood," casuarinas have been planted for erosion control, dune stabilization, windbreaks, fuelwood plantations, beautification, and watershed cover. In the lowlands, the most extensively planted species has been *Casuarina equisetifolia*. In the uplands, *Casuarina glauca* has been most commonly used, primarily for erosion control.

Experience accumulated over the years suggests that in Hawaii casuarinas are best suited for elevations below 300 m, with annual rainfall of 500–1,000 mm. Elsewhere they cannot compete in growth rate with other species, especially on good sites. In high rainfall areas, most casuarina species have been slow starters and have ended up as understory trees in mixed planted stands. On the other hand, in low rainfall coastal sites casuarinas can and often do compete successfully with *Prosopis* and *Leucaena*, which are also fast-growing nitrogen fixers.

## OTHER PLACES

*Casuarina equisetifolia* is extensively planted in Portugal, Tunisia, Corsica, and other Mediterranean sites. It is also common in West Africa, for example Togo and Benin, where casuarina is systematically grown for charcoal in sandy coastal soils on a 10- to 12-year rotation.

Two species of casuarina occur naturally in Fiji. *Casuarina equisetifolia* (nokonoko) and *Casuarina vitiense* (velau) are both popular firewoods; velau is also utilized in construction where hardness and strength are required.

Three species of casuarina occur in South Africa. *Casuarina cunninghamiana*, *Casuarina equisetifolia*, and *Casuarina glauca* grow rapidly and are hardy to both drought and frost. They are used for timber, poles, firewood, shelterbelts and windbreaks, and for reclamation.

\**C. cristata*, *C. cunninghamiana*, *C. decaisneana*, *C. equisetifolia*, *C. glauca*, *C. huegeliana*, *C. junghuhniana*, *C. littoralis*, *C. oligodon*, *C. papuana*, *C. stricta*, *C. torulosa*.



# 3

## The Plants


### FOLIAGE

As noted earlier, casuarina foliage consists of long “needles” with regularly spaced nodes. The needles are actually modified branchlets, termed cladodes, and differ from pine needles in that they are segmented and pull apart easily. The leaves are so reduced that they look like whorls of tiny teeth and are barely visible as small scales forming a ring around the node between each segment. The number of teeth in each whorl reflects the number of leaves and is, in part, diagnostic of the species.

The branchlets bear tiny ridges along their length formed from the fused leaves. The stomata (the pores for gas exchange) lie between the ridges, mostly in furrows, which protect them from the environment. This protection of the stomata endows casuarinas with great tolerance to drought and salt. But casuarinas have more features that help them resist stress. A thick, waxy cuticle covers the branchlets, and the roughly circular cross-section reduces the surface area exposed to the sun. Both of these features help reduce the plant’s water losses. In this way, casuarinas avoid desiccation in dry areas. Furthermore, the cylindrical form and high amount of structural tissue prevents wilting of the foliage at low tissue-water potentials. Thus, in drought conditions casuarinas can survive without danger of tissues collapsing.

### FLOWERING

Casuarinas have very reduced male and female flowers, arranged in catkin-like inflorescences. Most species have separate male and female trees (dioecious), but in a few species the male and female flowers occur on the same tree (monoecious). They are wind pollinated. After




*Casuarina equisetifolia*. Although these drawings are of one species, they are generally representative of the parts of most casuarinas. (P. B. Tomlinson)

pollination the female inflorescence develops into a small woody "cone," which has beak-like valves that open at maturity to release small winged seeds.

For many Australian species of casuarina, flowering takes place during a brief period of the year and is more or less consistent from year to year. Other species are less regular. *Casuarina cristata*, for example, flowers either in spring or following a heavy rain.

## NITROGEN FIXATION

As discussed previously, the root hairs of most casuarina species are invaded by the filamentous soil actinomycete, *Frankia*. When the microorganism's infectious threads reach the cortical root cells they divide and expand, forming lobes on the outsides of the root. The result is a woody perennial "nodule" that is spherical and up to 10 cm in diameter.



Leaf teeth, *Casuarina cunninghamiana*. The true casuarina leaves occur as tiny toothlike remnants fused to cylindrical branchlets. The stomates are found in grooves on the faces of the branchlet, a location that helps protect them from salt, desiccation, and other hardships. (D. J. Boland)

The association with *Frankia* benefits the casuarina, and the resulting symbiosis provides nitrogenous compounds that fertilize the plant's growth. Only the young tips of each nodule appear to actively fix nitrogen. The woody nodules are found in large masses from the base of the trunk out to near the drip line. They are most readily visible just beneath the soil surface, but they have been located as deep as 10 m in the soil.

The woody nodules are long lived, but eventually they decay, releasing *Frankia* spores and particles into the soil. The survival time and mechanisms of distribution of *Frankia* in the soil are unknown and deserve research.

The greatest number of nodules have been found where the soil activity is close to neutral. However, natural stands of *Casuarina glauca* are well nodulated in soils that are quite acidic (to pH of about 4). Nevertheless, nodulation normally is most successful within the range of soil pH from 6 to 8.

Nodulation is very much influenced by the available moisture and the soil aeration. Oxygen is necessary for the fixation process, so that active nodules normally occur only on surface roots that are obtaining adequate moisture. The symbiosis cannot occur in waterlogged sites. However, nodules are abundant in natural stands of *Casuarina glauca* where the water table is within 30 cm of the soil surface\* and nodules occur on the root system of *Casuarina cunninghamiana* growing at sites that are periodically submerged, such as river banks.

Also necessary for nitrogen fixation are minute amounts of mineral elements such as molybdenum, cobalt, and copper. These, however, are usually provided by even the poorest of soils.

Casuarinas fix about as much nitrogen as legumes. In a study in Senegal, the amount of nitrogen in soil under casuarina trees increased annually at rates of 58 kg per hectare when compared with nearby unplanted sand dunes.† In a natural forest of *Casuarina littoralis* near Sydney, Australia, the annual nitrogen accumulated in the litter under the trees was 290 kg per hectare, largely because of nitrogen fixation.‡ In one experiment, nodulated seedlings of *Casuarina glauca* grown in a greenhouse increased their shoot nitrogen content about thirteenfold within 60 days of the first appearance of nodules.§

The actual rate of nitrogen fixation appears to depend on environmental factors and on the casuarina species (and perhaps even on the

\*Information from P. D. Coyne.

†Dommergues, 1963 and 1976.

‡Clarke, L. D., and N. J. Hannon, 1970. Mangrove swamp and salt marsh communities of the Sydney district III. Plant growth in relation to salinity and waterlogging. *Journal of Ecology* 58:351-369.

§Information from P. D. Coyne.

particular strain of tree) as well as on the strain of the *Frankia* symbiont.

The *Frankia* that inhabits root nodules of actinorrhizal plants, such as casuarinas and alders, was first isolated and cultured from alders only in 1978. Since that time, microbiologists have isolated a dozen strains of *Frankia* from the roots of different host plants maintained in a worldwide collection in Middlebury College, Vermont (USA). Some of them have been shown to fix atmospheric nitrogen when grown in pure culture.

In 1983 pure strains of *Frankia* were, for the first time, isolated from casuarina nodules and cultured in artificial media. Tests have shown that these pure cultures are suitable as inoculum for casuarina seedlings and rooted cuttings.\*

Using such isolates and pure cultures, growers may in the future be able to match the best strain of *Frankia* to the appropriate casuarina species so as to maximize nitrogen fixation.

Whether a single *Frankia* strain will infect all of the casuarina species or whether more than one strain will be needed is not yet known, but some recent experiments suggest that different strains of *Frankia* are needed to infect and nodulate the different casuarina species.


## MYCORRHIZAL SYMBIOSES

Casuarina roots, in common with roots of many other tree species, have symbioses with a number of genera of soil fungi that help the trees scavenge mineral nutrients from the soil. These fungi are called mycorrhizae ("fungus-root"); two main types occur:

1. Ectomycorrhizae. These form an intricate meshwork of threads, or hyphae, clinging to the root surface and penetrating between root cells. From this mantle of fungal hyphae, thin filaments, barely visible to the naked eye, radiate out into the soil where they absorb nutrients and water and transport them back to the root. Ectomycorrhizal fungi on casuarinas include such genera as *Cenococcum*, *Pisolithus*, *Hymenogaster*, *Thelephora*, *Rhizopogon*, and *Amanita*.

2. Endomycorrhizae. The hyphae of these fungi actually invade the cortex of the root and proliferate between, and actually penetrate into, the root cells themselves. They are identified by microscopic arbuscules (hyphae-branched-like trees) and vesicles (thick-walled oval structures that develop within the cells). These fungi belong to the family Endogonaceae and are abundant in many soils. The most common genus is *Glomus*.

\*Diem, Gauthier, and Dommergues, 1983; Diem, Gauthier, and Dommergues, in press; Lundquist and Torrey, in press; and Zhang, Lopez, and Torrey, in press.



**Mycorrhizae.** Casuarinas have symbioses with two types of mycorrhizal fungi. Shown here are ectomycorrhizae that ensheath casuarina roots and facilitate the absorption of minerals such as phosphorus. These beneficial fungi help casuarinas survive and thrive in mineral-poor soils. (K. J. Mullette and R. K. Bamber)

The association of casuarina roots with both types of mycorrhizae significantly enhances the trees' adaptability as well as their ability to grow in harsh environments. Specifically, the fungi help the trees by:

- Improving mineral nutrition. Mycorrhizae have been shown to be particularly important in making phosphorus available to the tree, but they also help absorb soil nitrogen and microelements.
- Increasing tolerance to drought. This is of special importance during field plantings, as it prevents or improves recovery from wilting.
- Influencing the nitrogen-fixing activity of *Frankia*. In phosphorus-deficient soils, nitrogen fixation in root nodules is markedly reduced in the absence of mycorrhizae.
- Improving soil structure. Hyphal mats contribute to the binding of soil particles and thus reduce the harmful effects of wind and water erosion.
- Increasing resistance to some disease infections by preventing access of the organisms to the plant root.
- Alleviating the effects of acid soils, excessive aluminum, and certain other toxic soil conditions.



**Proteoid roots.** An unknown microorganism causes casuarinas to form dense, mat-like roots near the soil surface. These proteoid roots are very adept at absorbing nutrients because of their surface area and inherent efficiency. In experiments in low-phosphorus soils, they have doubled the plant's growth. (P. Reddell)

## PROTEOID ROOTS

Casuarina roots also interact with unidentified soil microorganisms that cause the development of "proteoid roots." These unusual structures occur as tight-packed rows of lateral rootlets that form thick mats just below the soil surface. They greatly increase the surface area that casuarina roots have for nutrient absorption.

To date, proteoid roots usually have been observed in association with organic debris in the soil and in sand under the tree drip-lines. Little is known about their benefits. However, other plants that form such roots demonstrate increased absorption of phosphate from soil.\*

\*Information from G. D. Bowen and N. Malajezule. The species tested was *Banksia grandis*; the proteoid roots on casuarinas have not yet been evaluated.

# 4

## Management

### PROPAGATION FROM SEED

Casuarinas are normally propagated by seed. As previously noted, most species have separate male and female trees, but some have both flowers on one tree. (See table.) Trees normally flower at an early age and produce seed in good quantities most years. Extracting the seeds is relatively simple: cones, or cone-bearing branches are placed in the sun to dry. In 2 or 3 days the woody valves (bracteoles) open and the winged seeds are released and can be collected.

The seed of most Australian casuarinas stores well when dried and kept in closed containers at room temperature. Tests showed little loss in germination after 12 years of storage for *Casuarina cristata*, *Casuarina decaisneana*, *Casuarina glauca*, *Casuarina stricta*, and *Casuarina torulosa*. On the other hand, seed of *Casuarina equisetifolia* is best stored at low temperatures, and although *Casuarina cunninghamiana* seeds will keep well for 1 or 2 years, the viability of even freshly collected seed is sometimes low. In general, seed of Asian and Pacific Island casuarinas seems to store poorly.

Seed of the different species varies greatly in size and germination requirements. Most germinate in 2–3 weeks, but *Casuarina decaisneana* is quicker, probably because it is adapted to take advantage of sporadic rainfall in its arid native habitat.

To produce seedlings in a nursery, conventional supplies—such as pots, tubes, and sleeves—can be employed. Even extremely primitive techniques are successful in many cases. Young trees are often transplanted bare-rooted, usually when 30–50 cm tall.

On many reforestation sites direct seeding is likely to be successful because casuarinas naturally reseed themselves very well on open, disturbed areas. Road banks in Australia, for example, are often revegetated naturally by casuarinas. Deep volcanic ash, that erupted



## Flowering Character and Seed Size

	Monoecious <sup>a</sup> (M) Dioecious <sup>b</sup> (D)	Mean Numbers of Seeds per Kg
<i>Casuarina campestris</i>	D	780,000
<i>C. cristata</i>	D	420,000
<i>C. cunninghamiana</i>	D	1,800,000 (>3,000,000)
<i>C. decaisneana</i>	D	85,000
<i>C. dielsiana</i>	D	270,000
<i>C. equisetifolia</i>		
var. <i>equisetifolia</i>	M	760,000 (616,000–1,488,000)
var. <i>incana</i>	M	470,000
<i>C. fraseriana</i>	D	235,000
<i>C. glauca</i>	D	1,760,000
<i>C. huegeliana</i>	D	510,000
<i>C. junghuhniana</i>	D	—
<i>C. littoralis</i>	D	480,000
<i>C. luehmannii</i>	D	—
<i>C. obesa</i>	D	840,000
<i>C. oligodon</i>	D	—
<i>C. stricta</i>	D	200,000
<i>C. torulosa</i>	D	260,000

<sup>a</sup> Male and female flowers occur on the same plant.

<sup>b</sup> Male and female flowers occur on different plants.

Source: D. J. Boland, J. C. Doran, R. D. Johnson, S. J. Midgley, and J. W. Turnbull.

in 1960 at Kapoho, Hawaii, has become completely revegetated by *Casuarina equisetifolia*, seeding itself during the years since the eruption. Natural seeding seldom occurs directly beneath a parent tree or under the canopy of a casuarina stand. However, abundant reproduction occurs in shallow, windblown litter in open areas adjacent to the trees.

## VEGETATIVE PROPAGATION

Probably all casuarinas can be propagated vegetatively. A sterile hybrid of *Casuarina equisetifolia* and *Casuarina junghuhniana* is reproduced almost entirely by cuttings in Thailand and India, and it seems likely that other species can also be propagated this way, although most have not been tried. Cuttings have the benefit of being a rapid, sure method of perpetuating the traits of superior specimens and also of allowing trees of a single sex to be planted so as to avoid weedy regeneration through the spread of natural seedlings.

A vegetative propagation technique recently developed in India for both *Casuarina equisetifolia* and the *Casuarina junghuhniana* hybrid involves the rooting of clusters of shoots (sprigs). These are dipped in rooting hormone (sometimes this step is omitted) and placed in a humid atmosphere—for example, in a sealed plastic tent in the light shade

under coconut trees. After 20 days they establish roots and can be planted out.\* This seems like a promising breakthrough for the rapid propagation of desirable casuarina specimens and it may be applicable to species other than the two used by the Indian researchers.

## MANAGEMENT

As discussed earlier, several species of casuarinas are characterized by "root suckering," that is, by the abundant formation of root-originated shoots. This ability makes them candidates for management under a sort of "coppice" rotation. However, it can create a random, dense mass that makes management difficult. Root suckering occurs in *Casuarina cristata*, *Casuarina glauca*, *Casuarina junghuhniana*, *Casuarina luehmanii*, *Casuarina obesa*, and *Casuarina stricta*. Some of these species sucker more readily than others, and it is not known whether this characteristic is genetic, and therefore limited only to them, or whether it is a general property of casuarinas that is elicited only when environmental conditions are favorable.

## INOCULATION

The *Frankia* actinomycete that forms the symbiosis with casuarinas is found naturally in soils where casuarinas are native. However, when the trees are planted outside their normal range the microorganism could be missing. In nursery plantations where sterile soil mixtures are used, the seedlings must be inoculated with *Frankia* from pure cultures or from ground-up nodules. It seems probable that even where casuarinas do form nodules without inoculation, the rate of nitrogen fixation might be improved by introducing a more effective strain of *Frankia*. Techniques for handling the microorganism were worked out in 1983.†

A practical inoculation technique for general use in the field is to shovel some surface soil from under a casuarina tree into the planting medium used in the nursery. Its *Frankia* spores then infect the seedlings.

A more sophisticated technique involves obtaining some root nodules from established trees and soaking them in 70 percent ethanol for a few seconds (to reduce the risk of contamination from microbes on the outside of the nodules). They are then washed, crushed in water, and the coarser particles filtered out. The resulting suspension is

\*Information from S. Kondas (see Research Contacts).

†Information from J. G. Torrey.

applied to the roots of casuarina seedlings or injected into the soil within the seedling's area. Fresh nodules can be dried over silica gel and stored for up to a year without losing their infective ability. Ground-up nodule suspension can be stored also in saline solution or in a polyvinylpyrrolidone solution.

*Frankia* are easier to handle than rhizobia (the bacteria that live symbiotically with many legumes and help them the way *Frankia* helps casuarinas) as the spores can be dried and stored at room temperature without losing viability.

In phosphorus-deficient soils, casuarina plantings should also include inocula of the mycorrhizal fungi that help the trees obtain phosphorus. In soils devoid of all available phosphorus (as might occur in some sand dunes, for instance), it is necessary to apply a small quantity of phosphate fertilizer, since this induces a good response from both plants and mycorrhizae. The mycorrhizae are not available in inoculum form; a sample of soil from around established trees is currently the only source.

## PLANTATION ESTABLISHMENT

Normally, casuarina plantings are given little site preparation. Even in the aridity of Egypt the saplings receive only a little watering during the first year, and after that they have to survive on their own.\*

Spacings differ with the end use and the site. For rain-fed plantations, 1,000–2,500 trees per hectare are used if billet-sized materials are desired. With irrigation, it is possible to plant 10,000 or more per hectare, although the products are smaller in diameter. In Egypt, for example, planters use an in-row spacing of 0.5 m for fuelwood and 2.5 m for scaffolding poles. Where plantations are established for wind protection in harsh sites, 3 m x 3 m spacing is appropriate. In Papua New Guinea 4 m x 4 m is used for shading coffee plantations.

## DISEASES

Although some diseases and pests have been noted in specific locations, casuarinas are generally healthy plants.

In India, the fungus *Trichosporium vesiculosum*, which causes a serious peeling of bark and may kill the trees, has been noted on

\*However, from Hawaii, Roger Skolmen reports that fertilizer (26 g of 10-30-10 per tree, or manure) is important to survival when planting in subsoil of eroded spots. Nodulation and mycorrhizae in the nursery are not enough. A mulch of casuarina litter can be used in place of fertilizer, but a combination is best.

*Casuarina equisetifolia*. It infects roots, especially where excessive watering and crowding occurs, such as along river banks. Another, but less serious, fungus, *Ganoderma lucidum*, starts as a saprophyte on nearby decaying stumps and then becomes a parasite on living trees by entering through wounds.

In the Guanica area of Puerto Rico, many casuarinas were killed in 1940 by a dieback disease, and stem canker and dieback have affected trees there since that time. Investigators attribute these syndromes to the fungus *Diplodia natalensis*. A similar problem was noted in some casuarinas in Florida in 1965.

In Florida there has been a high rate of mortality from mushroom root rot caused by the fungus *Clitocybe tabescens*, primarily on sandy soils previously populated by oaks. *Casuarina cunninghamiana*, *Casuarina equisetifolia*, and *Casuarina glauca* are all affected.

## PESTS

To date, no major insect problems have been encountered in adult plantations except for *Apte monachus*, the black borer, which tunnels into the stems and makes them susceptible to wind damage.

Nursery seedlings are liable to attack by crickets, chiefly *Brachytrupes achaetinus*. Caterpillars and aphids also occur in nurseries, but not in older trees and not when the nursery is near the sea.

In the sapling stage in Puerto Rico, grubs of *Coelosterna scabrata* have been noted boring into the root column, eventually killing the plant. *Arbela tetraonis* is a bark- and wood-boring caterpillar that can be easily detected by a characteristic covered trail winding over the bark. The caterpillar of *Phassus malabaricus* bores into the stem close to the ground and pupates in the larval burrow; it kills the tree by girdling it.

In Nigeria casuarinas up to age 3 years have been damaged or killed by termites.

## LIVESTOCK AND WILDLIFE

Young trees can be damaged by browsing or trampling, but they are resilient and tend to recover quickly. In Florida, deer sometimes browse seedlings so severely that the plants are killed. In California, as in Australia, young seedlings are susceptible to browsing by rabbits. Damage can be severe in some areas and seasons when other palatable vegetation is sparse, especially in desert environments. In Hawaii, on the goat-infested island of Kahoolawe, *Casuarina equisetifolia* survived nicely behind a fence but were eaten when the fence rusted away.

## TEMPERATURE

Most casuarinas are severely damaged or killed by frost. Non-Australian species, such as *Casuarina oligodon* and *Casuarina jung-huhniana*, grow in regions that are largely frost free. Coastal species such as *Casuarina glauca* rarely experience temperatures below  $-3^{\circ}\text{C}$ , while *Casuarina equisetifolia* occurs naturally only in frost-free areas.

However, in southern inland Australia, a few species such as *Casuarina cunninghamiana* and *Casuarina stricta* are found where occasional temperatures of  $-10^{\circ}\text{C}$  occur. And in California, in the Sacramento Valley, *Casuarina cunninghamiana* is more frost tolerant than either *Casuarina stricta* or *Casuarina equisetifolia*. *Casuarina cunninghamiana* has survived temperatures of  $-8^{\circ}\text{C}$  with no apparent injury.

## FIRE RESISTANCE

Most species of *Casuarina* are moderately tolerant of fire.\* *Casuarina decaisneana*, for instance, can tolerate extremely hot summer fires in Central Australia—the black color of its thick bark is a result of fire scorch. If tops are killed the trees will coppice, develop shoots from the roots, or (if not too severely damaged) produce new crown shoots. Some species shed seed soon after the seeds mature, so at times of fire there is usually no seed stored on the tree and seedling regeneration is therefore sparse.

Some casuarinas, however, appear to be fire-tender. These include *Casuarina fraseriana*, *Casuarina littoralis*, and *Casuarina torulosa*. They can be damaged by fire but produce vigorous new shoots afterwards. And because they carry cones for several years without shedding seed, seedling regeneration may be profuse in the seedbed created by the fire.

\*A major exception is the widely used *Casuarina equisetifolia*, which is killed outright by fire. This is a serious problem in Guam.

# 5

## Uses

### FUEL

Casuarina wood, as mentioned, has been called the best firewood in the world, and fuel is its most universal use. It is very dense, with a specific gravity ranging from 0.8 to 1.2. It is easy to split, has a high calorific value (about 5,000 kcal per kg), and tends to burn slowly with little smoke or ash. It also can be burned when green, an important advantage in fuel-short areas.



Fuel. Firewood cutters in stand of *Casuarina equisetifolia* near Madras, India. (N. D. Vietmeyer)



**Fuel.** Southern Highland villagers in Papua New Guinea with cords of *Casuarina oligodon* wood for firewood wait beside the road for passing trucks from nearby Mendi. Casuarina wood has been called the best firewood in the world. (N. D. Viemeyer)

When large trees are cut, the branches and brushwood are often used for fuel. In addition, the needles and cones burn well, so that the trees provide fuel even before they are cut down. From their fourth year, trees shed cones in quantities amounting to almost 4,000 kg per hectare. In India the cones are a cheap and handy fuel for hawkers selling roasted peanuts (groundnuts) from carts. They are also used to fuel pottery kilns.

Casuarina wood is excellent for charcoal making. Whereas most woods lose about three-quarters of their weight when made into charcoal, casuarinas lose only about two-thirds. The entire tree—stems, branches, and roots—can be converted to charcoal.

## WOOD

The wood of most casuarinas is so hard and heavy that it is difficult to saw and is not a good source of lumber. Moreover, it tends to split, crack, and warp as it dries.


Nevertheless, this dense, straight-grained wood has many important uses. Australian aboriginals use it to make boomerangs, and the first Europeans to settle in Australia found it useful to have about their farms for shingles, tool handles, fences, farm buildings, masts and oars, yokes, walking sticks, and other items. Even today in Western Australia, casuarina wood is commercially valued for wood turnery, tool handles, piano legs, shingles, veneer, and panelling.

In general, casuarina wood is useful as roundwood for fencing, pilings, beams, and rafters; as split wood for fencing, pilings, and roofing shingles; and as comminuted wood for particle board, pulp, and parquetry. In India it is used for scaffolding and structural members for buildings, as well as for masts for country fishing boats. In Egypt three particle-board factories are consuming casuarina windbreaks and shelterbelts so excessively that agronomists are becoming concerned that crops will suffer. The wood of *Casuarina equisetifolia* has been found to make a good paper pulp through use of the neutral sulfite semichemical process. But, as noted earlier, the difficulty of breaking up this extremely hard wood complicates pulping.



Roofing. Roof shingles being cut from *Casuarina torulosa* near Wauchope, New South Wales, Australia. (CSIRO Division of Forest Research)





Shelterbelt. Egyptian desert highways are often protected by *Casuarina glauca* and *Casuarina cunninghamiana* shelterbelts, many planted by university and high school students. (M. H. El-Lakany)

## SHELTERBELTS

*Casuarina equisetifolia* is often planted as a windbreak in North Africa, West Africa, Yemen, Somalia, the Middle East, India, and South China. In southern Australia *Casuarina stricta* has been widely used as a windbreak. The abundance of highly branched twigs on casuarinas absorbs wind energy amazingly well. A wind strong enough to blow hats off can be stripped of its force by a belt of casuarinas two or three deep, leaving the leeward air calmed. By reducing windspeed to almost zero, the shelterbelts stop wind erosion, and in areas with hot, dry winds they protect crops and animal herds and increase yields.

An advantage of casuarinas is that their roots do not readily harbor nematodes that affect the neighboring crops, a problem that is increasingly recognized as a serious limitation of shelterbelts. (In Egypt, however, *Casuarina equisetifolia* has been found to carry some nematodes, although *Casuarina glauca* has not.)

Egyptian farmers have solved the problem of root suckers coming

up in their fields. They dig a ditch between the crop and their *Casuarina glauca* shelterbelts. Sheep and goats then eat the exposed shoots and keep them from becoming pests.

In addition to their wind firmness casuarinas have desirable characteristics for shelterbelts: adaptability to many soils and climates, self-sufficiency for nitrogen, rapid early growth, adequate height and longevity, dense crown, and useful wood. It is most unusual for a single tree to have all of these attributes; to reduce wind adequately, shelterbelts normally require two or more species.

## EROSION CONTROL

Casuarinas can protect soil by reducing wind erosion, but they also do it with their network of fine subsurface roots and by building up a litter of intertwined needles that protects against rain and wind.

*Casuarina equisetifolia* is much used for stabilizing sandy soils (see below) and *Casuarina cunninghamiana* is valued for protecting riverbanks. For erosion control, the copious root suckering of species such as *Casuarina glauca* can be an advantage, because it helps the trees spread and hold down the land, especially on severe slopes or washed areas. In addition, the litter from the trees blows over the bare ground,



Streambank protection. *Casuarina glauca* is commonly used to protect canals that may otherwise clog up with just one sandstorm. (M. H. El-Lakany)



**Erosion control.** *Casuarina equisetifolia* near Fremantle, Haiti. The persistent, interlocked mat of fallen casuarina foliage encourages soil microflora, provides a good mantle against erosion, helps rebuild soil, and protects unstable sites. Infertile soils are improved through the nitrogenous compounds they receive through leaf fall, as well as through the decay of nodules and fine roots. (T. F. Geary)

protecting it from erosion and providing a good seedbed for natural reproduction.

## SAND DUNE STABILIZATION

Because it is salt- and drought-tolerant and can grow and reproduce in sand, *Casuarina equisetifolia* is used to control erosion along coastlines and estuaries. Thousands of square kilometers of casuarina forests have been planted in coastal sand dunes in southern China (see chapter 2). Casuarina is also used in Egypt to stabilize sand dunes that are drifting into the Nile Delta and Nile Valley. And in Mexico it is being planted in saline sandy areas near Mexico City to reduce dust storms. *Casuarina equisetifolia* is also stabilizing moving dunes on the coast of northern Senegal.

On the leeward side of sand dunes, when lower branches touch the ground they often root and develop upright branches. In this manner, they firmly fix the shifting sand with an expanding cover of trees.



Protection of agricultural land through dune afforestation. Under a World Food Program grant, almost 16,000 men and women planted *Casuarina equisetifolia* in a \$9 million project aimed at recovering impoverished or sand-choked soils in the area near Da Nang, Vietnam. (W.F.P. photo by F. Mattioli)

## TANNING

The bark of *Casuarina equisetifolia* contains 6–18 percent tannin and has been used extensively in Madagascar for tanning purposes. In Australia the tannin content is considered too low for commercial interest; however, casuarina bark enjoys a certain standing among amateur tanners. The tannins penetrate hides quickly and furnish a swollen, pliant, and soft leather of pale reddish-brown color. Other casuarina species probably could also be used for tanning.

## HONEY PRODUCTION

Casuarinas furnish no nectar for honeybees. However, *Casuarina littoralis* and *Casuarina cunninghamiana* are rated as having medium value as pollen sources; *Casuarina glauca* and *Casuarina torulosa* are rated as being of minor value.



**Ornamental.** A hybrid casuarina makes a pleasing backdrop to an ornamental planting at the Asian Institute of Technology, near Bangkok, Thailand. (N. D. Vietmeyer)



**City beautification.** *Casuarina cunninghamiana*, Tucson, Arizona. (T. R. Bundy, the University of Arizona)



**Roundwood. Lagaip River, Papua New Guinea. Flexible young casuarinas joined to form the arch of a bridge. (J. Siegel)**

## FODDER

While domestic animals will graze seedlings and suckers of casuarinas, the foliage is high in tannin and is astringent and constipating and may interfere with the animal's ability to utilize protein.

# 6

## Best-Known Species

Because of their proven performance, *Casuarina cunninghamiana*, *Casuarina equisetifolia*, *Casuarina glauca*, *Casuarina junghuhniana*, and *Casuarina oligodon* are each described in more detail in this chapter. Lesser-known species with promise are discussed in chapter 7.

Distribution maps of the Australian species were provided courtesy of CSIRO and are from J. C. Doran and N. Hall, 1981. *Notes on Fifteen Australian Casuarina Species*. CSIRO.

**Botanic Name** *Casuarina cunninghamiana* Miq.

**Common Names** River sheoak, creek oak, river oak, casuarina

**Main Attributes** The tree is one of the largest of the casuarinas. Handsome of appearance, it can be used for ornamental planting as well as for shade and shelter. It is a good street tree and is suitable for semiarid regions. It provides heavy shade and is almost trouble free. It has some cold tolerance, is very adaptable, and withstands periodic inundation so that its major use in Australia is for protecting stream banks from erosion. Because of its importance for this purpose, in New South Wales it may not be felled without a permit.

**Description** *Casuarina cunninghamiana* grows up to 40 m in height and more than 1 m in diameter. Its thin deciduous branchlets are soft and short and carry whorls of 6–8 leaf-teeth.

**Distribution** Native to eastern and northern Australia, it is found growing from the south of New South Wales (latitude 37°S) to the north of Queensland (latitude 12°S). Typically, it occurs fringing freshwater streams and rivers on both sides of the Great Dividing Range. A distinct race, possibly a separate species, is found along the larger rivers in the higher rainfall areas of the Northern Territory.

*Casuarina cunninghamiana* is extensively planted in Argentina and neighboring countries for windbreaks and to protect stream banks. It is also established in Florida. In Egypt it is important for shelterbelts. In Israel it is widely planted along railroads and highways for windbreaks and woodlots. And in Bulawayo, Zimbabwe, it is planted as a street tree.

### **Environmental Requirements**


- **Temperature.** Adapted to climates varying from temperate to tropical. The climatic range in its native habitat varies from warm subhumid to warm semiarid. The species is able to tolerate up to 50 light frosts per year—such incidences occurring mainly in the southern parts of its distribution.

- **Altitude.** Sea level to about 800 m.

- **Rainfall.** From 500 to 5,000 mm per year, but, as the tree is riverine, rainfall alone is no indication of the total moisture available. The incidence of the principal rainfall varies from a weak summer-autumn maximum in southern New South Wales to a strongly defined summer maximum (monsoonal) in the north.

- **Soil.** Soils are usually alluvial and vary from silty loams to sands





D. J. Boland

to gravels. In Hawaii it grows well on histosols developed over acidic lava (pH 5.0).

**Nodulation** Prolific.

**Root Suckering** Occurs, definitely in eastern Australia.

**Limitations** Not as salt tolerant as the closely related *Casuarina glauca*. On highly calcareous soils it is likely to turn chlorotic. Seedlings are susceptible to browsing damage and need protection from grazing in the early years.

**Botanic Name** *Casuarina equisetifolia* Forst. & Forst.

**Synonym** *Casuarina litorea* L.

**Common Names** Casuarina, sheoak, horsetail oak, Australian pine, ironwood, whistling pine, agofo (Philippines), ru (Malaysia), filao (Senegal), nokonoko (Fiji)

**Main Attributes** This is a large and usually long-lived tree and has the widest distribution both natural and planted of all the casuarinas. It has proved itself in scores of tropical countries. It is particularly valuable for stabilizing sand dunes, as it withstands salt spray, tolerates infertile soils, and is drought tolerant. Most of the experiences described in chapter 2 are with this species.

This is a species for warm to hot subtropical and tropical climates. Although it is not frost hardy, it tolerates a wide range of temperatures. The calorific value of its charcoal, 7,181 kcal per kg, is among the highest of any firewood species. The trees grow fast, sometimes reaching heights of 3 m within 8 months of transplanting to the field.

**Description** Two subspecies are recognized. Subspecies *incana* is a small tree that could be useful for low-growing shelterbelts but that has been little tried as an exotic. Subspecies *equisetifolia* has a tall stem and narrow crown. Unless otherwise stated, any reference to *Casuarina equisetifolia* in this book refers to this subspecies.


**Distribution** Subspecies *incana* occurs only along the coast of Queensland and northern New South Wales. The natural distribution of subspecies *equisetifolia* extends along seacoasts from Malaysia to subtropical Australia, Melanesia, Micronesia, the Philippines, and Polynesia.

### Environmental Requirements

- **Temperature.** In Australia these trees occur in the hot humid climatic zone, but they extend into the hot subhumid zone (while subspecies *incana* has its main distribution in the warm subhumid zone). Frosts are absent in most of its habitats, although in the south there may be 1–3 frosts a year.

- **Altitude.** This is generally a lowland tree but is grown up to about 600 m elevation in Hawaii.

- **Rainfall.** In its natural habitat, annual rainfall is from 700 to 2,000 mm, often with a dry season of 6–8 months. However, it has been planted successfully in areas with annual rainfall as little as 200–300 mm or as much as 5,000 mm.



R. D. Haller

● **Soil.** The species tolerates calcareous and slightly saline soils. Reportedly it grows poorly on fine-textured soils but it grows well on clays in Hawaii. It can withstand partial waterlogging for a time.

**Nodulation** Prolific.

**Root Suckering** Occurs.

**Limitations** Initially, the trees have poor ability to compete with weeds, especially in dense grass cover. Seedlings are vulnerable to attack by ants, crickets, and other insect pests. The trees are susceptible to root rot. *Casuarina equisetifolia* can exhaust the moisture in the soil, lower the water table of the site, and restrict growth of a healthy understory. The tree is fire sensitive and can be browsed only lightly without being damaged. Although other *Casuarina* species coppice readily, *Casuarina equisetifolia* coppices only to a limited extent and only when cut young (3–4 years).

**Botanic Name** *Casuarina glauca* Sieb. ex Spreng.

**Common Names** Swamp sheoak

**Main Attributes** This tall tree can survive on difficult sites where other trees fail because of salinity, waterlogging, or shallow water table. It withstands even periodic tidal inundation. Sometimes it occurs where it gets drenched in salt spray from the sea. In Egypt it has proved more drought tolerant than *Casuarina cunninghamiana* and *Casuarina equisetifolia*. In Israel it outperforms all the other casuarinas, reaching 20 m after 12–14 years, even on saline water tables.

It is favored for shelter purposes. Its canopy tends to commence at ground level, owing to low branching or the development of root suckers, which arise spontaneously around the parent tree.

**Description** This is an erect, fast-growing tree, typically 10–14 m tall, although in favorable localities it may reach 30 m. The main stem, which may be buttressed and fluted, is moderately straight and typically dominant for most of the tree's height. Free-growing specimens have a somewhat sparse and narrow crown. The slender deciduous branchlets (about 1 mm in diameter) carry 12–20 small leaf-teeth in widely spaced whorls. They are much coarser (and a little longer) than those of *Casuarina equisetifolia* or *Casuarina cunninghamiana*.


**Distribution** *Casuarina glauca* is found in a narrow belt hugging the coast of eastern Australia from Bega in New South Wales to Rockhampton in Queensland. It has also been successful in the marshes and saline soils of Israel, Cyprus, India, Kenya, Malawi, South Africa, Egypt, and Florida.

This casuarina is most common along the edges of swampy flats near estuaries and along tidal reaches of rivers. It is also sometimes found on or near beach fronts. The flats may be only marginally above tidal limits; the water table is usually close to the surface (often within 30 cm of the surface).

#### **Environmental Requirements**

- Temperature. 5°–33°C.
- Altitude. Sea level to 900 m (in Hawaii).
- Rainfall. 500–4,000 mm (in Hawaii).
- Soil. In its native habitat soils are often sandy or clayish and are underlain with rock. Salt content is often high. However, the trees occasionally occur on moist, rocky headlands.

In Israel the tree grows on very dry sites in the Negev Desert. It has grown where a layer of salt covers the soil surface (50,000 ppm).



D. J. Boland

Although most natural stands are on acidic soils, *Casuarina glauca* has shown good growth on alkaline clay-loam soils with shallow water tables in hot semiarid areas of Central Australia. In Thailand the seedlings have tolerated high levels of calcium in the soil and as much as 30 percent limestone. In southern Florida it flourishes on oolitic limestone. In Hawaii it is frequently planted on much-weathered parent basalt in eroded blowouts, sometimes in holes blasted by dynamite. It is also planted in pure limestone sand and does well.

**Nodulation** Prolific. The tree sometimes produces huge growths from which root suckers later arise.

**Root Suckering** Even without root damage it forms thickets of suckers. Suckering is increased by pruning the trees.

**Limitations** Suckers are grazed by livestock, but they have little value as forage. Because it regenerates vigorously from root sprouts, this species is a possible weed and, as already noted, is a pest in parts of Florida and is regarded as noxious in Hawaii.

In Hawaii it does not attain the height of *Casuarina cunninghamiana* or *Casuarina equisetifolia*, probably because it stagnates from abundant suckering.

**Botanic Name** *Casuarina junghuhniana* Miq.

**Synonym** *Casuarina montana* Jungh.

**Common Name** Jemara (Indonesia)

**Main Attributes** This tall, straight casuarina has good form for a plantation species. It is adapted to hot, humid conditions, to monsoonal climates with a long dry period, and to highland areas in the tropics. It grows quickly in compact clay soils near Bangkok, Thailand. It has some salinity tolerance and thrives on many types of soil. It reportedly competes well with vigorous tropical weeds such as *Imperata cylindrica*. So far, it is widely used only in Thailand. There the trees do not produce seed and are reproduced artificially using vegetative methods.

**Description** This is a long-lived pioneer tree that grows to majestic size—35 m in height and 1 m in diameter. The branchlets are gray-green to dark green and carried in a narrow conical crown.


**Distribution** *Casuarina junghuhniana* is native to highland regions of eastern Indonesia—to East Java, Bali, and the Lesser Sunda group of islands. There it occurs in extensive pure stands on mountain summits. It pioneers the natural revegetation of deforested grassland, volcanic ash and sand, gravelly stream beds, and screes. Man-made grassland has allowed it to extend its area manyfold, at the cost of mixed mountain forest and scrub forest that formerly prevailed.

A hybrid of *Casuarina junghuhniana* and *Casuarina equisetifolia*\* was introduced to Thailand around 1900. Today, its good, straight stems and symmetric conical crown make it a popular ornamental and cultivated crop throughout the country.

#### **Environmental Requirements**

- **Temperature.** Unknown.
- **Altitude.** Its natural distribution is up to 3,000 m. In Thailand and South India, however, it is mostly grown near sea level.
- **Rainfall.** In its native habitat the rainfall pattern is monsoonal with a well-defined summer maximum and a range from 700 to 1,500 mm. The tree is moderately drought tolerant. (In 1951 the hybrid was introduced into South India from Thailand. In 1972 a severe drought killed all 6-year-old *Casuarina equisetifolia* trees, but not the *C. junghuhniana* x *C. equisetifolia* hybrid.)

\*The identification of this tree as a hybrid was made in 1982 by L. A. S. Johnson. Information from Pravit Chittachumnonk.



**Pravit Chittachumnonk**

- **Soil.** It can grow well in various types of soil, from compact clay to light volcanic soil.

**Nodulation** Nodules have been found on a tree in the Bandung region of Indonesia.

**Root Suckering** The root system extends laterally and produces abundant root suckers.

**Limitations** In Thailand and India the hybrid trees do not produce seed and are propagated only by vegetative means, usually by air-layering, using the abundant suckers or branches.

**Botanic Name** *Casuarina oligodon* L. Johnson

**Common Names** Yar or soft yar (Papua New Guinea), sheoak

**Main Attributes** The Papua New Guinea government is encouraging local people to grow *Casuarina oligodon* as firewood in the Highlands. The Office of Forests in the Highland provinces distributes seedlings, which are planted by local landowners. The trees are highly self-regenerating, and they help in reforesting grasslands. The wood is used in construction, but because it is durable, it is also suitable for fences. Its main use is for fuelwood in the Highlands where nights are cool. It is also used for windbreaks, and most villages have *Casuarina oligodon* planted around their boundaries. The tree grows well even with competition from vigorous grasses (such as *Imperata cylindrica*, *Saccharum robustum*, and *Themeda australis*). Coffee is a major cash crop in the Papua New Guinea Highlands, and *Casuarina oligodon* is useful on coffee plantations because it shades the coffee bushes and improves the soil by fixing nitrogen.

**Description** This is a medium to tall tree, reaching 30 m in height and 0.6 m in diameter. The bark is gray-brown and fissured, peeling off in irregular flakes. The tree produces small cones (diameter less than 1 cm).


**Distribution** The species is native to New Guinea and is found mostly in upland valleys. It forms extensive pure stands along river beds, but at times it is associated with *Casuarina papuana* (see page 94). It occurs commonly on stream banks and ridge tops and on abandoned garden and village sites. In the Papua New Guinea Highlands, people fallow the land 10–20 years under *Casuarina oligodon*. It is part of their traditional land-use pattern.

This species is little known outside New Guinea, but in Hawaii it has grown very rapidly.

#### **Environmental Requirements**

- **Temperature.** This is truly a tropical highland species.
- **Altitude.** The tree is found up to 2,500 m or higher in Papua New Guinea.
- **Rainfall.** It occurs naturally in areas where the rainfall ranges from 1,900 mm to 2,600 mm and humidity is high throughout the year. In Hawaii it is planted in an area with 5,000 mm rainfall but good drainage.
- **Soil.** It is mostly found in sandy soils along creeks and rivers but





N. D. Vietmeyer

grows well in colluvial soils, humic brown-clay soils, and alluvial and meadow soils.

**Nodulation** Occurs.

**Root Suckering** The species mainly spreads through seed and generally does not spread through suckers. However, it has been known to form epicormic shoots after fire or when damaged by other natural causes.

**Limitations** *Casuarina oligodon* may not grow successfully on poor soils. It seems to be very sensitive to areas of high salt concentrations. It has also proved susceptible to wind breakage in Hawaii; it appears to be much more brittle than any of the other species.



# 7

## Other Promising Species

In addition to the five species described in chapter 6, there are several lesser-known casuarinas that have real or potential value. As more introductions are attempted and more is learned about them, they could make an important place for themselves. None has yet been tested significantly outside its native habitat.

The following pages describe:

*Casuarina campestris*

- |                       |                      |
|-----------------------|----------------------|
| <i>C. cristata</i>    | <i>C. littoralis</i> |
| <i>C. decaisneana</i> | <i>C. luehmannii</i> |
| <i>C. dielsiana</i>   | <i>C. obesa</i>      |
| <i>C. fraseriana</i>  | <i>C. stricta</i>    |
| <i>C. huegeliana</i>  | <i>C. torulosa</i>   |

**Botanic Name** *Casuarina campestris* Miq.

**Synonym** *Allocasuarina campestris* (Miq.) L. Johnson

**Main Attributes** This species provides perhaps the ultimate in wind protection. Single rows with plants 2–3 m apart provide excellent shelter for livestock and dwellings. *Casuarina campestris* protects soil well and is planted for conservation purposes. It is now used mainly for low, hedge-type shelterbelts. It will grow on a wide range of soils, including coastal sands.

**Description** Although it may also grow as an erect tree with a dense crown, this species is usually a very bushy, multistemmed shrub only 1–3 m high.

**Distribution** *Casuarina campestris* is native to the southwestern part of Western Australia.

### **Environmental Requirements**

- **Temperature.** This species is native to temperate areas of Western Australia. Throughout most of its range it receives 1–12 frosts per year.

- **Altitude.** It occurs from near sea level to about 375 m.

- **Rainfall.** It is found mainly in the semiarid climatic zone, but it also extends into both subhumid and arid zones. The 50th percentile rainfall is 225–400 mm, and the lowest precipitation on record in the area is 140–250 mm. There is a clearly defined winter maximum.

- **Soil.** *Casuarina campestris* has been recorded on some extremely poor soils such as those derived from cemented laterite, greenstone, ferric-conglomerate, broken quartzite, and limestone. It has been also recorded on loams, shallow clays over laterite, and coarse gravelly soils.

**Nodulation** Unreported.

**Root Suckering** Unreported.

**Limitations** The wood is available only in small diameters. Its technical properties are unknown.



**CSIRO Division of Forest Research**

**Botanic Name** *Casuarina cristata* Miq. There are two subspecies: *Casuarina cristata* subspecies *cristata* and *Casuarina cristata* subspecies *pauper*.

**Synonyms** *Casuarina lepidophloia* F. Muell.; *Casuarina glauca* Benth., in part; not Sieb.

**Common Names** Black sheoak, belah

**Main Attributes** This tree is valued as a source of light shade and shelter on farms, for timber, and, to a limited extent, as an emergency drought fodder.

Subspecies *pauper* is more suited to drier (as low as 250 mm annual rainfall) and harsher conditions than subspecies *cristata*, as well as to areas where there is limestone in the soil. It also appears suitable for saline soils.

Subspecies *cristata* seems to require a rainfall greater than 400 mm. It is an excellent tree for windbreaks and is an attractive ornamental. Specimens growing in the open develop a dense crown, almost to ground level.

**Description** This smallish tree reaches a height of 20 m. It is erect in form, and when trees are planted close together the main stem is straight for more than half the total height.

**Distribution** Subspecies *cristata* is native to eastern Australia. It grows in a belt on the inland side of the Great Dividing Range in New South Wales and southern Queensland. Subspecies *pauper* is native to a wide belt from western New South Wales across the central latitudes of South Australia to Western Australia.


### Environmental Requirements

- **Temperature.** Subspecies *pauper* occurs in an arid region with burning hot summers and a few winter frosts.

- **Altitude.** The tree is found mainly below 350 m.

- **Rainfall.** Subspecies *pauper* grows where annual rainfall is very erratic, as low as 175 mm, and where there is a potential annual evaporation of 2,500 mm or more. In this region of low, open woodland or shrub steppe, this casuarina is often the only tree to be seen.

- **Soil.** This species seems to withstand compact clay soils and high alkalinity—pH 8.8 and lime content up to 3.5 percent calcium in the surface soil, for example.



D. J. Boland

**Nodulation** Good at most sites. In dry inland areas nodules are more difficult to locate as they are generally found deeper in the soil.

**Root Suckering** It reproduces readily by root suckers, often forming large, dense stands of one sex, which could make it a pest in some areas.

**Limitations** Young trees must be protected from grazing animals.

**Botanic Name** *Casuarina decaisneana* F. Muell.

**Synonym** *Allocasuarina decaisneana* (F. Muell.) L. Johnson

**Common Name** Desert sheoak

**Main Attributes** This is one of the remarkable trees of central Australia, the hottest and driest part of the continent. In parts of this area it may be the dominant tree of the landscape, occurring as scattered individuals or in groves. Frequently, it is the only tree species present in a landscape where even shrubs are not common.

**Description** This tree has a graceful pendulous habit when mature, but the young plants are more upright. The trunk is straight and may be more than half of the total height of 10–15 m. The cones are ovoid and oblong and are 3–5.5 cm long and 2.5–3.5 cm thick.

**Distribution** It occurs mainly south and west of Alice Springs in the southern half of the Northern Territory and in adjacent areas of Western Australia and South Australia.

#### **Environmental Requirements**

- **Temperature.** The trees withstand incredible conditions. Mean maximum temperature of the hottest month is about 35°C, with readings up to 47°C common. On the other hand, winter temperatures may fall to –7°C.

- **Altitude.** Within the range of 250–700 m.

- **Rainfall.** The mean annual rainfall value is 220 mm, based on 13 years of observation; in the lowest recorded year, only 38 mm of rain fell. The potential evaporation may exceed 3,000 mm a year.

- **Soil.** In its native habitat, *Casuarina decaisneana* grows on deep sandy soils and on undulating or low sites where the maximum seepage into the subsoil can be expected during the rare heavy rain.

**Nodulation** Nodules have been found on this species in the field, but they were not common at the sites examined.

**Root Suckering** Unreported.

**Limitations** The species has not been widely tested, and there is no knowledge of it under cultivation. It appears to be slower growing than other casuarinas. Some specimens, drip-irrigated at Alice Springs, reached 4 m in 6 years. The trees may have restricted soil requirements perhaps being intolerant of alkali.





CSIRO Division of Forest Research

**Botanic Name** *Casuarina dielsiana* C.A. Gardn.

**Synonym** *Allocasuarina dielsiana* (C.A. Gardn.) L. Johnson

**Common Name** Diels' sheoak

**Main Attributes** This small, compact tree provides good shelterbelt or shade trees when young (with age, the lower branches die). The Western Australian Forests Department recommends it for use in multirow shelterbelts. It merits trials for amenity planting or for the production of fuelwood and timber in areas with high summer temperatures, low and erratic rainfall, and rocky and freely draining soils.

**Description** The tree is commonly 5–8 m tall on moderately good sandy loams, but it is sometimes reduced to a bush only 4 m high on the top of dry gravelly ridges. The crown tends to be carried in the upper part of the trunk, while the lower parts are thinly branched.

**Distribution** It is a native of Western Australia, mainly within 300 km of the west coast and between latitudes 26°–30°S.

#### **Environmental Requirements**

- **Temperature.** Summers are long and very hot, with the mean maximum of the hottest month approaching 38°C.
- **Altitude.** Its habitat ranges in elevation from near sea level to about 300 m.
- **Rainfall.** Its native area averages 300–500 mm, with a well-defined winter maximum and only moderate variability.
- **Soil.** This species is typically found around or on the more rocky and rugged sites, and while it may edge sand plains, it is rarely found on them. The soils are mainly sand, sandy loams, or stony skeletal. Other soil types include gravels on ironstone ridges and light-red loams associated with granitic outcrops. Although it occurs naturally on well-drained, poor, and rocky soils, it is thought to have potential for compact clay soils in areas where the rainfall may be as low as 300 mm.

**Nodulation** Unreported.

**Root Suckering** Unreported.

**Limitations** Unreported.



SIRO Division of Forest Research

**Botanic Name** *Casuarina fraseriana* Miq. (formerly spelled *C. fraserana*.)

**Synonym** *Allocasuarina fraseriana* (Miq.) L. Johnson

**Common Name** Fraser's sheoak

**Main Attributes** This is a tall tree with promise for winter rainfall areas with nutrient-poor lateritic clays and sandy soils. In the southwest of Western Australia, one company is licensed to cut casuarinas and it utilizes only *Casuarina fraseriana*. Its products are wood turnery items (goblets, planters, saucers, bowls, and piano legs), roof shingles, and veneer for paneling. Reject logs are made into dowelling, moldings, and picture frames. Firewood is still cut from this species for the Perth market.

**Description** At its best this is an erect, somewhat pyramidal tree with a straight bole two-thirds of the tree height and an open crown. In particularly favorable areas it may reach heights of 14–15 m, but it is commonly only 9–12 m tall. Even on lateritic clay the height may be 10 m, but on poor sites such as dry sandy plains or rocky areas, the trees may be reduced to tall shrubs 2–5 m high.

**Distribution** This is a species of coastal southwestern Western Australia. It occurs mainly on a narrow belt on the seaward side of the ranges.

#### **Environmental Requirements**

- **Temperature.** This is typically a species of warm humid climates, but it extends slightly into the adjacent warm subhumid zone. Areas immediately adjacent to the sea are frost free, while further inland, frosts may average up to 11 per year.

- **Altitude.** From 50 m to about 500 m above sea level.

- **Rainfall.** The 50th percentile rainfall has a wide range, varying from 900 to 1,250 mm; in the lowest year on record, 500–750 mm of rainfall was recorded. There is a well-developed winter maximum rainfall.

- **Soil.** This casuarina is found on a wide range of soils, including deep, white, acid sands, oxisols, ultisols, and gravels, as well as yellow (iron stained) calcareous sands near Perth, Western Australia. It does not grow on the adjacent compact clay soils. It can handle nutrient-poor soils, but they must be well drained.

**Nodulation** Very variable; absent on some specimens, good on others.



CSIRO Division  
of Forest Research

**Root Suckering** Cut stumps produce vigorous new shoots and may be suitable for coppice management. Living trees are often burnt into hollow trunks in very hot wildfires; however, the trees recover, and new branches break out from beneath the very thick bark to form badly disfigured survivors.

**Limitations** Unreported. Actually, the species exhibits no features to suggest that it is invasive or would cause problems difficult to control.

**Botanic Name** *Casuarina huegeliana* Miq.

**Synonym** *Allocasuarina huegeliana* (Miq.) L. Johnson

**Common Name** Rock sheoak, granite sheoak

**Main Attributes** This is one of the largest casuarinas in the inland dry areas of Western Australia. The Forests Department of Western Australia considers that this species merits greater recognition for shade planting on poor, shallow sandy soils with low rainfall. The species also deserves consideration for ornamental planting on some difficult sites.

**Description** This tree is commonly only 5–12 m tall, although under favorable conditions it sometimes attains 14 m. In close stands the trees have a columnar habit, while open-growing specimens have a wide crown and a main stem that divides at one-fourth to one-third of plant height.

**Distribution** This casuarina is native to semiarid and subhumid areas in the southwest of Western Australia. Much of its occurrence is in the wheat belt.

### **Environmental Requirements**

- **Temperature.** In its native habitat summers are hot, with mean maximum temperatures of the hottest month 29°–35°C, while mean minimum temperatures of the coldest month are as low as 4°C. Areas near the coast may be frost free, but inland there are 1–10 frosts a year.

- **Altitude.** It grows from just above sea level to 450 m.

- **Rainfall.** This is in the range of 300–750 mm. It is a winter rainfall zone with dry summers.

- **Soil.** The species derives its common names—rock sheoak and granite sheoak—from its occurrence on massive rocky outcrops (inselbergs). It is also found on sandy plains and on highly ferruginous banded ironstone. It has been recorded on a wide range of other soil types: yellow clay-sands, sandy and gravelly loams, sandy clays, sandy types of all gradations, and oxisols and ultisols.

**Nodulation** It is difficult to obtain quantitative data because of problems excavating root systems at sites where this species grows.

**Root Suckering** Not known.



CSIRO Division of Forest Research

**Limitations** The tree is less suitable for shelterbelts, since the moderately dense crown is carried on the upper part of the stem, though it can be used in association with lower-story shrubs such as *Casuarina campestris* (page 68).

**Botanic Name** *Casuarina littoralis* Salisb.

**Synonyms** *Casuarina suberosa*, *Allocasuarina littoralis* (Salisb.) L. Johnson

**Common Name** Black sheoak

**Main Attributes** Probably the most plentiful casuarina in Australia, this is mainly an understory species of tall open forests, but it occurs in woodland areas and in the more open patches of closed forest. It probably provides the neighboring trees with the nitrogen required for growth. This is a tree for relatively low, narrow shelterbelts and for use in avenues where larger trees are undesirable. It recolonizes sand blows and regenerates vigorously from seed drop after forest fires.

**Description** *Casuarina littoralis* is usually a small open tree or sometimes a large shrub in the height range of 3–12 m. In forest and shrubland associations this species has an erect habit and narrow crown, but on windswept coastal heaths it may be reduced to a procumbent plant.

**Distribution** This tree has the widest range of latitudes of any Australian casuarina. It is native to a belt (mainly within 100 km of the sea) from the northern tip of Queensland (latitude 12°S) to Tasmania (latitude 43°S).

### **Environmental Requirements**


- **Temperature.** The species is found in widely varying climates. Most of the trees occur in warm subhumid and humid climates, but appreciable stands are found in the cool subhumid zone and some in the hot humid zone. At the highest altitudes in New South Wales the trees are subjected to as many as 70 frosts a year.

- **Altitude.** While most plentiful at altitudes under 300 m, the tree is found at elevations up to 1,200 m.

- **Rainfall.** The 50th percentile rainfall is 650–1,250 mm and the lowest on record is 300–500 mm. In the far north, rainfall follows a strong monsoonal pattern with very dry winters and springs. This pattern changes southwards and has a more or less uniform distribution in Tasmania.

- **Soil.** Black sheoak grows on a variety of sites from lowland flats to undulating topography and mountain peaks, though it is most common on well-drained hills and mountain slopes. It may be found in rocky gorges, near the edge of swamps, on heathlands, and on





T. F. Geary

sandy lowlands behind sandy dunes. The most common soil types are sand, podzolics, skeletal, and rocky areas where soil is almost absent. Well-drained sites are preferred.

**Nodulation** Variable; absent on some specimens, prolific on others. Environmental influences are critical for nodulation.

**Root Suckering** Unreported.

**Limitations** Not suited for very humid (5,000 mm rainfall) histosols in Hawaii.

**Botanic Name** *Casuarina luehmannii* (R. T. Baker)

**Synonym** *Allocasuarina luehmannii* (R. T. Baker) L. Johnson

**Common Name** Bull oak

**Main Attributes** This is mainly an open woodland species that is tolerant of light frost, poorly aerated soils, waterlogging, and slight salinity. It is often an understory tree that can be planted in association with tall trees. It is suitable for amenity planting in parks or avenues. It will provide satisfactory but somewhat open shelterbelt protection. On moderately good sites its early growth is relatively fast, and there are reports that it will withstand some seasonal inundation by floodwaters.

**Description** This is a medium-sized tree, reaching heights of 20 m and diameters of 60 cm. On all except the poorest sites, the main stem is straight and often up to half the total height, after which it divides into several large ascending stems that carry the rather open crown.

**Distribution** This casuarina has a wide natural occurrence in eastern Australia, from northern Queensland to eastern South Australia.

#### **Environmental Requirements**

- **Temperature.** In its native habitat summer temperatures are warm in the south and hot in the north, with mean maximum temperatures of the hottest month varying from 29°–35°C.

- **Altitude.** *Casuarina luehmannii* commonly grows on plains, gently undulating slopes, and, especially in the drier areas, around the edges of swamps and depressions where surface drainage is poor, but it also extends—as a poorer tree—to slopes and ridges.

- **Rainfall.** For most of its area the rainfall is 380–630 mm, but in northern New South Wales and Queensland it may be as high as 800 mm.

- **Soil.** Soils show a wide range, from poor, well-drained sands to sandy loams and poorly drained clays, all usually low in nutrients.

**Nodulation** No nodulation has yet been found.

**Root Suckering** Under some conditions the species may develop a suckering habit.

**Limitations** Seed is difficult to obtain and germinates poorly.



CSIRO Division of Forest Research

**Botanic Name** *Casuarina obesa* Miq.

**Common Name** Western Australian swamp sheoak

**Main Attributes** This species grows well in wetlands near somewhat saline waterways in both coastal and semiarid inland situations. The species is salt tolerant, and it forms abundant nodules near the soil surface. This tree is closely related to *Casuarina glauca* and can probably be grown in the same locations and used in the same ways. For instance, it can be planted as a decorative row of shade trees along the seaside. However, it grows in more inland situations than *Casuarina glauca* and may be even more drought tolerant.

**Description** This species is a handsome, well-shaped tree that may reach about 14 m high. The main stem is branchless for much of its height, and in close stands, if the growing situations is good, it is usually moderately straight.

**Distribution** The tree is predominantly found along seacoasts and shores of inland saline depressions of Western Australia.

#### **Environmental Requirements**

- **Temperature.** For most of the area the mean temperature of the hottest month is 29°–34°C and the mean minimum of the coolest month 5°–9°C. Coastal areas, especially from Perth northwards, are frost free, but elsewhere there are 1–12 heavy frosts a year.

- **Altitude.** From near sea level to about 300 m.

- **Rainfall.** Its native habitat has a pronounced winter rainfall. The South Australia occurrence is in the arid zone, but near the boundary in the semiarid zone. It is drought tolerant down to 200 mm.

- **Soil.** It often occupies low-lying, swampy flats just above high-tide limits and along river banks and the edges of inland salt lakes. In these areas it may grow as pure, dense thickets or as the principal small tree. Soils include a wide range of sands and silts. Other soils on which it has been recorded include pink granitic sands, red clay loams, and swampy soils.

**Nodulation** Prolific.

**Root Suckering** Occurs.

**Limitations** Unreported.



**CSIRO Division of Forest Research**

**Botanic Name** *Casuarina stricta* Ait.

**Synonym** *Allocasuarina verticillata* (Lam.) L. Johnson

**Common Names** Drooping sheoak, coast sheoak, mountain oak

**Main Attributes** *Casuarina stricta* is a graceful ornamental with a spreading form and beautiful, drooping blue-green branches. The branchlets are among the largest (46 cm) of any casuarina. It has demonstrated good growth on irrigated lands in Egypt and has shown early success in Cyprus, India, Israel, and several countries of southern Africa. It grows rapidly (up to 2 m per year), even on poor soils, but after a few years it stops, so that it remains a small tree. The trees form effective shelterbelts and tolerate lopping well, which allows most of the smaller branches of the crowns to be cut off to provide fodder in times of severe drought without killing the trees. On even moderately good soils, open-grown specimens are useful for shade and ornament.

**Description** Usually only about 6–10 m high, this species typically has a straight, but usually short, main stem. When it is grown in the open, the crown is dense, deep, and almost as wide as the tree is high. The smallest branchlets droop.

**Distribution** The tree is found on the coastal plains and ranges of New South Wales, Victoria, South Australia, and Tasmania and grows from near the high water mark to inland sand dunes.

#### **Environmental Requirements**

- **Temperature.** The tree probably requires a temperate climate. In its native habitat mean minimum temperature in winter is from  $-2^{\circ}$  to  $+5^{\circ}\text{C}$ ; mean maximum in summer is  $15^{\circ}$ – $32^{\circ}\text{C}$ . The number of frosts varies from 0 to 40 per year.

- **Altitude.** The tree is found from coastal headland areas to tablelands at about 800 m elevation.

- **Rainfall.** The 50th percentile rainfall is within the limits of 600–900 mm, with the lowest on record 250–375 mm.

- **Soil.** *Casuarina stricta* grows on dry ridges and rocky ground, poor coastal sands, and saline soils. It is found growing in salt-spray and wind-affected coastal sites.

**Nodulation** It nodulates well.



CSIRO Division of Forest Research

**Root Suckering** The trees coppice vigorously with shoots from the cut stumps. It is a species likely to produce thickets of suckers, but it is not an invasive weed.

**Limitations** Where the tree is lopped for fodder in times of drought, little natural regeneration is found.

**Botanic Name** *Casuarina torulosa* Ait.

**Synonym** *Casuarina tenuissima*, *Allocasuarina torulosa* (Ait.) L. Johnson

**Common Names** Forest sheoak, rose sheoak

**Main Attributes** This species produces better timber than other casuarinas. The wood is less liable to checking and splitting and this is the only species that is commercially sawn for construction timber in Australia.

**Description** This is a handsome tree, attaining 25 m in height and 1 m in diameter on favorable sites. The bole is often more than half the tree height, the main branches are spreading to erect, and the crown has an open appearance.

**Distribution** Native to the coastal ranges of eastern Australia from northern Queensland to southern New South Wales, this tree is associated with wetter forests than most Australian casuarinas. It is shade tolerant and typically grows as an understory species in open forest or grows in tall open-forest formations.

#### **Environmental Requirements**

- **Temperature.** Winter minimum, 2°C; summer maximum, 31°C in its native habitat.
- **Altitude.** Sea level to 1,100 m.
- **Rainfall.** 900–2,500 mm evenly distributed or with a summer maximum often in areas with a relatively dry spring.
- **Soil.** *Casuarina torulosa* adapts to a wide range of soils from sandy alluvials to compact clays. It has been recorded on steep slopes, moderate hillsides, tablelands, and undulating lowlands, while the soils noted include tertiary basalts, rhyolite, quartzite, various sedimentary rocks, sandstone, and shale.

**Nodulation** Observed on most specimens examined. No quantitative information.


**Root Suckering** Unreported.

**Limitations** This species grows on poor soils, but not as poor as those where the other common eastern Australian species, *Casuarina littoralis*, grows (page 80).





**K. B. Walsh**



# 8

## Recommendations and Research Needs

The work yet to be done on casuarinas challenges researchers in many parts of the world and in disciplines such as botany, forestry, soil science, microbiology, ecology, and ethnobotany. For philanthropic institutions, foundations, and international development agencies concerned with problems of fuel, fiber, and other resources, casuarina research presents an area worthy of financial support.

Specific research needs and recommendations follow.

### ASSESSMENT OF GENETIC RESOURCES

In the past, the introduction of casuarinas has been made haphazardly, without regard to relationships between species, provenances, and site requirements. Yet there is much genetic diversity within the broad group of casuarinas and within individual species. It is now important to learn about this diversity and to broaden the genetic base of casuarina plantations. The time has come to make extensive field collections, with a view to selecting species and individual plants with the greatest potential for specific sites and purposes.

In particular, it is important to establish seed collections and to study populations from many and diverse provenances. Almost all seeds so far distributed have come from southern Australia. Species from the Pacific Islands and more tropical areas have been little used, nor have their populations been studied. Few of the many species of Western Australia have been collected, studied, or tested elsewhere. Because so many casuarinas are native to Australia, it is clear that much of the research on the genetic resources will be done there and will be crucial for casuarina development in the rest of the world.

Assessment of genetic resources should include:

- Accurate mapping of worldwide distribution;
- Seed collections of known species from identified provenances;
- Assessment of the importance of hybridization, both natural and experimental; and
- Seed distribution and genetic-information exchange.

## COOPERATIVE WORLDWIDE PLANTATION TRIALS

Commercial experience with casuarinas is restricted to a small number of species planted randomly throughout the tropics. It is now imperative to establish a series of carefully planned and replicated plantation trials at selected sites throughout the tropical and subtropical world to allow comparisons of growth and performance. A common methodology should be used at each trial location. Sites for the trials should be selected that test the responses of various species to different soils, altitudes, latitudes, temperatures, moisture levels, and pests. Information from the trials, brought together for assessment and comparison, will provide the technical foundation to enable rational choices about the establishment of large casuarina plantations under the most favorable conditions.

This effort in international scientific cooperation will require a coordinator with sufficient funding to collect and distribute seeds, to establish contact and maintain correspondence with research groups, and, ultimately, to collect data and publish results of the trials.

## SILVICULTURAL RESEARCH

Information available in the published literature, though suggesting a promising future for casuarinas, is insufficient to support recommendations for the immediate establishment of large plantations throughout the tropics and subtropics. Much of the information still needed must come from carefully planned and coordinated silvicultural trials. However, basic research in a number of areas can make future forestation efforts more effective, less expensive, and more widely applicable.

Silvicultural research on casuarina species is needed in the following areas:

- Phenology of flowering and fruiting;
- Methods of seed storage, especially of tropical species;
- Nutrient requirements, including microbial interactions;

- Methods of establishing plantations, including planting densities and cluster planting;
- Methods of vegetative propagation from mature superior trees, including the influence of tree age, sex, and type of cuttings, the use of hormone treatments, potential of tissue culture and of air-layering;
- Intercropping with other tree crops such as *Eucalyptus* for biomass, timber, or pulp and of interplanting with agricultural field crops;
- Coppice cropping and root suckering;
- Fire tolerance and regeneration after fire;
- Photosynthetic effectiveness, including effects of stress;
- Investigation of potential plant pathogens, nematodes, and other plant pests; and
- Accurate measurements of growth and productivity of casuarinas.

Casuarinas are pioneer species with the ability to establish and persist on poor sites. They exhibit capacities to flourish under stressed environments that include extremes of soil acidity and salinity, water stress (both flooding or drought), high temperatures, and other adverse conditions. Physiological studies of these tolerances will make possible the best use of the genetic capacities within the group.

## MICROBIOLOGICAL RESEARCH

Specific research needs into the casuarinas' symbioses include the following:

- Isolation and culture of *Frankia* strains, optimum methods of inoculation and nodule establishment, and combinations that most effectively boost symbiotic nitrogen fixation;
- Vegetative propagation of trees selected on the basis of their ability to nodulate and fix nitrogen; and
- Identification of mycorrhizal fungi, as well as development of inoculation methods and analysis of phosphorus and micronutrient requirements of casuarinas.

## UTILIZATION RESEARCH

Examples of research areas needed on casuarina utilization include:

- The design and planning of shelterbelts;
  - Methods of improving land stabilization and controlling erosion;
  - Use of casuarinas for biomass plantations;
  - Methods for making pulp and particle board from casuarina wood;
- and
- Conversion of casuarina wood to firewood and charcoal.

## RESEARCH ON OTHER SPECIES

This report has highlighted 18 of the 67 species of casuarina. Among the remainder are species certain to be just as valuable as those we describe here—they are merely less well known. These species offer interesting research challenges to botanists and foresters. Species worth particular study include the following:

- *Casuarina collina* A little-known tree native to New Caledonia. It seems to have properties similar to those of *Casuarina glauca*, but it occurs in more humid and tropical areas. It is well worth more-widespread testing.

- *Casuarina grandis* Often larger than *Casuarina cunninghamiana* (to 50–60 m), but commonly 35–40 m. This tree is closely related to *Casuarina cunninghamiana* and *Casuarina junghuhniana*. It is known only from southeastern Papua New Guinea from near Tufi in the Northern Province to the Gwariu River in the Milne Bay Province. It forms dense stands along rivers, regenerating on gravel banks and other open sites, possibly partly from root suckers.\*

- *Casuarina papuana*† One of the most attractive species. It makes an excellent ornamental in Hawaii and has pleasing foliage display and red adventitious root masses on the trunk. A medium-sized tree native to Papua New Guinea, it is found at low altitudes and is also scattered in the Highlands. It grows mostly on rocky creek banks or in secondary forests on ridges. The species is useful because it can grow on poorer soils than *Casuarina oligodon* and tolerates a wide range of rainfall and temperatures in the country. It is used mostly as firewood but can also be used for heavy construction (usually in the round), tool handles, shuttles, charcoal, and pilings permanently submerged in fresh water.

The tree grows to 30 m with dark open crown composed of tufted branches; its branchlets are not obviously drooping. It grows commonly in hill and mountain upland areas, notably on poor or variously nutrient-imbalanced soils, but it may also occur along rocky river banks and on top of limestone hills and pinnacles. It is restricted to the Moluccas and New Guinea. The tree is often cultivated, both in villages and in towns; it is known in New Guinea Pidgin as “Hat Yar.”

- *Casuarina nobilis*‡ Tree reaches at least 25 m, with slender bole, growing in various types of forest and bush, usually on nutrient-poor or unbalanced soils. It has been reported by Whitmore and others working on Borneo vegetation as occurring in heath forest (*kerangas*),

\*Information from D. G. Frodin.

†*Gymnostoma papuanum* (S. More) L. Johnson in sched.

‡*Gymnostoma nobile* (Whitmore) L. Johnson in sched.

peat-swamp forest, forest on ultramafic rocks, and on limestone hills and pinnacles. It has, when young, a bushier habit that *Casuarina papuana*. It is found in Borneo and New Guinea and is cultivated in towns and settlements (for instance, Kepong, West Malaysia, and Lae, Papua New Guinea).

● *Casuarina rumphiana*\* Found in Moluccas, Celebes, and Philippines, in mountain forests, 10–14 m high. It is a slow-growing species.

The tree grows to 20–25 m with large open crowns when mature. The “needles” have a tendency to drop somewhat and not to form tufts as bushy as those of *Casuarina papuana* and *Casuarina sumatrana*. It grows on dry fire-prone slopes, limestone hills, and volcanic blast areas from low elevations to about 1,000 m. It is a common local tree and is perhaps “encouraged” in villages. On the karstic Lelet Plateau in New Ireland all trees of this species form a striking feature of the landscape, sometimes giving a park-like aspect on the limestone hills. It is the characteristic upland casuarina of New Britain, as well as New Ireland, the Philippines, Moluccas, and the Bismarck Archipelago. It was originally described by Rumphius in the 17th century.

● *Casuarina sumatrana*† A moderately large tree up to 30 m tall with a clear cylindrical bole. It occurs in heath forest in Borneo on very poor, acid soils from sea level to 1,000 m. It grows quickly and has been planted as a fuelwood crop in Sarawak. It produces a very high grade of charcoal and is a popular ornamental tree. The wood seasons well with little degrade. Nodules were found on all 72 trees examined in Bogor. It is slow growing.

The tree grows to 30 m with crowns similar to those of *Casuarina papuana*, but its cones are about twice the size. In Papua New Guinea it is only known from one locality where it grows in swamps mixed with sago palms, but records from Irian Jaya suggest it also grows in nutrient-poor upland areas. The trunk has better form than that of *Casuarina papuana*; in swamps it may be stilt-rooted. It is found in Burma, Sumatra, Java, Borneo, Celebes, Philippines, western Irian Jaya, and New Britain. It is also cultivated in towns and settlements (for example, Kepong, West Malaysia).


● *Casuarina deplancheana* An efficient colonizer of ferrallitic soils in New Caledonia. It is resistant to fire. Both endomycorrhizae and nitrogen-fixing nodules with *Frankia* have been identified on its roots.

● *Casuarina spicigera*‡ Grows to perhaps 20 m along rocky banks of lowland rivers, like *Casuarina grandis* (see above). The crowns are fairly open. It is restricted to the northern side of the north coast

\**Gymnostoma rumphianum* (Miq.) L. Johnson in sched.


†*Gymnostoma sumatranum* (Jungh. ex de Vriese) L. Johnson in sched.

‡*Gymnostoma spicigerum* L. Johnson in sched.



*Casuarina nobilis*, Brunei, Borneo. (P. S. Ashton)





*Casuarina papuana*, Hawaii. (R. G. Skolmen)



*Casuarina grandis*, Southeastern Papua New Guinea. (D. G. Frodin)

ranges of Papua New Guinea and also to the Van Rees range of Irian Jaya. The tree is locally cultivated, for example, at Vanimo in Papua New Guinea.

## DISSEMINATION OF INFORMATION

The panel recommends that casuarina researchers undertake to publish three documents:

1. A field guide to the identification of members of the family *Casuarina* and hybrids between them. This is an urgent task if confusion and wasted effort are to be avoided as researchers in various parts of the world begin work with the different species.

2. Planting guides. Handbooks should be published with practical, step-by-step information on propagating, planting, managing, and utilizing the plants.

3. A newsletter. In exploring the potential of casuarinas as useful crops, it will be important to maintain communication among researchers working with the plants. Since these researchers are likely to be situated in far-flung research stations, universities, missions, and villages, their findings may not be widely noted in technical journals. A newsletter would bring together results from botany and forestry science, would provide rapid exchange of information, and would be a forum for informal opinions, observations, and preliminary experimental data that are usually not accepted by journals.

## Appendix A

### Selected Readings

A report that complements this one is: Midgley, S. J., J. W. Turnbull, and R. D. Johnston. 1983. *Casuarina Ecology, Management and Utilization*. Proceedings of an International Workshop. Canberra, Australia, August 17–21, 1981. Commonwealth Scientific and Industrial Research Organization, Melbourne, Victoria, Australia. 286 pp.

Copies are available from: Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600, Australia.

A newsletter that keeps up with research developments on casuarina is published by the Casuarina Working Group of the International Union of Forestry Research Organizations. Copies are available from: M. H. El-Lakany, Department of Forestry, Faculty of Agriculture, University of Alexandria, Egypt.

A bibliography containing 447 casuarina references, with annotation, was prepared for an International Meeting on Casuarina, Canberra, August 17–21, 1981, by Heather Howard, Division of Forest Research, CSIRO. The following is a selection from the bibliography.

#### GENERAL

Burkill, I. H. 1966. *A Dictionary of the Economic Products of the Malay Peninsula*. Governments of Malaysia and Singapore by the Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia.

Conover, C. A., and E. W. McElwee. 1971. *Selected Trees for Florida Homes*. Florida Cooperative Extension Service Bulletin 182. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, Florida, USA. 77 pp.

Little, E. L., Jr., and F. H. Wadsworth. 1964. *Common Trees of Puerto Rico and the Virgin Islands*. Agricultural Handbook No. 249. Forest Service, U.S. Department of Agriculture, Washington, D.C., USA. 548 pp.

Maiden, J. H. 1889. *The Useful Native Plants of Australia* (facsimile edition of the 1889 edition,) produced by Compendium Pty, Ltd., Melbourne, Victoria, Australia, 1975.

Midgley, S. J., J. W. Turnbull, and R. D. Johnston. 1983. *Casuarina Ecology Management and Utilization*. Proceedings of an International Workshop, Canberra, Australia, August 17–21, 1981. Commonwealth Scientific and Industrial Research Organization, Melbourne, Victoria, Australia. 286 pp.

Rhoads, A. H. 1952. *The destructiveness of Clitocybe root rot to plantings of casuarinas in Florida*. *Lloyda* 15(3):161,184.

Schubert, T. H. 1979. *Trees for urban use in Puerto Rico and the Virgin Islands*. General Technical Report SO-27. Forest Service, U.S. Department of Agriculture, Washington, D.C., USA. 91 pp.

Watt, G. 1889. *A Dictionary of the Economic Products of India*. Printed by the Superintendent of Government Printing, Calcutta, India.

## SOIL AND SYMBIOSES

- Diem, H. G., D. Gauthier, and Y. R. Dommergues. 1983. Inoculation of *Casuarina* using a pure culture of *Frankia*. Nitrogen Fixing Tree Research Reports 1:18-19.
- Diem, H. G., D. Gauthier, and Y. Dommergues. In Press. An effective strain of *Frankia* from *Casuarina* sp. Canadian Journal of Botany.
- Dommergues, Y. 1963. Evaluation du taux de fixation de l'azote dans un sol dunaire reboise en Filao [Nitrogen-fixation rate of a dune soil afforested with *Casuarina equisetifolia*.] Agrochimica, Pisa 7(4):335-340.
- Analysis of the nitrogen content of a plot at the Cape Verde peninsula established 13 years earlier showed an addition of 64 kg to the 80 kg per ha present in the nonafforested soil. Taking into account contribution by rainwater and losses through leaching, the annual fixation is estimated at 58.4 kg per ha, chiefly by the nodules of *Casuarina equisetifolia*.
- Junor, R. S. 1978. Control of wind erosion on coal ash. Journal of the Soil Conservation Service of New South Wales 34(1):8-13.
- Coal ash from the Lake Tallawarra power station is sluiced with water from a saline lake and allowed to settle in large ponds. Once dry, it is subject to water and wind erosion. *Acacia decurrens* and *Casuarina glauca* were successfully planted on a stabilization trial and have since spread naturally over the site.
- Lundquist, R., and J. G. Torrey. In press. the propagation of *Casuarina* species from rooted stem cuttings. Botanical Gazette.
- Zhang, Z., M. F. Lopez, and J. G. Torrey. In press. A comparison of cultural characteristics and infectivity of *Frankia* isolates from root nodules of *Casuarina* species. Plant and Soil.

## GENERAL BOTANY

- Barlow, B. A. 1959. Chromosome numbers in the *Casuarinaceae*. Australian Journal of Botany 7(3):230-237.
- Chromosome numbers of 37 species of *Casuarina* are reported and found to conform with a suggested division of the genus into major groups. The extra-Australian species ( $\times = 8$  or 9) are probably ancient. Somatic chromosome numbers from 28 to 20 occur in Australian stocks.
- Narashimhan, M. J. 1918. Preliminary study of the root nodules of *Casuarina*. Indian Forester 44:265-268.
- Preliminary study of root nodules showed that *Casuarina* trees improved sandy soils so much that inland flora regenerated.

## SYSTEMATIC BOTANY

- Doran, J. C., and N. Hall. 1981. Notes on fifteen Australian *Casuarina* species. Division of Forest Research, CSIRO, Canberra, A.C.T., Australia. 34 pp.
- Notes are provided for 15 Australian *Casuarina* species and subspecies with value or potential for planting. The description of each species includes a map of geographical distribution and photos showing tree, habit, "cone," and bark. The main characteristics of each species are tabulated.
- Godfrey, A. L. 1979. *Casuarina*—the neglected tree. Victoria's Resources 21(4):31-33, 35.
- Describes species occurring in Australia and covers some of their useful qualities.

Johnson, L. A. S. 1982. Notes on Casuarinaceae II. Journal of the Adelaide Botanic Gardens 6:73-87.

Terry, M. 1941. Desert oak. Australian Timber Journal 7(2):82-83, 85.

A popular account of *Casuarina decaisneana*, which is widely distributed in dry sandy country in south central Australia. It grows 35-50 ft in height and has a tough wood that is apparently highly resistant to termites.

Treub, M. 1891. Sur les Casuarinees et leur place dans la systeme naturelle. Annales du Jardin Botanique de Buitenzorg 10:145-231.

Article describing the *Casuarina*. Text in French. No English summary.

Troup, R. S. 1921. Order LVII. Casuarinaceae. *Casuarina*, Forst. Pp. 900-908 in *Silviculture of Indian Trees*, R. S. Troup, ed. Clarendon Press, Oxford, England.

Describes distribution, habitat, reproduction, and plantation methods for *Casuarina equisetifolia*.

## PLANT ECOLOGY

Becking, J. H. 1970. Plant-endophyte symbiosis in non-leguminous plants. *Plant and Soil* 32(3):611-654.

This review includes a survey of the non-leguminous plant species (14 genera of 7 families) that bear root nodules and are capable of nitrogen fixation. Most of these plants are woody and are important to the colonization of bare soil. The particular roles of *Alnus glutinosa* and *Casuarina equisetifolia* are described, with quantitative data on nitrogen fixation.

Dommergues, Y. 1976. Mycorrhizes et fixation d'azote. [Mycorrhizae and fixation of nitrogen.] *Anales de Edafologia y Agrobiologia* 35(11/12):1039-1056.

A literature review, drawing attention to the lack of knowledge about the reasons why nonsymbiotic nitrogen fixation occurs in the mycorrhizosphere of some plants but not others. Suggests lines for further research into known cases of tripartite associations between plant host, nitrogen-fixing bacterium, and mycorrhizal fungus; for example, in *Casuarina*.

Gauthier, D., H. G. Diem, and Y. Dommergues. 1981. In-vitro nitrogen fixation by two actinomycete strains isolated from *Casuarina equisetifolia* nodules. *Applied and Environmental Microbiology* 41(1):306-308.

Lawrie, A. C. 1982. Field nodulation in nine species of *Casuarina* in Victoria. *Australian Journal of Botany* 30(4):447.

Mendoza, V. B. 1978. Adaptability of six tree species to cogonal areas: additional information on the possible role of phenols and sugars. *Sylvatrop* 3(1):1-7.

A leaching experiment was conducted to investigate whether *Imperata cylindrica* produced compounds that may be harmful to plant growth. *Casuarina equisetifolia* displayed the best height growth in *Imperata* grass.

Torrey, J. G. 1978. Nitrogen fixation by actinomycete-nodulated angiosperms. *Bio-Science* 28(9):586-592.

Describes the infection process in the root system and the ultrastructure of the host-endophyte relationship. Evaluates the significance of actinomycete-nodulated plants in providing nitrogen through mixed plantations, succession plantings, and other forestry management practices.

## REGENERATION AND FORMATION OF STANDS

Anonymous. 1954. Research studies: the mountains; shallow loam problem area. *Caribbean Forester* 15(1/2):8-9.

Of the trees tested on the shallow, erodible soils of Puerto Rico, *Casuarina equisetifolia* displayed the best growth.

Barr, D. A. 1965. Restoration of coastal dunes after beach mining. *Journal of the Soil Conservation Service of New South Wales* 21(4):199-209.

Restoration procedures include stripping, stockpiling, and replacement of topsoil, leveling of tailings, and revegetation. After initial stabilization with brush matting, primary stabilizers (grasses) are sown, followed by secondary stabilizers (creepers and shrubs) and tertiary stabilizers (trees: *Casuarina equisetifolia*).

Esbenshade, H. W., and A. Grainger. 1980. Bamburi reclamation project. *International Tree Crops Journal* 1(2/3):199-203.

Eighty acres of the Bamburi limestone quarry near Mombasa, Kenya, have been reclaimed since 1971 and 30,000 trees have been planted. *Conocarpus lancifolia* and *Casuarina equisetifolia* are grown for timber and fuelwood on the quarry floor. The quarry sides grow fodder for bees and livestock.

Geary, T. F., and C. B. Briscoe. 1972. Tree species for plantations in the granitic uplands of Puerto Rico. USDA Forest Service Research Paper ITF No. 14. Forest Service, U.S. Department of Agriculture, Washington, D.C., USA. 8 pp.

Gives details of the survival and growth of 32 tree species tested for adaptability in Puerto Rico's humid granitic uplands, a region of sandy, well-drained erosive soils. *Casuarina equisetifolia* is recommended for post and pole crops.

Husain, A. M. M., and P. K. Ponnuswamy. 1980. Propagation of *Casuarina junghuhniana* by planting shoots and root suckers. *Indian Forester* 106(4):298-299.

*Casuarina junghuhniana* does not produce seeds, as the original stock is only male. Therefore, it is essential to propagate only by vegetative means. This paper describes rooting of sprigs and root suckers vegetatively.

Karschon, R. 1960. Effects of spacing and irrigation on yield of eucalypts and *Casuarina*. *La-Yaaran* 10(1/4):9-11, 61-58.

Shows the advantage of wider spacing and irrigation in the production of wood chips, posts, pulpwood, and poles. With 9-year-old *Casuarina* spp., irrigation improved the volume m.a.i. almost twofold.

Karschon, R. 1962. *Casuarina* introduction trials (preliminary results). *La-Yaaran* 12(2):41-43, 69-67.

Some results of small-scale trials, in Ilanot and the northern Negev, of 15-16 species of *Casuarina*. Most successful are *C. cunninghamiana*, *C. decaisneana*, *C. fraseriana*, *C. glauca*, *C. lepidophloia*, *C. luehmanni*, and *C. torulosa*.

Le Roux, P. J. 1974. Establishing vegetation in saline soil to stabilise aeolian sand at Walvis Bay, South-West Africa. *Forestry in South Africa* 15:43-46.

Describes the climate and soil conditions of the area and reports on trial plantings on sand dunes and saline wet silt since 1970. *Casuarina equisetifolia* failed on saline silt and dune sand irrigated with seawater. *Casuarina equisetifolia* grew well under irrigation with sewage water.

Maheut, J., and Y. Dommergues. 1959. Fixation par le reboisement des dunes de la presqu'île du Cap-Vert et l'évolution biologique des sols. [Sand dune fixation by afforestation on the Cape Verde peninsula and the biological development of the soil.] *Bois et Forêts des Tropiques* 63:3-16.

Describes successful afforestation with *Casuarina equisetifolia*, including topography, climate, and soils of the site; nursery techniques, spacing, irrigation, and protection.

Malcolm, C. V., and S. T. Smith. 1971. Growing plants with salty water. *Journal of Agriculture of Western Australia, Series 4* 12(2):41-44.

For those using saline irrigation water, this article gives some hints on how to reduce salt damage to plants. Includes a table of plants that may be irrigated with water of specified degrees of salinity and lists precautions to be taken with each group. *Casuarina cristata* and *Casuarina glauca* are listed.

Mitchell, B. A. 1957. Malayan tin tailings—prospects of rehabilitation. *Malayan Forester* 20(4):181–186.

Describes conditions to be dealt with and research into possible methods of rehabilitation and afforestation, as well as improved methods of dumping tin tailings. Species trials show that *Casuarina equisetifolia* is promising on very infertile tailings.

Patro, C., and R. N. Behera. 1979. Cashew helps to fix sand dunes in Orissa. *Indian Farming* 28(12):31–32.

An outline of pilot dune-stabilization and shelterbelt trials with plantations of *Casuarina* /cashew and cashew/coconut. Details are given of establishment and tending methods.

Somasundaram, T. R., and S. S. Jagadees. 1977. Propagation of *Casuarina equisetifolia* Forst. by planting shoots. *Indian Forester* 103(11):735–738.

Although *Casuarina equisetifolia* is usually produced from seed, cuttings taken from young lateral shoots, with a heel smeared with Seradix 2 and planted under mist, showed up to 90 percent rooting success (maximum with July plantings). Shoots without treatment did not survive.

Spurway, B. J. D. 1950. Ru Ronang (*Casuarina sumatrana*) as a plantation crop. *Malaysian Forester* 13(2):73–74.

A successful experiment was made in planting wildling seedlings of *Casuarina sumatrana* at a spacing of 7 ft × 7 ft in prepared lines. The seedlings averaged 8 in. from root tip to top shoot, and survival was good. This species is considered suitable for establishing fuel plantations.

Toth, J. 1965. Aspect forestiere d'une plantation Saharienne. (Forest aspect of a plantation in the Sahara.) *Revue Forestiere Française* 17(10):674–695.

Describes and illustrates the establishment, since 1959, on coarse and fine sands with pH 8 near the surface of 35 ha of shelterbelts and amenity blocks in the north Sahara. *Casuarina torulosa* has been planted as windbreaks. Propagation by seeds and cuttings, and natural and artificial regeneration under the plantations, are briefly discussed.

van der Westhuyzen, J. J. N. 1957. Combating sand dunes at Port Edward: trees and shrubs used in reclamation. *Farming in South Africa* 33(7):37–39.

Recommendations for the fixation of sand dunes (probably the result of overgrazing) at Port Edward on the south coast of South Africa include the control of livestock and the planting of at least three zigzag rows of *Casuarina equisetifolia* 4–6 ft apart and 30–50 yards to windward.

## COMBINATIONS OF FORESTRY WITH AGRICULTURE AND PASTORAL HUSBANDRY, IRRIGATED WOODLAND FORESTS, AND SHELTERBELTS

Anonymous. 1980. Planim diwai yar. [Grow *Casuarina*.] Mapping Branch, Office of Forests, Papua New Guinea. 18 pp.

Booklet with illustrations showing how the indigenous population plant and grow *Casuarina* with their other crops.

Badran, O. A., and M. H. El-Lakany. 1978. Breeding and improving *Casuarina* for shelterbelt plantations in Egypt. Pp. 573–578 in *Third World Consultation on Forest Tree Breeding*, Volume 2. CSIRO, Canberra, A.C.T., Australia.

In Egypt, and in many countries with no natural productive forests, *Casuarina* spp. are receiving increased attention because of their superiority as windbreaks and their high salt and drought resistance. Gives some primary results of a program to provide fast-growing, drought-resistant casuarinas that have desirable stem, crown, and branch characteristics.

Brockway, G. E. 1959. Tree establishment in the wheatbelt. Bulletin, No. 2616. Department of Agriculture, South Perth, Western Australia. 21 pp.

Examines the problems of establishing trees, mainly for cover, recreation, and fodder in semiarid areas in Western Australia. Includes a comprehensive tabulated planting guide listing requirements and uses of four *Casuarina* species.

Leigh, J. H., A. D. Wilson, and W. E. Mulham. 1979. A study of sheep grazing on a belah (*Casuarina cristata*)—rosewood (*Heterodendrum oleifolium*) shrub woodland in western New South Wales. Australian Journal of Agricultural Research 30:1223–1336.

Koki, Z. 1978. Studies on flying salt in Okinawa from the viewpoint of seashore conservation. Science Bulletin of the College of Agriculture, University of the Ryukyus 25:429–554.

As part of a study on the distribution and effects of wind-borne salt, salt deposition was measured on various trees. *Casuarina equisetifolia* with needle-like leaves collected more salt than the broad-leaved trees examined. It was estimated that tree shelterbelts should reduce downwind salt deposition by 60 percent, as compared with 40–50 percent for nets. Text in Japanese.

Morton, J. F. 1976. Pestiferous spread of many ornamental and fruit species in South Florida. Proceedings, Florida State Horticultural Society 89:348–353.

The massive invasion of large tracts of South Florida by *Casuarina equisetifolia* and other introduced species is an environmental problem now receiving serious attention. Further plantings are being discouraged to reduce the maintenance load of cultivated grounds and the threat to undeveloped areas being overrun by vigorous alien vegetation.

Sanger, C., G. Lessard, and G. Poulsen. 1977. Trees for people; an account of the forestry research program supported by the International Development Research Centre. International Development Research Centre, Ottawa, Canada. 52 pp.

Describes 10 different projects including breeding and improvement of *Casuarina* spp. for shelterbelts in Egypt.

Wilson, A. D., and W. E. Mulham. 1980. Vegetation changes and animal productivity under sheep and goat grazing on an arid belah (*Casuarina cristata*)—rosewood (*Heterodendrum oleifolium*) woodland in western New South Wales. Australian Rangeland Journal 2(2):183–188.

## ARBORICULTURE FOR ORNAMENTAL PURPOSES

Garnet, J. R. 1965. *Casuarina*. Australian Plants 3:185–187.

Descriptions of casuarina as street trees.

Ratnasabapathy, M. 1974/75. *Acacia auriculaeformis* and *Casuarina equisetifolia*—the urban invaders. Malayan Nature Journal 28(1):18–21.

Presents notes on the ecology, propagation, and pests (relatively few) of these two popular street trees in the new town of Petaling Jaya, Malaysia, which are also the dominant trees in pioneer vegetation on disturbed sites in the same area.

Weinstein, A., and G. Schiller. 1979. Afforestation and tree planting in Sinai. II. Northwestern Sinai. La-Yaaran 29(1–2):12–16, 32.

A list of species planted in agricultural and urban settlements between Rajah and El Arish is presented. *Casuarina glauca* and *Casuarina cunninghamiana* are among the ornamental species successfully grown under irrigation.

## FOREST PRODUCTS AND THEIR UTILIZATION

Bawagan, B. O., and A. A. Faulmino. 1978. Qualities of agocho del monte dissolving pulp. Forpride Digest 7(1):12–17.



A good-quality dissolving pulp was obtained from *Casuarina rumphiana* wood using the pre-hydrolysis-sulfate process. Describes physical and chemical properties.

Guha, S. R. D., and R. N. Madan. 1963. Chemical pulps for writing and printing papers from *Casuarina equisetifolia*. *Indian Forester* 89(5):365-367.

Laboratory experiments in sulphate pulping of *Casuarina equisetifolia* are described. Mean fibre length of the pulp was 1.08 mm and mean fibre diameter 0.011 mm. Easy-bleaching pulps in good yield could be prepared, and the species is regarded as promising raw material for writing and printing papers.

Guha, S. R. D., Y. K. Sharma, R. Pant, and S. N. Shoundiyal. 1970. Chemical, semi-chemical and mechanical pulps from *Casuarina equisetifolia*. *Indian Forester* 96(11):830-840.

Laboratory-scale experiments indicate that *Casuarina equisetifolia* is a suitable raw material for chemical and semi-chemical pulps. It is unsuitable for mechanical pulps.

Labate, P. J. 1973. [Industrial use of the wood of *Casuarina*.] Folleto Tecnio Forestal No. 32. Instituto Forestal Nacional, Argentina. 38 pp.

Describes some characteristics of wood of *Casuarina cunninghamiana* from three different sources in Argentina and offers information on the suitability of the wood for flooring, packing cases, and veneer. Text in Spanish.

## FORESTS AND FORESTRY FROM THE NATIONAL POINT OF VIEW

Alston, A. S. 1981. Fiji timbers and their uses. (Available from Department of Forestry, P.O. Box 2218, Suva, Fiji.) 6 pp.

Nolan, C. 1980. Loss of a treescape—a report from Bermuda. *Quarterly Journal of Forestry* 74(3):165-176.

An account of the destruction of the natural tree species by scale insects. *Casuarina equisetifolia* was successfully used to replace tree cover quickly, despite insufficient spacing and pruning.

Shatta, H. 1969. Forestry proceeds in Saudi Arabia. *World Wood*, San Francisco 10(9):9,28.

A short, illustrated note on forestry in the less-arid parts of the country. In trials of exotic trees, *Casuarina* spp. have shown promising results.

Williams, L. 1969. Forest and agricultural resources of Dahomey, West Africa. *Economic Botany* 23(4):352-372.

The forest resources of Dahomey have been much reduced by excessive exploitation and clearing for agriculture and are now sufficient for only 75 percent of the country's domestic requirements. *Casuarina equisetifolia* has been found useful as a fuel species.

# Appendix B

## Research Contacts

### ARGENTINA

L. A. Mendonza, Coordinator, Programa Arboles Forestales, Centro de Investigaciones de Recursos Naturales—INTA, 1712 Castelar

### AUSTRALIA

- M. Alexandra, Forester, Australian Seed Co., Box 136, Albert Park, Victoria 3206  
A. Baird, University of Western Australia, Nedlands, W. A. 6009  
B. A. Barlow, Division of Plant Industry, CSIRO, P.O. Box 1600, Canberra, A.C.T. 2601  
D. J. Boland, Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600  
C. J. Borough, Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600  
G. D. Bowen, Division of Soils, CSIRO, Private Bag No. 2, Glen Osmond, S.A. 5064  
M. Chuk, Forestry Unit, Conservation Commission of Northern Territory, P.O. Box 1046, Alice Springs, N.T. 5750  
D. M. Churchill, Director and Government Botanist, Royal Botanic Gardens and National Herbarium, Birdwood Ave., South Yarra, Victoria 3141  
J. Clemens, Department of Agronomy and Horticultural Science, University of Sydney, Sydney, N.S.W. 2006  
Conservator of Forests, Forests Department, P.O. Box 104, Como, W.A. 6152  
P. D. Coyne, National Parks and Wildlife Service, P.O. Box 636, Canberra, A.C.T. 2601  
G. M. Crowley, Post Office, James Cook University, Townsville, Queensland 4811  
P. Davidson, Ornithologist, Airports Division, Department of Aviation, P.O. Box 367, Canberra, A.C.T. 2601  
M. F. C. Day, 12 Melbourne Avenue, Deakin, A.C.T. 2600  
Department of Forestry, G.P.O. Box 944, Brisbane, Queensland 4001  
J. Doran, Division of Forest Research, CSIRO, P.O. 4008, Canberra, A.C.T. 2600  
Forest Commission of Victoria, 1 Treasury Place, Melbourne, Victoria 3002  
A. Fleming, Forestry Department, Australian National University, P.O. Box 4, Canberra, A.C.T. 2600  
A. H. Gibson, Division of Plant Industry, CSIRO, P.O. Box 109, Canberra, A.C.T. 2600  
H. Gray, Country Roads Board, Godenmark St., Kew, Victoria 3101  
D. M. Griffin, Forestry Department, Australian National University, P.O. Box 4, Canberra, A.C.T. 2600  
L. A. S. Johnson, Director, Royal Botanic Gardens, Sydney, N.S.W. 2000  
R. D. Johnston, Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600  
A. C. Lawrie, Department of Applied Biology, R.M.I.T., G.P.O. Box 2476V, Melbourne, Victoria 3001

- J. Leigh, Division of Plant Industry, CSIRO, P.O. Box 1600, Canberra, A.C.T. 2601  
T. B. S. Mahat, Forestry Department, Australian National University, P.O. Box 4, Canberra, A.C.T. 2600  
S. J. Midgley, Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600  
N. J. Moore, Department of Forestry, Australian National University, P.O. Box 4, Canberra, A.C.T. 2600  
H. A. Nix, Division of Land Use Research, CSIRO, P.O. Box 1666, Canberra, A.C.T. 2601  
P. Reddell, Division of Soils, CSIRO, Private Bag No. 2, Glen Osmond, S.A. 5064  
J. Redmond, Appraisals and Evaluation Unit, Australian Development Assistance Bureau, Department of Foreign Affairs, P.O. Box 887, Canberra, A.C.T. 2601  
I. Riley, School of Environmental and Life Sciences, Murdoch University, Murdoch, W.A. 6153  
J. Saunders, Division of Land Use Research, CSIRO, P.O. Box 1666, Canberra, A.C.T. 2601  
K. R. Shepherd, Forestry Department, Australian National University, P.O. Box 4, Canberra, A.C.T. 2600  
W. A. Shipton, Botany Department, James Cook University, Townsville, Queensland 4811  
Tasmanian Forestry Commission, P.O. Box 207B, Hobart, Tasmania 7001  
J. W. Turnbull, Division of Forest Research, CSIRO, P.O. Box 4008, Canberra, A.C.T. 2600  
K. B. Walsh, Department of Botany, University of Queensland, St. Lucia, Queensland 4067  
Wood Technology and Forest Research Division, Forestry Commission of New South Wales, P.O. Box 100, Beecroft, N.S.W. 2119

## BANGLADESH

- M. Omar Ali, Director, Forest Research Institute, Forestdale, P.O. Box 273, Chittagong

## BENIN

- A. G. Agbahungba, Unité de Recherche Forestière, B.P. 884, Cotonou

## CANADA

- O. Sziklai, Faculty of Forestry, University of British Columbia, Vancouver, B.C. V6T 1W5  
L. G. Lessard, Associate Director, Forest Science, IDRC, P.O. Box 8500, Ottawa, K1G 3H9

## CHINA

- Pan Chih-kang, Chinese Academy of Forestry, Wan Shou Shan, Beijing 100091  
Zhu Zhi-Song, Deputy Director, Forest Research Institute, Shahe, Guangzhou, Guangdong

## EGYPT

- Adly Bishay, American University in Cairo, Desert Development Project, 113 Sharia Kasr El-Aini, Cairo
- M. H. El-Lakany, Department of Forestry, Faculty of Agriculture, University of Alexandria, El-Shatby, Alexandria
- M. A. Haikel, Horticulture Research Institute, Cairo University St., Giza, Cairo
- S. A. E. Kandeel, Department of Forestry and Wood Technology, University of Alexandria, El-Shatby, Alexandria
- M. M. Megahed, Department of Forestry, Faculty of Agriculture, University of Alexandria, El-Shatby, Alexandria
- T. A. Omran, Department of Forestry, Faculty of Agriculture, University of Alexandria, El-Shatby, Alexandria
- M. M. Youness, Department of Forestry, Faculty of Agriculture, University of Alexandria, El-Shatby, Alexandria

## FRANCE

Centre Technique Forestier Tropical, 45 bis Avenue de la Belle-Gabrielle, 94130 Nogent-sur-Marne

## INDIA

- M. Stephen Duraivaj, Associate Professor, Department of Agriculture Botany, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu
- R. Jambulingam, Associate Professor of Forestry, Faculty of Horticulture, Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu
- S. Kondas, Managing Director, Tamil Nadu Forest Plantation Corporation Ltd., No. 4, Promenade Road, Cantonment, Tiruchirapalli 620001, Tamil Nadu
- R. S. Mathur, Director, Forestry Research Institute and Colleges, New Forest P.O., Dehra Dun, Uttar Pradesh
- J. S. P. Yadav, Director, Central Soil Salinity Research Institute, Karnal 132001, Haryana

## INDONESIA

Syafi Manan, Senior Lecturer, Faculty of Forestry, Bogor Agricultural University, Bogor

## ISRAEL

- A. Weinstein, Forestry Division, Agricultural Research Organisation, Ilanot, Doar Na Lev Hasharon 42805
- Y. Zohar, Agricultural Research Organisation, Forestry Division, Ilanot, Doar Na Lev Hasharon 42805

## KENYA

- René de Haller, Baobab Farms Ltd., P. O. Box 90202, Mombasa
- P. A. Huxley, ICRAF, P.O. Box 30677, Nairobi
- P. K. Arap Konuche, Silviculturist, Forest Department, P.O. Box 74, Kikuyu

P. K. R. Nair, Senior Scientist, ICRAF, P.O. Box 30677, Nairobi  
K. Oka, IDRC Regional Office, P.O. Box 30677, Nairobi

## MALAYSIA

Director, Forest Research Institute, Kepong, Selangor

## NIGERIA

Forestry Research Institute of Nigeria, P.M.B. 5054, Ibadan

## PAKISTAN

M. Ahmed, Department of Botany, University of Karachi, Karachi-32  
A. H. Chaudhary, Associate Professor, Department of Biological Sciences, Quaid-I-Azam University, Islamabad

## PAPUA NEW GUINEA

A. Allison, Wau Ecology Institute, Box 77, Wau  
A. Bryon, Environmental Science Council, Port Moresby  
Chief Research Officer, Forest Management Research Branch, Office of Forests, Boroko  
K. Thiagalingam, Senior Lecturer, Faculty of Agriculture, P.O. Box 320, University of Papua New Guinea, Port Moresby

## THE PHILIPPINES

R. B. Aspiras, BIOTECH, University of the Philippines at Los Baños, College, Laguna 3720  
S. C. Halos, Chief, Science Research Specialist, Forest Research Institute, College, Laguna 3720  
B. N. Ganguli, Senior Forestry Specialist, Asian Development Bank, P.O. Box 789, Manila 2800

## SAUDI ARABIA

M. L. El-Osta, Department of Plant Production, College of Agriculture, King Saud University, P.O. Box 2460, Riyadh

## SENEGAL

Y. Dommergues, Office de la Recherche Scientifique et Technique Outre Mer, Centre Ostrom de Dakar, B.P. No. 1386, Dakar  
N. A. Moctar, Direction des Eaux et Foret, Parc Forestier de Hann, B.P. 1831, Dakar

## SOUTH AFRICA

R. J. Poynton, South African Forestry Research Institute, P.O. Box 727, Pretoria 0001

## SPAIN

C. Rodriguez-Barrueco, Unit of Nitrogen Fixation, C.S.I.C., Aptdo. 257, Salamanca

## SWEDEN

M. Zumer-Linder, Department of Ecology and Environmental Sciences, Swedish University of Agriculture, 75007 Uppsala

## SRI LANKA

K. Vivekanandan, Research Division, Forest Department, P.O. Box 509, Colombo 2

## TANZANIA

J. E. Maphole, Arid Zone Afforestation Research, P.O. Box 840, Dodoma  
I. M. Shehaghilo, Forest Division, Silvicultural Research Station, P.O. Box 95, Lushoto

## THAILAND

Pravit Chittachumnonk, Division of Silviculture, Royal Forest Department, Bangkok 10900

## TUNISIA

Mohamed Charfi, Institut National de Recherche Forestières, B.P. 2, Ariana

## USA

J. L. Brewbaker, NFTA, Box 680, Waimanalo, Hawaii 96795  
C. B. Davey, Department of Forestry, North Carolina State University, Raleigh, North Carolina 27650  
H. W. Esbenschade, International Tree Crops Institute, USA, P.O. Box 665, Winters, California 95694  
T. F. Geary, USDA Forest Service, P.O. Box 2417, RPE, 606A, Washington, D.C. 20013  
Hawaii Division of Forestry and Wildlife, Department of Land and Natural Resources, 1151 Punchbowl Street, Room 325, Honolulu, Hawaii 96813

- R. K. Hermann, School of Forestry, Oregon State University, Corvallis, Oregon 97331  
W. Jones, Landscape Architecture Program, Bioscience East, University of Arizona,  
Tucson, Arizona 85721  
W. S. Silver, Department of Biology, University of South Florida, Tampa, Florida  
33620 (Florida casuarina)  
R. G. Skolmen, Institute of Pacific Islands Forestry, 1151 Punchbowl Street, Honolulu,  
Hawaii 96813  
J. G. Torrey, Professor of Botany, Cabot Foundation, Harvard University, Petersham,  
Massachusetts 01366  
R. Van den Beldt, Department of Horticulture, University of Hawaii at Manoa, Honolulu,  
Hawaii 96822

## ZIMBABWE

- L. J. Mullin, Forest Research Center, P.O. Box HG 595, Highlands, Harare

## Index of Species

- Agoho, 2, 58  
*Allocauarina*, 2  
*Allocauarina campestris*, 68  
   *decaisneana*, 72  
   *dielsiana*, 74  
   *fraseriana*, 76  
   *huegeliana*, 78  
   *littoralis*, 80  
   *luehmannii*, 82  
   *torulosa*, 88  
   *verticillata*, 86  
*Araucaria cunninghamii*, 19  
   *hunsteinii*, 19  
 Australian pine, 2, 58  
*Azadirachta indica*, 24  
 Belah, 70  
 Black sheoak, 70, 80  
 Bull oak, 82  
*casuarina*, 56, 58  
*Casuarina* spp., 2, 11, 14, 45, 98  
   *angularis*, 32  
   *campestris*, 41, 67, 68-69, 79  
   *collina*, 94  
   *cristata*, 29, 32, 35, 40, 41, 42, 67, 70-71, 102, 104  
   *cunninghamiana*, 3, 8, 10, 22, 23, 24, 25, 26, 29, 31, 32, 35, 36, 40, 41, 44, 45, 49, 50, 52, 53, 55, 56-57, 60, 94, 102, 104, 105  
   *decaisneana*, 6, 10, 25, 32, 40, 41, 45, 67, 72-73, 101, 102  
   *decossata*, 29  
   *deplancheana*, 6, 95  
   *dielsiana*, 41, 67, 74-75  
   *equisetifolia*, 2, 3, 4, 5, 6, 8, 10, 11, 13, 14, 16, 17, 18, 19, 21, 22, 24, 25, 26, 28, 29, 30, 32, 34, 40, 41, 44, 45, 46, 48, 49, 50, 51, 52, 55, 58-59, 60, 62, 100-105  
   *fraseriana*, 41, 45, 67, 76-77, 102  
   *glauca*, 3, 10, 12, 13, 18, 22, 24, 27, 29, 30, 31, 32, 36, 40, 41, 42, 44, 45, 49, 50, 52, 55, 57, 60-61, 70, 84, 94, 100, 102, 104  
   *grandis*, 94, 95, 97  
   *helmsii*, 29  
   *huegeliana*, 32, 41, 67, 78-79  
   *junghuhniana*, 3, 5, 10, 19, 22, 24, 25, 32, 41, 42, 45, 55, 62-63, 94, 102  
   *lepidophloia*, 70, 102  
   *litorea*, 58  
   *littoralis*, 32, 36, 41, 45, 52, 67, 80-81, 88  
   *luehmannii*, 41, 42, 67, 82-83, 102  
   *montana*, 62  
   *nobilis*, 94, 96  
   *nodiflora*, 32  
   *obesa*, 10, 41, 42, 67, 84-85  
   *oligodon*, 10, 18, 19, 32, 41, 45, 47, 55, 64-65, 94  
   *papuana*, 32, 64, 94, 95, 97  
   *rumphiana*, 95, 105  
   *spicigera*, 95  
   *stricta*, 29, 32, 40, 41, 42, 45, 49, 67, 86-87  
   *suberosa*, 80  
   *sumatrana*, 95, 96, 103  
   *tenuissima*, 88  
   *torulosa*, 24, 32, 40, 41, 45, 48, 52, 67, 88-89, 102, 103  
   *Vitiense*, 32  
   Coast sheoak, 86  
*Conocarpus lancifolius*, 24, 102  
 Creek oak, 56  
 Desert sheoak, 72  
 Diels' sheoak, 74  
 Drooping sheoak, 86  
*Eucalyptus* spp., 15, 93  
   *camaldulensis*, 28  
   *tereticornis*, 29  
 Filao, 58  
 Forest sheoak, 88  
*Frankia*, 11, 12, 14, 15, 35-38, 42-43, 93, 95, 100  
 Fraser's sheoak, 76  
 Gago, 18  
*Gmelina arborea*, 4  
 Granite sheoak, 78  
*Gymnostoma nobile*, 94  
   *papuanum*, 94  
   *rumphianum*, 95  
   *spicigerum*, 95  
   *sumatranum*, 95  
*Gynosperma*, 2  
 Hat Yar, 94  
 Horsetail oak, 58  
 Ironwood, 32, 58  
 Jemara, 62  
*Leucaena*, 32  
*Leucaena leucocephala*, 4  
 Mountain oak, 86  
 Neem, 24  
 Nokonoko, 32, 58  
*Pinus caribaea*, 5, 28, 29  
   *occidentalis*, 28  
*Prosopis*, 32  
 River oak, 56  
 River sheoak, 56  
 Rock sheoak, 78  
 Rose sheoak, 88  
 Ru, 58  
 Sheoak, 2, 58, 64  
 Soft yar, 64  
 Swamp sheoak, 60  
 Velau, 32  
 Western Australian swamp sheoak, 84  
 Whistling pine, 58  
 Yar, 64



### **Advisory Committee on Technology Innovation**

HUGH POPENOE, Director, International Programs in Agriculture, University of Florida, Gainesville, Florida, *Chairman*

#### **Members**

WILLIAM BRADLEY, Consultant, New Hope, Pennsylvania

HAROLD DREGNE, Director, International Center for Arid and Semi-Arid Land Studies, Texas Tech University, Lubbock, Texas (member through 1981)

ELMER L. GADEN, JR., Department of Chemical Engineering, University of Virginia, Charlottesville, Virginia

CARL N. HODGES, Director, Environmental Research Laboratory, Tucson, Arizona

CYRUS MCKELL, Native Plants, Inc., Salt Lake City, Utah

FRANÇOIS MERGEN, Pinchot Professor of Forestry, School of Forestry and Environmental Studies, Yale University, New Haven, Connecticut (member through 1982)

DONALD L. PLUCKNETT, Consultative Group on International Agricultural Research, Washington, D.C.

THEODORE SUDIA, Deputy Science Advisor to the Secretary of Interior, Department of Interior, Washington, D.C.

### **Board on Science and Technology for International Development**

GEORGE BUGLIARELLO, President, Polytechnic Institute of New York, Brooklyn, New York, *Chairman*

#### **Members**

SAMUEL P. ASPER, Educational Commission for Foreign Medical Graduates, Philadelphia, Pennsylvania

DAVID BELL, Department of Population Sciences, Harvard School of Public Health, Boston, Massachusetts

LEONARD BERRY, Professor, Graduate School of Geography, Clark University, Worcester, Massachusetts

ERNEST J. BRISKEY, Dean, School of Agriculture, Oregon State University, Corvallis, Oregon

HARRISON S. BROWN, Director, Resources Systems Institute, East-West Center, Honolulu, Hawaii

ROBERT H. BURRIS, Department of Biochemistry, University of Wisconsin, Madison, Wisconsin

**CLAUDIA JEAN CARR**, Associate Professor, Conservation and Resource Studies, University of California at Berkeley, Berkeley, California  
**NATE FIELDS**, Director, Developing Markets, Control Data Corporation, Minneapolis, Minnesota  
**ROLAND J. FUCHS**, Chairman, Department of Geography, University of Hawaii at Manoa, Honolulu, Hawaii  
**ELMER L. GADEN, JR.**, Department of Chemical Engineering, University of Virginia, Charlottesville, Virginia  
**JOHN HOWARD GIBBONS**, Director, U.S. Congress, Office of Technology Assessment, Washington, D.C.  
**N. BRUCE HANNAY**, Foreign Secretary, National Academy of Engineering, Washington, D.C.  
**WILLIAM HUGHES**, Director, Engineering Energy Laboratory, Oklahoma State University, Stillwater, Oklahoma  
**WILLIAM A. W. KREBS**, Vice President, Arthur D. Little, Inc., Acorn Park, Cambridge, Massachusetts  
**GEORGE I. LYTHCOTT**, University of Wisconsin, School of Medicine, Madison, Wisconsin  
**JANICE E. PERLMAN**, Executive Director, Committee for a New New York, New York City Partnership, New York, New York  
**HUGH POPENOE**, Director, International Programs in Agriculture, University of Florida, Gainesville, Florida  
**FREDERICK C. ROBBINS**, President, Institute of Medicine, National Academy of Sciences, Washington, D.C.  
**WALTER A. ROSENBLITH**, Foreign Secretary, National Academy of Sciences, Washington, D.C.  
**FREDERICK SEITZ**, President Emeritus, The Rockefeller University, New York, New York  
**RALPH HERBERT SMUCKLER**, Dean of International Studies and Programs, Michigan State University, East Lansing, Michigan  
**GILBERT F. WHITE**, Institute of Behavioral Science, University of Colorado, Boulder, Colorado  
**BILL C. WRIGHT**, Assistant Dean for International Programs, Oklahoma State University, Stillwater, Oklahoma

**JOHN G. HURLEY**, Director  
**MICHAEL G. C. McDONALD DOW**, Associate Director/Studies  
**MICHAEL P. GREENE**, Associate Director/Research Grants

Board on Science and Technology for International Development  
(JH-217D)  
Office of International Affairs  
National Research Council  
2101 Constitution Avenue, Washington, D.C. 20418, USA

### **How to Order BOSTID Reports**

Reports published by the Board on Science and Technology for International Development are sponsored in most instances by the U.S. Agency for International Development and are intended for free distribution primarily to readers in developing countries. A limited number of copies are available without charge to readers in the United States and other industrialized countries who are affiliated with governmental, educational, or research institutions, and who have professional interest in the subjects treated by the report. Requests should be made on the institution's stationary.

Single copies of published reports listed below are available free from BOSTID at the above address while the supplies last.

### **Energy**

19. **Methane Generation from Human, Animal, and Agricultural Wastes.** 1977. 131 pp. Discusses means by which natural process of anaerobic fermentation can be controlled by man for his benefit and how the methane generated can be used as a fuel.

33. **Alcohol Fuels: Options for Developing Countries.** 1983. 128 pp. Examines the potential for the production and utilization of alcohol fuels in developing countries. Includes information on various tropical crops and their conversion to alcohols through both traditional and novel processes.

36. **Producer Gas: Another Fuel for Motor Transport.** 1983. 112 pp. During World War II Europe and Asia used wood, charcoal, and coal to fuel over a million gasoline and diesel vehicles. However, the technology has since been virtually forgotten. This report reviews producer gas and its modern potential.

38. **Supplement to Energy for Rural Development: Renewable Resources and Alternative Technologies for Developing Countries.** 1981. 240 pp. Updates the 1976 BOSTID publication and offers new material on direct and indirect uses of solar energy. Provides index to both volumes.

39. **Proceedings, International Workshop on Energy Survey Methodologies for Developing Countries.** 1980. 220 pp. Report of a 1980 workshop organized to examine past and ongoing energy survey efforts in developing countries. Includes reports from rural, urban, industry, and transportation working groups, excerpts from 12 background papers, and a directory of energy surveys for developing countries.

## **Technology Options for Developing Countries**

**8. Ferrocement: Applications in Developing Countries.** 1973. 89 pp. Assesses state of the art and cites applications of particular interest to developing countries—boat building, construction, food and water storage facilities, etc.

**14. More Water for Arid Lands: Promising Technologies and Research Opportunities.** 1974. 153 pp. Outlines little-known but promising technologies to supply and conserve water in arid areas. (French language edition is available from BOSTID.)

**21. Making Aquatic Weeds Useful: Some Perspectives for Developing Countries.** 1976. 175 pp. Describes ways to exploit aquatic weeds for grazing, and by harvesting and processing for use as compost, animal feed, pulp, paper, and fuel. Also describes utilization for sewage and industrial wastewater treatment. Examines certain plants with potential for aquaculture.

**28. Microbial Processes: Promising Technologies for Developing Countries.** 1979. 198 pp. Discusses the potential importance of microbiology in developing countries in food and feed, plant nutrition, pest control, fuel and energy, waste treatment and utilization, and health.

**31. Food, Fuel, and Fertilizer for Organic Wastes.** 1981. 150 pp. Examines some of the opportunities for the productive utilization of organic wastes and residues commonly found in the poorer rural areas of the world.

**34. Priorities in Biotechnology Research for International Development: Proceedings of a Workshop.** 1982. 261 pp. Report of a 1982 workshop organized to examine opportunities for biotechnology research in developing countries. Includes general background papers and specific recommendations in six areas: 1) vaccines, 2) animal production, 3) monoclonal antibodies, 4) energy, 5) biological nitrogen fixation, and 6) plant cell and tissue culture.

## **Plants**

**16. Underexploited Tropical Plants with Promising Economic Value.** 1975. 187 pp. Describes 36 little-known tropical plants that, with research, could become important cash and food crops in the future. Includes cereals, roots and tubers, vegetables, fruits, oilseeds, forage plants, and others.

**22. Guayule: An Alternative Source of Natural Rubber.** 1977. 80 pp. Describes a little-known bush that grows wild in deserts of North America and produces a rubber virtually identical with that of the rubber tree. Recommends funding for guayule development.

**25. Tropical Legumes: Resources for the Future.** 1979. 331 pp. Describes plants of the family Leguminosae, including root crops, pulses, fruits, forages, timber and wood products, ornamentals, and others.

**37. The Winged Bean: A High Protein Crop for the Tropics.** (Second Edition). 1981. 59 pp. An update of BOSTID's 1975 report of this neglected tropical legume. Describes current knowledge of winged bean and its promise.

47. **Amaranth: Modern Prospects for an Ancient Crop.** 1983. Before the time of Cortez grain amaranths were staple foods of the Aztec and Inca. Today this extremely nutritious food has a bright future. The report also discusses vegetable amaranths.

#### **Innovations in Tropical Reforestation**

26. **Leucaena: Promising Forage and Tree Crop for the Tropics.** 1977. 118 pp. Describes *Leucaena leucocephala*, a little-known Mexican plant with vigorously growing, bushy types that produce nutritious forage and organic fertilizer as well as tree types that produce timber, firewood, and pulp and paper. The plant is also useful for revegetating hillslopes, providing firebreaks, and for shade and city beautification.

27. **Firewood Crops: Shrub and Tree Species for Energy Production.** 1980. 237 pp. Examines the selection of species suitable for deliberate cultivation as firewood crops in developing countries.

35. **Sowing Forests from the Air.** 1981. 64 pp. Describes experiences with establishing forests by sowing tree seed from aircraft. Suggests testing and development of the techniques for possible use where forest destructions now outpaces reforestation.

40. **Firewood Crops: Shrub and Tree Species for Energy Production.** Volume II. 1983. A continuation of BOSTID report number 27. Describes 27 species of woody plants that seem suitable candidates for fuelwood plantations in developing countries.

41. **Mangium and Other Fast-Growing Acacias for the Humid Tropics.** 1983. 63 pp. Highlights ten acacias species that are native to the tropical rain forest of Australasia. That they could become valuable forestry resources elsewhere is suggested by the exceptional performance of *Acacia mangium* in Malaysia.

42. **Calliandra: A Versatile Small Tree for the Humid Tropics.** 1983. 56 pp. This Latin American shrub is being widely planted by villagers and government agencies in Indonesia to provide firewood, prevent erosion, yield honey, and feed livestock.

43. **Casuarinas: Nitrogen-Fixing Trees for Adverse Sites.** 1983. These robust nitrogen-fixing Australasian trees could become valuable resources for planting on harsh, eroding land to provide fuel and other products. Eighteen species for tropical lowlands and highlands, temperate zones, and semiarid regions are highlighted.

#### **Managing Tropical Animal Resources**

32. **The Water Buffalo: New Prospects for an Underutilized Animal.** 1981. 118 pp. The water buffalo is performing notably well in recent trials in such unexpected places as the United States, Australia, and Brazil. Report discusses the animal's promise, particularly emphasizing its potential for use outside Asia.

44. **Butterfly Farming in Papua New Guinea.** 1983. 36 pp. Indigenous butterflies are being reared in Papua New Guinea villages in a formal government program that both provides a cash income in remote rural areas and contributes to the conservation of wildlife and tropical forests.

45. **Crocodiles as a Resource for the Tropics.** 1983. 60 pp. In most parts of the tropics crocodylian populations are being decimated, but programs in Papua New Guinea and a few other countries demonstrate that, with care, the animals can be raised for profit while the wild populations are being protected.

46. **Little-Known Asian Animals with a Promising Economic Future.** 1983. 124 pp. Describes banteng, madura, mithan, yak, kouprey, babirusa, Javan warty pig and other obscure, but possibly globally useful wild and domesticated animals that are indigenous to Asia.

### **General**

29. **Postharvest Food Losses in Developing Countries.** 1978. 202 pp. Assesses potential and limitations of food-loss reduction efforts; summarizes existing work and information about losses of major food crops and fish; discusses economic and social factors involved; identifies major areas of need; and suggests policy and program options for developing countries and technical assistance agencies.

30. **U.S. Science and Technology for Development: Contributions to the UN Conference.** 1978. 226 pp. Serves the U.S. Department of State as a major background document for the U.S. national paper, 1979 United Nations Conference on Science and Technology for Development.

The following topics are now under study and will be the subjects of future BOSTID reports:

- **Leucaena: Promising Forage and Tree Crop for the Tropics (Second Edition)**
- **Jajoba**

For a complete list of publications, including those that are out of print and available only through NTIS, please write to BOSTID at the address above.







**ORDER FORM**

While the limited supply lasts, a free copy of *Casuarinas: Nitrogen-Fixing Trees for Adverse Sites* will be sent to institutionally affiliated recipients (in government, education, or research) upon written request on your organization's letterhead or by submission of the form below. Please indicate on the labels the names, titles, and addresses, including country, of qualified recipients and their institutions who would be interested in having this report.

Please return this form to:

BOSTID (JH-217D)  
National Research Council  
2101 Constitution Avenue  
Washington, D.C. 20418, USA

20418

43

43

43

43

43

43




## The National Academy of Sciences

The National Academy of Sciences was established in 1863 by Act of Congress as a private, nonprofit, self-governing membership corporation for the furtherance of science and technology, required to advise the federal government upon request within its fields of competence. Under its corporate charter the Academy established the National Research Council in 1916, the National Academy of Engineering in 1964, and the Institute of Medicine in 1970.

## The National Research Council

The National Research Council was established by the National Academy of Sciences in 1916 to associate the broad community of science and technology with the Academy's purposes of furthering knowledge and of advising the federal government. The Council operates in accordance with general policies determined by the Academy under the authority of its congressional charter of 1863, which establishes the Academy as a private, nonprofit, self-governing membership corporation. The Council has become the principal operating agency of both the National Academy of Sciences and the National Academy of Engineering in the conduct of their services to the government, the public, and the scientific and engineering communities. It is administered jointly by both Academies and the Institute of Medicine. The National Academy of Engineering and the Institute of Medicine were established in 1964 and 1970, respectively, under the charter of the National Academy of Sciences.



## The Office of International Affairs

The Office of International Affairs is responsible for many of the international activities of the Academy and the Research Council. Its primary objectives are to enhance U.S. scientific cooperation with other countries; to mobilize the U.S. scientific community for technical assistance to developing nations; and to coordinate international projects throughout the institution.

## The Board on Science and Technology for International Development

The Board on Science and Technology for International Development (BOSTID) of the Office of International Affairs addresses a range of issues arising from the ways in which science and technology in developing countries can stimulate and complement the complex processes of social and economic development. It oversees a broad program of bilateral workshops with scientific organizations in developing countries and conducts special studies. BOSTID's Advisory Committee on Technology Innovation publishes topical reviews of unconventional technical processes and biological resources of potential importance to developing countries.

